



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





600030882R



E. BIBL. RADCL.

7.7.11

~~67~~ 13

19116 e. 38
1

CAROLI LINNAEI,

EQUITIS STELLAE POLARIS, ARCHIATRI REGII, PROF. MED.
ET REI HERB. IN UNIV. UPSAL.

GENERA PLANTARUM.

EDITIO NONA,

CURANTE

CURTIO SPRENGEL,

EQUITE STELLAE POLARIS, AQUILAE RUBRAE ET LEONIS
BELGICI, PROF. MED. ET REI HERB. IN UNIVERS. HALENSI.

TOM. I.

CLASSIS 1—13.

GOTTINGAE,

SUMTIBUS DIETERICHIANIS.

MDCCCXX.



LINNAEI
GENERA PLANTARUM.



CLASSIS I. M O N A N D R I A.

C O N S P E C T U S

CLASSIS PRIMAE

M O N O G Y N A E

Canneae n. 1 — 6.
Scitamineae n. 7 — 16.
Vochisieae n. 17 — 22.
Aliae — 23 — 39.

D I O G Y N A E n. 40 — 51.

T R I G Y N A E n. 52.

P O L Y G Y N A E n. 53, 54.

M O N O G Y N I A.

C A N N E A E.

1. CANNA C. Gesn. (1561.)

Cal. duplex, uterque tripartitus, subcoloratus, superus.
Cor. monopetala, tripartita, irregularis, subringens. *Lab.*
superius tripartitum, bifidum vel nullum.

Stam. Filamentum petaloideum, cuius margini anthera
adnata.

Pistill. petaloideum, cuius margini stigma lineare adnatum.

Caps. infera, membranacea, papilloso-muricata, trilocu-
laris, trivalvis.

Sem. in quovis loculo plura, receptaculo centrali affixa,
globosa. *Albumen* durum, cartilagineum. *Embryo*
axilis, longiusculus, subinvolutus.

Gärtn. t. 12. *Mirbel* in ann. mus. 16. t. 16. *Hegetschweiler*
diss. t. 1.

2. CALATHEA G. F. W. Meyer. (1818.)

Cal. Perianth. commune imbricatum. *Cal.* proprius supé-
rus trisepalus.

Cor. monopetala. Limbus duplex, exterior tripartitus, la-
ciniis lanceolatis, interior bipartitus, laciniis obtusis.

Stam. Filamentum petaloideum, anthera laterali adnata.

Pist. petaloideum. *Stigma* trigonum.

Caps. infera, chartacea, trilocularis, trivalvis.

Sem. in quovis loculo solitaria, subangulata, media basi foveola exsculpta, quae operculo embryonifero clausa est. *Albumen* farinaceum, cavitatibus duabus axilibus instructum, in quarum altera embryo vermicularis.

Bot. reg. 932. 961. praecipue 385. (*Maranta zebrina*.)

3. **MARANTA** Plum. (1703.)

Cal. duplex, uterque tripartitus, superus.

Cor. monopetala, tripartita, resupinata. Lacinia tertia bipartita.

Stam. Filamentum petaloideum bipartitum, hinc antheriferum, illinc stylum incurvum carnosum involvens. Stigma subtrigonum.

Caps. baccata, monosperma. *Sem.* rugosum, durum.

Hegetschweiler diss. t. 4. t. 23—28. *Roscoe* in Linn. transact. 8. t. 20. f. 2. *Redout.* lil. t. 57. Bot. reg. 786.

4. **THALIA** L. (1740.)

Cal. trisepalus, coloratus, superus.

Cor. monopetala sexpartita. Laciniae tres exteriores aequales obtusae: quarta obovata, quinta lateralis unguiculata; sexta lateralis, processibus binis filiformibus.

Stam. Filamentum filiforme laciniae sextae insertum. Anthera lateralis.

Stylus crassus spiralis. Stigma concavum, inflexum, perforatum, subbilabiatum, labio inferiori elongato.

Bacca monosperma. Ad latera embryonis duae cavitates ipsi parallelae.

Rostoe l. c. f. 3.

Plum. ic. ed. Burm. t. 108. f. 1.

Fraser et Sowerby tab. singularis. 1794. [*Peronia Cand.* lil. t. 312.]

5. **PHRYNIUM** Willd. (1797.)

Cal. trisepalus,

Cor. exterior tripetala subaequalis: interioris limbus quadripartitus.

Stam. Filamentum breve sublaterale.

Stylus superne uncinatus, stigmate incurvo subinfundibuliformi.

Caps. trilocularis, trivalvis. *Sem.* solitaria, basi arillata.

Embryo uncinatus.

Roscoe scitama. t. 2. *Asiat. research.* 11. t. 3.

6. **MYROSMA** L. fil. (1781.)

Cal. superus, duplex: uterque tripartitus.

Cor. monopetala, inaequalis, quinquepartita: laciniiis duabus superioribus brevioribus, tribus inferioribus longioribus apice trifidis.

Stam. Filamentum margini laciniae mediae inferioris adnatum, subulatum. Anthera ovata, compressa.

Stylus crassus, trigonus, deflexus, longitudinaliter fissus.

Stigma bilabiatum.

Caps. trilocularis, trivalvis, polysperma. *Sem.* angulata.

Roscoe in Linn. transact. 8. t. 20. f. 4.

SCITAMINEAE.

Familia, tum *Orchideis* tum *Irideis* *Haemodoreis*que affinis, intra tropicos tantum obvia, aromatum ferax. *Caulis* e radice tuberosa surgit, e vaginis foliorum involutis ortus; vasis spirales apertis, membranaceis insignis; saepe abbreviatus aut nullus. *Folia* simplicia, costata. *Flores* speciosi bracteati. *Perianthium* duplex tripartitum. *Filamentum* sulco stylum amplexans. *Anthera* biloba, lobis bilocularibus. *Stamina* duo sterilia basin styli stipantis nectarifera. *Capsula* infera trilocularis polysperma. *Semina* subarillata. *Albumen* farinaceum. *Vitellus* embryonem ambiens, aroma spirans.

7. HEDYCHIS König. (1783.)

Cal. tubulosus.

Cor. monopetala, tubulosa, limbo sexpartito, resupinato, lobo medio seu labello fissio.

Stam. Filamentum filiforme articulaturn. *Anthera* terminalis.

Stylus filiformis. *Stigma* infundibuliforme.

Caps. familiae.

Roscoe scitam. t. 3. 4. etc. Bot. reg. 526. 774. 1022.

8. ROSCOEA Sm. (1804.)

Cal. tubulosus.

Cor. duplex, exterior tripartita irregularis, lacinia superiore fornicata erecta, inferiore bilabiata.

Anthera incurva, appendice ad basin bicalcarata.

Stigma globulare, perforatum.

Smith. exot. bot. t. 100. *Hooker* exot. flor. 2. t. 144.

9. KIMPFERIA L. (1737.)

Cal. tubulosus, fissus.

Cor. duplex, exterior tripartita, inferiore sublongior, lacinii angustis; interior triloba, lobo intermedio labelliformi.

Stam. Filamentum carinatum, basi vel medio antheram bilocularem gerens, apice in laminam bipartitam expansum.

Stylus filiformis, a filamento receptus, stigmato urceolato ciliato.

Caps. familiae.

Roscoe in Linn. transact. 8. t. 20. f. 40. *Fischer* in Comment. soc. mosqu. t. 1. B. t. 2. C. *Redout.* liliac. t. 49. 144. 389. Bot. reg. t. 173.

10. CURCUMA Val. Cord. (1561.)

Cal. tubulosus, tridentatus.

Cor. duplex, exterior tripartita, interior triloba, lobo medio labelliformi.

Stam. Filamentum carinatum, petaloideum, apice trilobum, lobo medio antherifero. Anthera bicalcarata, *Stylus* filiformis, stigmate capitato.

Caps. familiae. *Sem.* corticata.

Roscoe scitam. t. 15. 23. 24. *Linn.* transact. l. c. f. 12.

11. *AMOMUM* L. (1737.)

Cal. tubulosus, apice trilobus.

Cor. duplex, exterior tripartita subaequalis: interior unilabiata, expansa.

Stam. Filamentum apice lateribusque in lacinulas duas acutas abiens, lobulo apicis medio fisso. Anthera inter lacinulas terminales et laterales adnata.

Stylus filiformis ab anthera receptus, stigmate infundibuliformi.

Caps. familiae.

Sims. in ann. of bot. 1. t. 13. *Bot. reg.* t. 329.

[*Hornstedtia Retz.*]

12. *ZINGIBER* Gärtn. (1788.)

Cal. tubulosus, hinc fissus.

Cor. duplex, exterior tripartita, subconnivens, interior unilabiata triloba.

Stamen. Filamentum basi planum, sursum involutum, marginibus incrassatis antheram fovens, apice indiviso, corniculato.

Stylus filiformis, stigmate infundibuliformi, filamento obvolutus.

Caps. familiae. *Sem.* in pulpa nidulantia.

Gärtn. de fruct. t. 12. *Fischer*, in comment. soc. mosqu. t. 2.

D. t. 4, F. *Roscoe* in *Linn.* transact. 8. t. 20. f. 8. scitam. t. 22.

[*Amomum* L. *Lam.*]

13. *COSTUS* L. (1737.)

Cal. tubulosus, trifidus.

Cor. duplex, exterior tripartita subconnivens, interior unilabiata, campanulata, postice fissa, lobis nullis lateralibus.

Stam. Filamentum oblongum, planum, petaloideum. Anthera medio adnata.

Stylus filiformis, fauci corollae insertus, antheram transiens, stigmate bilamellato, basi corniculis binis instructo.

Caps. familiae.

(Vagina petiolaris supra folii insertionem ochreae speciem format R. Br.)

Jacqu. ic. rar. 1. t. 1. *Bot. reg.* t. 665. *Roscoe* scitam. t. 32. in *Linn.* transact. 8. t. 20. f. 9.

14. *HELLENIA* Willd. (1797.)

Cal. tubulosus, hinc fissus.

- Cor.* duplex, exterior tripartita, patens; interior unilabiata, basi utrinque denticulo aucta.
Stam. Filamentum lineare, ultra antheram marginalem productum, lobulo brevissimo rotundato v. bilobo, *Stylus* filiformis, stigmate capitato.
Caps. familiae, crustacea.
Roscoe scitam. t. 30. Bot. reg. t. 1037.
 [Languas König. Heriteria Retz.]
15. *ALPINIA* Plum. (1703.)
Cal. tubulosus, tridentatus.
Cor. duplex, exterior tripartita, subconnivens; interior unilabiata, basi utrinque denticulo aucta.
Stam. Filamentum lineare, ultra antheram non productum. Anthera terminalis, stylum recipiens.
Stylus filiformis, antheram superans, stigmate concavo.
Caps. familiae, baccata.
 Bot. reg. 141. 328. 777. *Roscoe* scitam. t. 21. 54.
 [Gethyra Salisb. Eleteria White. Remeaulia Andr.]
16. *GLOBBA* L. (1767.)
Cal. subtubulosus, trifidus.
Cor. duplex, exterior tripartita; interior unilabiata, basi denticulis binis.
Stam. Filamentum lineare, ultra antheram non productum. Anthera basi vel nuda, vel appendiculata.
Stylus filiformis, stigmate concavo.
Caps. familiae.
 Smith exot. bot. 103. 117. Roxb. corom. 3. t. 228 — 230.
Roscoe scitam. t. 16. Linn. transact. 8. t. 20. f. 13.
 [Mantisia Curt. Ceratanthera Hornem.]

V O C H I S I E A E *Hilar.*

- Arbores aut frutices, resina scatentes, Americam australem tropicam inhabitantes, ramis foliisque subcoriaceis oppositis vel verticillatis: floribus vel axillaribus solitariis, vel terminalibus racemosis aut paniculatis. Calyx inferus, coloratus, subcalcaratus. Cor. 1—3-petala. Stamina quandoque sterilia. Stylus simplex, stigmate capitato. Caps. lignosa, trilocularis, trivalvis, polysperma. Semina axi centrali affixa. Albumen nullum. Embryo cotyledonibus foliaceis.*
17. *CUCULLARIA* Schreb. (1789.)
Cal. liber, quinquepartitus, lacinia summa maxima cucullata, basi calcarata.
Cor. tripetala.
Stam. fertile unicum, petalo medio oppositum. Anthera quadrilocularis. Filamenta duo sterilia lateraliter.
Stylus simplex, curvatus, stigmate capitato.
Caps. familiae. Semina solitaria apice alata.
 Mart. nov. gen. 1. t. 83 — 92.
 [Vochy Aubl. Vochysia Juss. Mart.]

18. *SALVERTIA* Hilar. (1820.)

Cal. liber, quinquepartitus, laciniis subaequalibus, suprema calcarata.

Cor. pentapetala subregularis.

Stam. fertile unicum, petalo infimo oppositum. Filamenta duo sterilia lateralia.

Stylus clavatus, stigmate elliptico.

Caps. prioris.

Mart. nov. gen. 1. t. 93.

19. *DITMARIA* * (1818.)

Cal. germini adnatus, quinquepartitus, irregularis, laciniis summa maxima basi calcarata.

Cor. petalum unicum, obcordatum.

Stam. unicum fertile, anthera quadriloculari dorso affixa. Filamenta sterilia terna lateralia.

Stylus lateralis filiformis.

Caps. ignota.

Mart. nov. gen. t. 82.

[*Erisma* Rudg. Mart.]

20. *QUALEA* Aubl. (1775.)

Cal. liber, subpentasepalus, summo maximo plerumque calcarato.

Cor. petalum unicum obcordatum.

Stam. fertile unicum, raro duo; filamentorum sterilius rudimentis obsoletis. Anthera quadrilocularis.

Stylus simplex, stigmate capitato.

Caps. lignosa, trilocularis, trivalvis. Placenta columnaris centralis libera. *Semina* in singulis loculis numerosa, apice alata.

Mart. nov. gen. t. 78 — 81.

21. *AMPHILOCHIA* Mart. (1824.)

Cal. liber, pentasepalus, sepalo summo maximo subcalcarato.

Cor. petalum unicum, obcordatum.

Stam. fertile unicum, anthera quadriloculari. Filamenta sterilia nulla.

Stylus simplex, stigmate capitato.

Caps. lignosa, trilocularis, trivalvis. *Sem.* unicum vel duo in quovis loculo.

Mart. nov. gen. t. 77.

[*Agardhia* * syst.]

22. *CALLISTHENIA* * (*Callisthene* Mart. 1824.)

Cal. liber, pentasepalus, sepalo supremo maximo calcarato.

Cor. petalum unicum, obcordatum.

Stam. fertile unicum, anthera quadriloculari. Filamenta sterilia nulla.

Stylus simplex, stigmate capitato.

Caps. lignosa, trilocularis, trivalvis, valvis absque dissepimento, loculis mono-dispernis.

Mart. nov. gen. t. 75. 76.

23. *MANGIFERA* L. (1748.)

Cal. quinquepartitus.

Cor. petala quinque lanceolata, calyce maiora.

Stam. unicum fertile: filamentis binis aut quaternis sterilibus.

Stylus filiformis, stigmatē capitato.

Drupa monosperma, putamine fibrillis lignosis. Sem. exalbuminosum, cotyledonibus crassis, maximis.

Terebinthaceae. Lam. illustr. t. 138. Gärtn. t. 100.

24. *USTERIA* Willd. (1792.)

Cal. inferus, quadridentatus, dente quarto maximo.

Cor. hypocrateriformis quadripartita.

Caps. oblonga, bilocularis: placenta laterali carnosā. Sem. numerosa, imbricata, margine alato. Albumen tenue.

Embryo cotyledonibus subrotundis, foliaceis.

— — *Willd. in Beobacht. der berl. Naturf. B. 4. t. 2. Sims. in ann. of bot. 1. t. 7.*

[*Monodynamis* J. F. Gmelin.]

25. *LOPEZIA* Cav. (1791.)

Cal. superus, tetrasepalus, caducus.

Cor. pentapetala irregularis. Petala duo lateralia spatulata, inferiori cucullato complicato genitalia impubia occultente; duo superiora basi biglandulosa.

Caps. subrotunda, quadrilocularis, polysperma.

Onagraea Jacq. ic. rar. t. 203. ecl. 2. t. 109. 110. Andrews repos. 551.

[*Pissura* Bonat.]

26. *CENTRANTHUS* Neck. Cand. (1805.)

Cal. superus, limbo integerrimo involuto.

Cor. quinqueloba, calcarata.

Stam. longitudine corollae, anthera subrotunda.

Stylus filiformis, longitudine staminis.

Achenium inferum, pappo plumoso coronatum.

Valerianaeae. Engl. bot. 1531. (Valeriana).

27. *PHILXDRUM* Banks. Gärtn. (1788.)

Spatha monophylla, perianthio longior. Perianthium tetrapetalum, coloratum, marcescens, petalis duobus angustioribus, in quae stamina sterilia transiisse videntur.

Stam. Filamentum subulatum, anthera biloba.

Stylus simplex, stigmatē capitato.

Caps. supera, trilocularis, trivalvis, valvis medio septiferis. Sem. numerosa, minuta. Embryo filiformis rectus in albumine peripherico.

Affine Juncos. Gärtn. t. 16. Roscoe in Linn. transact. 8. t. 20. f. 5.

[*Garciana* Lour.]

28. POLLICHIA Ait. (1789.)*Cal.* quinquedentatus, inferus.*Cor.* nulla.*Stam.* Filamentum filiforme. Anthera didyma subrotunda.*Stylus* filiformis, stigmate bifido.*Sem.* unicum, fundo calycis incrassati inclusum, squama receptaculi carnosae suffultum. Albumen carnosum.*Embryo* subperiphericus.*Paronychieae*, Smith, spicil. bot. 1. t. 1. Gärtn. t. 213.**29. CERDIA Sess. Cand. (1828.)***Cal.* quinquepartitus, intus corollinus, laciniis apice setigeris.*Cor.* nulla.*Stylus* filiformis apice bifidus.*Caps.* unilocularis polysperma.*Paronychieae*.**30. SALICORNIA L. (1737.)***Cal.* integer, squamatus, tumidus, spicae ita innidulans, ut post anthesin foveolae supersint.*Cor.* 0.*Stam.* filamentum unicum (aut duo), anthera subrotunda biloba.*Stylus* simplex, apice bi-vel trifidus.*Semen* calyci immersum, embryo peripherico.*Chenopodeae*, Engl. bot. 415. 1691.**31. HALOCNEMON Marsch. Bieb. (1819.)***Cal.* communis, amenti squama truncata: proprius e squamis ternis appressis: post anthesin nuda superest rhachis filiformis.*Cor.* 0.*Stam.* filamentum unicum, anthera subrotunda biloba.*Stylus* simplex, apice bi-vel trifidus.*Semen* calyce persistente tectum, embryo peripherico.*Chenopodeae*, Pallas illustr. t. 2 — 7. Gärtn. t. 127. (Salicornia).**32. HIPFURIS L. (1737.)***Perianthium* subnullum.*Stam.* Filamentum simplex, germi insidens, anthera subrotunda, biloba.*Stylus* filiformis, simplex.*Achenium* albumine carnoso, embryo axili filiformi.*Naiades*, Sturm. fl. germ. Reichenbach. icon. 1. t. 86. Gärtn. t. 87.**33. APHELIA R. Br. (1810.)***Squamae* spicae distichae uniflorae.*Cor.* univalvis.*Stam.* Filamentum cum anthera simplici.*Stylus* simplex, stigmate simplici.

Utriculus monospermus, line longitudinaliter dehiscens.

Restiæ R. Br. Ann. scienc. nat. 13. t. 2. f. 1.

14. CHONDRACHNE R. Br. (1810.)

Spica undique arcte imbricata, squamis cartilagineis.

Spicula sub singula squama multiflora androgyna, paleis fasciculatis, exterioribus masculis.

Stylus centralis bifidus.

Cyperoidææ.

35. CHORIZANDRA R. Br. (1810.)

Capitulum sessile e spiculis numerosis aggregata.

Spiculae nudae, multiflorae, squamulis fasciculatis.

Stamina singula intra singulas squamas periphericas.

Stylus bifidus, centralis.

Cyperoidææ.

36. ASCARINA Forst. (1775.)

Amenta filiformia, dioeca.

Perianthium, squama brevissima.

Stam. Filamentum abbreviatum. *Anthera* oblonga, quadrisulca.

Stylus nullus. *Stigma* subtrilobum germi adnatum.

Caryopsis.

Chloranthææ R. Br. Forst. charact. gen. t. 59.

37. TRICHOCLADUS Pers. (1807.)

Amenta subcapitata, polygama.

Perianth. squamae brevissimae.

Stam. Filamentum filiforme, anthera oblonga.

Stylus subclavatus.

Caps. unilocularis, quadrivalvis, monosperma.

— — [Dahlia Thaub.]

38. HYPOLEPIS Pers. (1807.)

Flores bracteati dioeci.

Perianthium monopetalum, sexpartitum, intus barbatum.

Stam. Filamentum crassum, anthera clavata.

Stylus brevis, stigmate capitato.

Caps. supera, sexlocularis, polysperma.

— — [Phelypaea Thurb.]

39. CYNOMORIUM Michel. (1729.)

Amentum crassum, squamis peltatis crassis sparsis et paleis rectis angustis dense obsessum.

Stam. e paleis connatis et peltula centrali. Filamentum filiforme, anthera subrotunda, biloba, lobis bilocularibus.

Styli inter stamina sparsi, e paleis pariter basi connatis et peltulis centralibus surgentes, stigmatibus capitatis.

Nucula monosperma, paleis coronata: albuminosa, embryone laterali, globuloso.

Balanophorææ Rich. Michel. nov. gen. t. 12. Richard mém. du mus. 8. t. 21.

DIGYNIA.

40. *Aspicarpa* Lag. (1814.)*Perianthium* quinquepartitum, tribracteatum.*Stamen* brevissimum, hypogynum; anthera subcordata, biloculari.*Styli* brevissimi; stigmatibus truncatis, bilobulatis.*Nucula* monosperma.*Malpighiaceae* Rich. Rich. in mēm. du mus. vol. 2. t. 13.41. *Ceratocarpus* Buxbaum. (1726.)*Flores* monoeci. *Perianthium* monophyllum cuneatum, utrinque carinatum, bicornue.*Stam.* Filamentum capillare, receptaculo insertum. Anthera ovalis, didyma.*Styli* capillares, stigmatibus simplicibus.*Semen* compressum, calyce tectum. Embryo periphericus.*Chenopodeae.* Gleditsch in nov. act. petrop. 16. t. 17.

Gärtn. t. 127.

42. *Corispermum* Ant. Jussieu. (1712.)*Cal.* disepalus,*Cor.* 0.*Filamentum* simplex, anthera ovali.*Styli* capillares, stigmatibus acutis.*Caryopsis* compressa, embryo peripherico.*Chenopodeae.* Gärtn. t. 75.43. *Mniarum* Forst. (1775.)*Cal.* quadrifidus, urceolatus, persistens.*Cor.* 0.*Stam.* Filamentum rectum; calycem excedens; anthera didyma.*Styli* capillares, stigmatibus acutis.*Utriculus* monospermus, calyce cinctus. Embryo periphericus.*Chenopodeae.* Forst. charact. gen. t. 1. Labillard. nov. holl. t. 2. [Ditoca Banks. Gärtn. t. 126.]44. *Blitum* L. (1737.)*Cal.* trifidus, post anthesin in baccam turgens.*Cor.* 0.*Stam.* Filamentum setaceum, anthera didyma.*Styli* capillares.*Semen* crustaceum in calyce baccato occultum. Embryo periphericus.*Chenopodeae.* Sturm. fl. germ. Gärtn. t. 126. Poiteau et Turpin flor. paris. t. 2. 3.45. *Mobelia* Lour. (1790.)*Flores* androgyni, amentacci.*Perianth.* Squamæ trigonæ subcaruosæ.*Cal.* et *Cor.* 0.

Stam. Filam. 0. Anthera subovata ad basia squamæ.

Styl. 0. Stigmata duo subulata, patentia.

Drupa nuce subovata lacunosa, biloculari.

6. *BROSIMUM* Sw. (1788.)

Flores dioeci, amentacei.

Amentum globosum, undique tectum squamis peltatis, maioribus basilaribus.

Cal. et Cor. 0.

Stam. Filamenta solitaria, brevissima, inter squamas.

Anthera orbicularis, peltata, bilamellata, polline globuloso inter lamellas erumpente.

Styli capillares, basi subcoherentes.

Bacca globulosa, monosperma, nucleo bilobo.

Urticæ. Swartz fl. ind. occ. t. 1.

47. *CALLITRICHE* Fab. Column. (1616.)

Cal. 0.

Cor. dipetala, petalis incurvis, canaliculatis.

Stam. Filamentum simplex subrecurvum: anthera didyma.

Styli capillares.

Caps. quaternæ, hinc alatae, monospermae. Embryo inversus, dicotyledoneus in albumine.

Haloragææ. Engl. bot. 722. Sturm. flor. germ.

48. *ZOSTERA* L. (1747.)

Flores androgyni.

Spadix planus in vaginis foliaceis, genitalibus sparsis obtectus.

Stam. Filamentum brevissimum. Anthera cylindrica unilocularis, polline fibrillis ramulosis mixto.

Stigmata duo capillaria.

Nucula striata. Embryo dicotyledoneus, incurvus.

Naiades. Engl. bot. 467. Hook. flor. londin. t. 35.

49. *CAULINIA* Willd. (1798.)

Flores monoeci.

Perianthium nullum, nisi vaginulae brevissimæ.

Stam. Anthera sessilis, apice denticulata.

Styli capillares.

Caryopsis.

Naiades. Michel. nov. gen. t. 8. Schrader Handh. 3. t. 296.

Naias muricata Delil. t. 50., quæ forte huc pertinet, stylos habet ternos.

50. *PSILURUS* Trin. (1820.)

Spiculæ unifloræ (subbifloræ) in spica articulata.

Cal. uni-bivalvis: valvæ laterales, corolla breviores.

Cor. bivalvis, valva exteriore apice subaristata.

Styli penicillares.

Gramineæ. Pal. Beauv. agrost. t. 20. f. 10. Host. gram. 4. t. 29.

[*Monerma* Pal. Beauv. *Asperella* Host.]

Bacca didyma, lobo altero saepe abortienta. Embryo erectus, cotyledonibus ovatis.

Bot. reg. 1. Gärtn. t. 42.

56. *NYCTANTHES* L. (1737.)

Flores aggregati, involucrati.

Cal. tubulosus, integer.

Cor. tubulosa, limbo plano, oblique torto, quinquelobo.

Stam. Antherae subsessiles intra tubum.

Stylus simplex, stigmate capitato.

Caps. duae, coniunctae, compressae, monospermae. Embryo, qualis Jasmini.

Bot. reg. 399. [Parilium Gärtn. t. 51. Scabrita Gärtn. t. 138.]

57. *LIGUSTRUM* Virgil.

Cal. tubulosus, abbreviatus, quadridentatus.

Cor. infundibuliformis, tubo calycem superante, limbo quadripartito.

Stam. Filamenta brevia, tubo corollae inserta. Antherae biloculares.

Stylus brevissimus, stigmate bifido obtuso.

Bacca globosa, bilocularis, loculis mono-dispermis.

Albumen duriusculum. Embryo inversus.

Engl. bot. 764. Gärtn. t. 92.

58. *PHILLYREA* Diosc.

Cal. tubulosus, abbreviatus, quadridentatus.

Cor. breviter campanulata, quadrifida.

Stam. subexserta, filamentis brevibus, antheris erectis.

Stylus simplex, stigmate crassiusculo.

Bacca globosa, bilocularis, loculo altero saepe abortiente.

Sem. unicum, albumine subfarinaceo, embryone inverso.

Tournef. t. 367. Gärtn. t. 92.

59. *OLEA* Varr. Cat.

Cal. tubulosus, abbreviatus, quadridentatus.

Cor. breviter campanulata, limbo quadripartito, plano.

Stam. subexserta, filamentis brevibus, antheris erectis.

Stylus simplex, brevissimus: stigma bifidum, lacini emarginatis.

Drupa bilocularis, loculo altero abortiente.

Nux oblonga; albumine subcarneo, embryone inverso.

Tournef. t. 370. Gärtn. t. 93.

60. *CHIONANTHUS* L. (1737.)

Cal. quadripartitus, persistens.

Cor. tubo brevissimo, limbi laciniis longissimis linearibus.

Stam. Filamenta brevissima intra tubum: antherae datae, erectae.

Stylus brevissimus, stigmate trifido.

Drupa monosperma, putamine striato.

Lam. ill. t. 9. Gärtn. t. 39.

51. SYRINGA Dodon. (1557.)

Cal. tubulosus, abbreviatus, quadridentatus, persistens.

Cor. infundibuliformis, tubo elongato, limbo quadripartito.

Stam. filamentis brevissimis, antheris ovalibus intra tubum.

Stylus filiformis, stigmate crassiusculo, bitido.

Caps. oblonga, bilocularis, bivalvis, valvis dissepimento contrariis. *Semina* solitaria, oblonga, compressa, margine membranacea. *Embryo* inversus.

Sturm. fl. germ. *Gärtn.* t. 49.

62. FORSYTHIA Vahl. (1804.)

Cal. quadripartitus.

Cor. subcampanulata, quadrifida.

Stam. filamentis brevissimis, antheris oblongis, calycem aequantibus.

Stylus longitudine calycis, stigmate bilobo.

Capsula?

[*Syringa Thunb.* fl. jap. t. 3.]

63. BOLIVARIA Schlechtend. (1826.)

Cal. quinquepartitus, persistens.

Cor. campanulata, quinquepartita, lobis acutiusculis.

Stam. corollam aequantia, filamentis tubo insertis, antheris medio affixis, oblongis, bilocularibus.

Stylus simplex, stigmate subcapitato.

Caps. cartilaginea, didyna, circumscissa, bilocularis, loculis dispermis. *Sem.* angulata reticulata. *Embryo* rectus.

Linnaea, 1. tab. 4. f. 1.

L A B I A T A E. (Cf. cl. XIV.)

64. SALVIA Plin.

Cal. tubulosus; vel trilobus ampliatus, vel tridentatus subcylindricus; vel tubulosus, ore bilabiato, labio superiori tri-inferiori bidentato.

Cor. tubulosa, ringens. Labium superius galeatum, inferius trifidum, lobo medio maiori.

Stam. ascendentia (raro declinata): filamentis tubo corollae insertis appendiculatis, seu infra medium processibus glanduliferis, antheris oblongis.

Stylus filiformis, stigmate bitido, acuto. Germina nectario (gynobasi) suffulta, styli basin circumstantia.

Caryopses quaternae in fundo calycis, seminibus exalbuminosis, embryo recto.

Sturm. fl. germ. *Mirbel.* in ann. mus. 15. t. 14. f. 18.

Gärtn. t. 66.

65. ROSMARINUS Columell.

Cal. tubulosus, ore bilabiato, labio superiori integro, inferiori bifido.

Cor. ringens, labio superiori bipartito, inferiori trifido.

Stam. corolla longiora, filamentis simplicibus.

Stylus simplex, stigmate acuto.

Caryopses ut in *Salvia*.

Lamarck ill. t. 19.

66. *MONARDA* L. (1737.)

Cal. tubulosus, striatus, ore quinqueidentato.

Cor. tubulosa, ringens, labio superiori angusto stamina includente, inferiori trilobo.

Stam. filamentis ascendentibus, intra labium superius, antheris incumbentibus.

Stylus simplex, stigmate bifido acuto.

Caryopses ut in *Salvia*.

Lam. ill. t. 49. Bot. reg. 87.

67. *ZIZIPHORA* L. (1736.)

Cal. cylindricus, striatus; ore minimo, quinqueidentato, fauce barbata.

Cor. ringens, tubo calycem aequante, limbo minori. *Lab.* *sup.* ovatum, integrum, reflexum: *infer.* trifidum.

Stam. filamentis simplicibus, antheris oblongis.

Stylus subulatus, stigmate acuto.

Caryopses ut in *Salvia*.

Lam. ill. t. 18. *Gärtn.* t. 66.

68. *CUNILA* L. (1764.)

Cal. cylindricus, striatus, ore quinqueidentato villosus.

Cor. ringens. *Lab.* *sup.* planum, acutiusculum emarginatum: *infer.* trilobum, lobo medio emarginato.

Stam. filamenta quatuor, quorum duo sterilia. *Antherae* didymae.

Stylus filiformis, stigmate bifido.

Caryopses ut in *Salvia*.

69. *LYCOPUS* L. (1737.)

Cal. tubulosus, quinquefidus.

Cor. subtubulosa, quadriloba, lobo superiori latiori emarginato.

Stam. filamentis simplicibus, antheris erectis bilocularibus.

Stylus simplex, stigmate bifido.

Caryopses ut in *Salvia*.

Schkuhr. Handb. t. 4. Engl. bot. 1105.

70. *AMETHYSTEA* L. (1748.) *Amethystina* Amman. (1739.)

Cal. campanulatus, semiquinquefidus.

Cor. tubulosa, limbo quinquepartito, subinaequali. *Lab.* *sup.* bipartitum: *inf.* tripartitum, lacinia media maiore integerrima.

Stam. exserta, filamentis simplicibus, antheris didymis.

Stylus curvatus, stigmate bifido.

Caryopses ut in *Salvia*.

Lam. ill. t. 16.

71. *STACHYTARPHETA* Vahl (1804.)*Cal.* tubulosus, quadridentatus.*Cor.* hypocrateriformis, tubo incurvo, limbo quinquelobo.*Stam.* quatuor, quorum duo sterilia.*Caryopses* duae, immaturae in drupae formam iunctae.*Reichenb. hort.* 1. t. 59. *Jacqu. ecl.* 2. t. 159. 160. (*Cymburus Salisb. parad.* 49. 53. *Verbena Jacqu. ic. rar.* 2. t. 208.)72. *HOSLUNDIA* Vahl (1804.)*Cal.* tubulosus, quinquedentatus.*Cor.* ringens, labio superiori concavo, infer. trifido.*Stam.* quatuor, quorum duo sterilia.*Semina* quatuor intra calycem baccatum.*Polis. Beauv. fl. ovar.* t. 33.73. *COLLINSONIA* L. (1737.)*Cal.* campanulatus, quinquefidus, subbilabiatus; lab. sup. tri-infer. bifido.*Cor.* tubulosa, limbo quinquefido irregulari, laciniis superioribus brevissimis, infima elongata, multifida fimbriata.*Stam.* longissima, filamentis simplicibus, antheris incurventibus.*Stylus* setaceus, stigmate bifido acuto.*Utriculi* duo globosi.*Lam. ill.* t. 21. *Schk. Handb.* t. 2. b.74. *MORINA* Tournef. (1717.)*Cal.* duplex: *exter.* inferus, tubulosus persistens, multidentatus, dentibus spinosis inaequalibus: *inter.* superus, bipartitus, laciniis obtusis persistentibus.*Cor.* ringens, tubo longissimo, inferne angustissimo, superne ampliato: labio superiori semibifido, inferiori trifido.*Stam.* filamentis simplicibus, antheris cordatis erectis.*Stylus* filiformis, staminibus longior. Stigma capitato-peltatum.*Achenium*, calyce interno coronatum.Desciscit a familia, ut maiore iure ad *Dipsaceas* Juss. pertinere videatur.*Lam. ill.* t. 21.

V E R B E N E A E. Cf. cl. XIV.

75. *MELASANTHUS* Pohl. (1827.)*Cal.* tubulosus, subplicatus, breviter quinquedentatus.*Cor.* tubulosa, subinfundibuliformis, limbo plano, irregulariter quinquelobo.*Stam.* quatuor, quorum duo sterilia.*Caps.* elliptica, bilocularis, disperma.*Pohl. ic. bras.* t. 60—65.

76. CORNUTIA Plum. (1703.)

Cal. parvus quinquedentatus, persistens.

Cor. tubulosa, tubo elongato, limbo quadrifido inaequ-

Stam. quatuor, quorum duo sterilia.

Stylus bifidus, stigmatibus crassiusculis.

Drupa, putamine quadriloculari. Semina solitaria, ebuminosa. Embryo erectus.

Lam. ill. t. 541. Hosta Jacqu. hort. schönbrunn. 1. t.

Gärtn. t. 213.

PERSONATAE. Cf. cl. XIV.

77. VERONICA Fuchs. (1542.)

Cal. quadri-quinquepartitus.

Cor. subrotata, quadriloba, lobo quarto angustiori: (rasime tubulosa aut subringens).

Stamina duo fauci inserta.

Stylus filiformis, stigmate capitato.

Caps. bilocularis, obcordata, polysperma, dissepime in verticales sibi impositas placentas abeunte, aut centa demum libera. Embryo in albumine erectus.

Sturm. fl. germ. *Gärtn.* t. 54.

78. PAEDEROTA L. (1764.)

Cal. quinquepartitus, aequalis.

Cor. tubulosa, limbo suberecto bilabiato, labio super. dentato, inferiori integro aut bidentato.

Stamina ascendentia.

Caps. compressa, bilocularis, bivalvis; placenta den libera. *Sem.* numerosa, minuta.

Sturm. fl. germ.

79. WULFFENIA Jacqu. (1781.)

Cal. quinquepartitus, persistens.

Cor. tubulosa, basi subgibba, limbo ad faucem bart bilabiato; labio super. integro stamina involvente, in trilobo.

Stamina ascendentia, antheris bilocularibus.

Stylus longissimus, stigmate capitato.

Caps. compressa, bilocularis, quadrialvis, placenta der libera.

Jacqu. miscell. austr. 2. t. 8. f. 1. ic. rar. I. t. 2.

80. GRATIOLA Dodon. (1557.)

Cal. quinquepartitus, bracteatus vel basi nudus.

Cor. resupinata, tubo calycem excedente, limbo quadrifido, lobo superiori emarginato latiori, reliquis aequali

Stam. filamenta quatuor, quorum duo sterilia.

Stylus simplex, stigmate bilamellato.

Caps. bilocularis, dissepimento medio placentifero, valvis parallelo.

Fl. dan. 363. Gärtn. t. 53.

81. **BONNAYA** Link. (1821.)

Cal. quinquepartitus, subbracteatus.

Cor. ringens. Lab. superius integrum, inflexum: inferius trilobum.

Stam. duo conniventia. Antherae biloculares, loculis oppositis.

Stylus simplex, stigmatе subdilatato.

Caps. linearis, subtetragona, valvis navicularibus: dissepimentum medio placentiferum.

Link et Otto icon. plant. t. 11.

[Chirita Don.]

82. **MICROCARPAEA** R. Br. (1810.)

Cal. tubulosus, pentagonus, quinquefidus.

Cor. bilabiata.

Caps. bivalvis, dissepimento contrario, demum libero.

83. **BEYRICHIA** Schlechtend. (1828.)

Cal. quinquepartitus, lacinis lateralibus maioribus, postica minima.

Cor. tubulosa, bilabiata, labio utroque integro.

Stam. quatuor, quorum duo sterilia inclusa.

Stylus apice incrassatus, stigmatе cupuliformi.

Caps. bilocularis, bivalvis, placenta centrali demum libera.

84. **PEPLIDIUM** Delil. (1813.)

Cal. tubulosus, quinqueidentatus.

Cor. tubulosa, quinqueloba.

Stam. subincurva.

Caps. bilocularis, non dehiscens, valvis fragilibus, placenta centrali.

Delil. fl. aegypt. t. 4.

85. **BAEA** Commers. Juss. (1789.)

Cal. quinquepartitus.

Cor. tubo brevi, limbo bilabiato: labio super. tri-infer. bilobo.

Stam. filamentis crassis, arcuatis, antheris conniventibus.

Caps. longa, corniculata, contorta, bilocularis, quadrivalvis, valvis demum contortis.

Lam. ill. t. 15.

86. **HENCKELIA** * (1817.)

Cal. quinquefidus.

Cor. tubuloso-ventricosa, limbo subaequaliter quinquelobo.

Stam. quatuor, quorum duo sterilia. Antherae biloculares, contiguae.

Stylus simplex, stigmatе bifido.

Caps. siliquosa, bilocularis, bivalvis, valvis demum tortis: dorso dehiscens, placentis revolutis, ut fructum bicapsularem mentiantur. Sem. numerosa.

[*Didymocarpus* Jack. Linn. transact. vol. 11. tab. 2. f. 3.]

Hook. exot. fl. 3. t. 227. Stenotocarpus Lindl. in Bot. 1847

87. *LYSIONOTUS* Don. (1825.)*Cal.* quinquepartitus.*Cor.* tubuloso-campanulata, limbo quinquelobo, bilabiato.*Stam.* quatuor inclusa: perfecta inferiora, antheris coalescentibus, superiora sterilia.*Stylus* simplex, stigmatē subinfundibuliformi.*Caps.* valvis e sutura dorsali dissilientibus. *Sem.* utrinque villis comosa.88. *CALCEOLARIA* Feuill. (1725.)*Cal.* quadrifidus, laciniis ovatis.*Cor.* bilabiata. Lab. sup. minimum acutum: inf. horizontaliter porrectum, calceiforme, marginibus involutis.*Stam.* filamentis brevibus, antheris bilocularibus, loculis divaricatis.*Stylus* brevis, stigmatē acutiusculo.*Caps.* bilocularis, dissepimento medio utrinque in placentas incrassato.*Hook.* exot. flor. t. 75. *Gärtn.* t. 62.89. *GYMNANDRA* Pall. (1776.)*Cal.* ovatus, longitudinaliter fissus, tridentatus.*Cor.* ringens, lab. sup. indiviso, infer. bi-tripartito, laciniis acutiusculis.*Stam.* filamenta fauci corollae inserta. Antherae cordatae.*Stylus* filiformis. Stigma capitatum.*Caps.* oblonga, bilocularis, dissepimento contrario, apice quadridentata, loculis monospermis. *Sem.* oblongum, sulcatum.*Willd.* in berl. Magaz. 5. t. 9. 10.90. *DIPLOPHYLLUM* Lehm. (1814.)*Cal.* disepalus, compresso-clausus, sepalis cordatis serratis.*Cor.* rotata, quadriloba, lobis oppositis minoribus.*Stam.* filamentis corolla brevioribus, antheris didymis.*Stylus* filiformis, stigmatē simplici.*Caps.* subrotunda, compressa, bilocularis, quadrivalvis, tetrasperma.[*Veronica Steven* in Linn. transact. 11. t. 31.]91. *SANCHEZIA* Ruiz et Pav. (1794.)*Cal.* quinquepartitus.*Cor.* tubulosa, quinquefida: lacinae revolutae, duae superiores breviores.*Stam.* quatuor, quorum duo sterilia. Antherae aristatae.*Caps.* bilocularis, bivalvis.*Ruiz et Pav.* flor. peruv. prodr. t. 32.92. *SCHWENKIA* Royen. L. (1767.)*Cal.* tubulosus, quinquefidus.*Cor.* tubulosa, limbo plicato quinquedentato aut undulato glandulae clavatae in limbo.

Stam. quinque, quorum tria sterilia.
Stylus simplex, stigmate subcapitato.
Caps. bilocularis, bivalvis, polysperma, valvis dissepimento parallelis.

Humb. nov. gen. 2. t. 180. 181.

93. *SCHIZANTHUS* Ruiz et Pav. (1794.)

Cal. quinquepartitus.

Cor. bilabiata, resupinata. Lab. sup. quinquepartitum, lacinis lateralibus bifidis, palato prominulo: lab. infer. tripartito, lacinia media emarginata, concava.

Stam. quatuor, quorum duo breviora sterilia, in cavitate lab. infer. ante pubem nidulantia. Antherae bilobae.

Stylus filiformis, stigmate simplici.

Caps. calyce minor, bilocularis, bivalvis: dissepimento utrinque placentifero. Sem. numerosa, rugosa.

Hook. exot. flor. t. 75. 86.

94. *SABMIENTA* Ruiz et Pav. (1794.)

Cal. quinquepartitus inaequalis.

Cor. obovato-ventricosa, basi fauceque coarctatis, limbo quinquefido aequali.

Stam. quinque, quorum tria sterilia inclusa.

Caps. (unilocularis circumscissa R. et P.) (Germen biloculare Schlechtend.)

[*Utricularia* *Feuill.* 3. t. 43.]

A C A N T H E A E. Cf. cl. XIV.

95. *JUSTICIA* Honst. L. (1737.)

Cal. quadri-quinquefidus bracteatus.

Cor. bilabiata.

Stam. Antherae biloculares, loculis saepe distantibus.

Caps. bilocularis, bivalvis: loculis dispernis: dissepimento valvis adnato contrario, retinacula seminum gestante.

Dianthera L. *Jacqu.* ecl. 1. t. 11. 12. 34. *Gärtn.* t. 51.

96. *DICLIPTERA* Juss. (1807.)

Cal. quadri-quinquefidus bracteatus.

Cor. bilabiata, labio inferiori subintegro.

Stam. Antherae biloculares.

Caps. subglobosa, subunilocularis: dissepimento demum libero, cui semina pauca ope retinaculorum adhaerent.

Ann. mus. 9. t. 23. f. 3. [*Diapedium* König 1806.]

97. *HYPOESTES* Soland. R. Br. (1810.)

Cal. quadrifidus, aequalis: involucri quadrifido subtrifloro inclusus.

Cor. bilabiata.

Stam. Antherae uniloculares.

Caps. *Justiciae*.

[*Justicia* Vahl. *symb.* 1. t. 1.]

98. *ERANTHEMUM* L. (1748.) R. Br. (1810.)
Cal. quadripartitus, aequalis.
Cor. hypocrateriformis, limbo quinquefido, subaequali.
Stam. quatuor, quorum duo sterilia, fertilia exserta.
Caps. Justiciae.
Andr. repos. 88. Bot. reg. 867. 879. [*Ruellia Venten. Cels.* t. 46.]
99. *NELSONIA* R. Br. (1810.)
Cal. inaequalis, quadripartitus.
Cor. infundibuliformis, limbo quinquefido, subaequali.
Stam. inclusa. Antherae loculis duobus divaricatis.
Caps. bilocularis, dissepimento adnato, elastico. Sem. numerosa, sine retinaculis.
100. *ELYTRARIA* Mx. (1803.)
Spicae bracteatae. *Cal.* quadri-quinquepartitus.
Cor. hypocrateriformis, limbo quinquefido, aequali.
Stam. quatuor, quorum duo sterilia. Antherae oblongae, biloculares.
Stylus setaceus, stigmate ligulato, recurvo.
Caps. bilocularis, bivalvis, valvis medio septiferis. Sem. pauca, ad dissepimenti imi axin (sine retinaculis) affixa.
Michaux flor. bor. amer. 1. t. 1.
101. *MENODORA* Humb. Bonpl. (1812.)
Cal. subcampanulatus, multipartitus.
Cor. infundibuliformis, limbo quinque-sexpartito, subaequali.
Stam. Antherae biloculares, longitudinaliter dehiscentes.
Stylus stigmate capitato.
Caps. bilocularis, bivalvis.
Humb. et Bonpl. plant. aequinoct. 2. t. 110.

BIGNONIEAE. Cf. cl. XIV.

102. *CATALPA* Juss. (1789.)
Cal. bipartitus.
Cor. campanulata, tubo ventricoso, limbo quadrilobo inaequali.
Stam. quinque, quorum tria sterilia.
Styl. stigmate bilamellato.
Caps. siliquaeformis, bivalvis, dissepimento valvis opposito.
Sem. plura, basi et apice membranaceo-papposa.
 [Biguonia L.] *Schkuhr* Handb. t. 175. Bot. mag. 1094.
- † 103. *ONCOMA* * (Oxera Labill. 1824.)
Cal. quadripartitus, scariosus.
Cor. tubulosa, fauce dilatata, limbo quadrifido subaequali.
Stam. quatuor, quorum duo sterilia, fertilia longe exserta, antheris bilocularibus.
Stylus incurvus, stigmate bifido.

Germen quadrilobum, carnosum.

Labill. nov. Caledon. t. 28.

104. *SCHREBERIA* Roxb. (1798.)

Cal. bilabiatus, inaequaliter dentatus.

Cor. campanulata, limbo quinque - septemfido, laciniiis emarginatis.

Stam. intra tubum, antheris ovatis erectis, bilocularibus.

Stylus simplex, stigmate bifido, acuto.

Caps. pyriformis, lignosa, bilocularis, bivalvis; loculis oligospermis. Sem. membranaceo - alata.

Roxburgh coromand. 2. t. 101.

105. *CYRTANDREA* Forst. (1775.)

Cal. quinquepartitus.

Cor. infundibuliformis, fauce ampliata, limbo subirregulari quinquelobo.

Stam. quatuor, quorum duo sterilia, intra tubum: fertilium filamenta curvata: antherae conniventes, biloculares.

Stylus, stigmate bilobo.

Bacca bilocularis, polysperma. Dissepimenta bifida, revoluta, laciniiis placentiferis.

Cyrtandreae Jack. *Forst. char. gen. t. 3. Linn. transact. vol. 14. t. 2. f. 1.*

106. *COLUMELLIA* Ruiz. et Pav. (1798.)

Cal. quinquepartitus, aequalis.

Cor. tubo brevissimo, limbo quinquepartito subaequali.

Stam. basi tubi inserta, antheris plicato - undulatis.

Stylus brevis, stigmate capitato.

Caps. bilocularis, bivalvis, polysperma, valvis apice bifidis, placentis septo adnatis.

Gesneriae? Flor. peruv. 1. t. 8. 12.

LENTIBULARIAE. Rich. R. Br.

107. *PINGUICULA* C. Gesner. (1555.)

Cal. bilabiatus, labio super. tri - infer. bipartito.

Cor. ringens calcarata: lab. sup. bilobo vel emarginato, infer. trilobo.

Stam. brevissima, germi adstantia. Antherae globosae.

Stylus brevissimus, stigmate bilobo, lobo inferiori incrassato.

Caps. unilocularis, polysperma: placenta globosa, libera.

Sem. cylindrica.

Sturm. fl. germ. Gärtn. t. 112.

108. *UTRICULARIA* L. (1737.)

Cal. bipartitus, coloratus, deciduus.

Cor. personata, calcarata. palato prominulo; lab. super. erecto, infer. declinato.

Stam. filamentis brevissimis incurvis, antheris cohaerentibus.

Stylus brevis, stigmate bilabiato, labio inferiori latiori.

Caps. unilocularis, polysperma, circumscissa, operculata: placenta libera globosa.

Sturm. fl. germ. Gärtn. t. 198.

PRIMULAEAE. Cf. cl. V.

109. HEMIANTHUS Nuttall. (1818.)

Cal. tubulosus, inferne fissus, limbo quadridentato.

Cor. bilabiata: lab. super. obsoleto, infer. tripartito, lacinia media ligulata, incurva, truncata.

Stam. filamentis bifidis, hinc antheriferis.

Stylus bifidus.

Caps. unilocularis, bivalvis, polysperma.

Journ. of the acad. of nat. scienc. of Philadelph. 1. tab. G. f. 2.

110. MICRANTHEMUM Mx. (1803.)

Cal. quadripartitus, lacinii superioribus minoribus.

Cor. subcampanulata, tubo brevissimo: limbo inaequaliter quadripartito, lacinia superiore minore.

Stam. convergentia, filamentis incurvis, basi appendiculatis, antheris subdidymis.

Stylus brevis, stigmate depresso-capitato, obliquo.

Caps. subglobosa, unilocularis, bivalvis, polysperma, placenta basilari.

Michaux flor. bor. amer. t. 2.

111. CAMPYLANTHUS Roth. (1821.)

Cal. quinquepartitus inaequalis.

Cor. infundibuliformis, tubo incurvo, limbo quinquepartito aequali.

Stam. filamentis incurvis, intra tubum: antheris peltatis unilocularibus.

Stylus stigmate simplici.

Caps. trilocularis, polysperma: placenta centralis: semina numerosa, compressa, margine membranaceo.

[*Eranthemum* L. suppl.]

112. STRIGA Lour. (1790.)

Cal. quadrifidus, persistens, lacinii subulatis aequalibus.

Cor. hypocrateriformis: tubo longo, superne incurvo: limbo quadripartito, lobo supremo maiori emarginato.

Stam. filamentis brevissimis, intra tubum, antheris oblongis.

Stylus, stigmate simplici.

Caps. unilocularis, polysperma.

113. CRYPTOHECA Blum. (1827.)

Cal. infundibuliformis, quadrifidus.

Cor. petala quatuor minima, aut nulla.

Stam. duo opposita, fundo calycis inserta. *Antherae* subrotundae.

Stylus lateralis, obtusus.

Caps. unilocularis, apice irregulariter circumscissa. *Sem.* numerosa.

Salicariae.

114. *BOERHAAVIA* Vaill. (1718.)

Involutum squamulatum, pauciflorum.

Cor. infundibuliformis, limbo plicato, sublobato.

Stam. 1. 2. 3.

Nectarium urceolatum, staminiferum.

Nucula monosperma, albumine centrali, embryo peripherico.

Nyctagineae. Jacqu. hort. vind. 1. t. 4. 7. *Cavan.* ic. 2. t. 112. *Gärtn.* t. 127.

115. *CODARIUM* Soland. Vahl. (1804.)

Cal. quinquepartitus, tubo persistente, laciniis aequalibus.

Cor. petalum unicum, lanceolatum, nectario annulari insertum.

Stam. calyci inserta, erecta: antheris bilocularibus.

Stylus incurvus, stigmate simplici.

Legumen pedicellatum, bivalve, monospermum, pulpa farinacea farctum.

Leguminosae. [Dialium Willd. *Röm. archiv.* 1. t. 6.]

116. *OLIGARRHENA* R. Br. (1810.)

Cal. quadri-partitus, bracteatus.

Cor. quadrifida, persistens.

Stam. inclusa.

Nectar. Squamulae quatuor, hypogynae.

Germen biloculare.

Epacrideae.

b. *Fructus inferi.*

117. *ANGURIA* Jacqu. (1763.)

Flores monoeci.

Cal. quinquedentatus.

Cor. cum calyce coalita, quinquefida.

Stam. antheris serpentino modo dehiscentibus.

Stylus semibifidus, stigmatibus bifidis.

Pepo subtetragonus, quadrilocularis, polyspermus. *Sem.* compressa.

Cucurbitaceae. Jacqu. amer. t. 155. *Plum.* ic. t. 22.

118. *DIPLARRHENA* Labill. (1798.)

Cal. corollinus, sexpartitus, laciniis inferioribus minoribus, superiore dissimili fornicata.

Stam. tria, quorum duo fertilia conniventia, tertium sterile.

Stigma tripartitum bilabiatum.

Caps. oblonga, trigona. *Sem.* plano-depressa, uniseriata.

Irideae. Labill. iter, 1. t. 15.

B. *Corolla pleiopetala.*

119. FONTANESIA Labill. (1791.)

Cal. quadripartitus, inferus.*Cor.* dipetala.*Stam.* elongata.*Styl.* stigmate bifido,*Caps.* bilocularis, chartacea, non dehiscens, loculis monospermis.*Jasmineae.* Labill. pl. syr. dec. 1. t. 1. Lam. ill. t. 22.[? *Tetrapilus* Lour.]

120. CIRCAEA Lobel. (1591.)

Cal. bipartitus, superus, coloratus.*Cor.* dipetala, petalis emarginatis.*Stam.* erecta, antheris bilocularibus.*Stylus* simplex, stigmate bicapitato.*Caps.* obovata, setis uncinatis hispida, bilocularis, loculis monospermis. Sem. embryo erecto.*Onagris* affinis. Sturm. fl. germ. Gärtn. t. 24.

121. MILLINGTONIA Roxb. (1821.)

Cal. trisepalus, calyculatus.*Cor.* tripetala, squamis nectariferis ternis.*Stam.* filamentis apice bifidis, antheris bilobis.*Stylus* simplex.*Drupa* subbilocularis, bivalvis, monosperma. Embryo curvatus, conduplicatus, albumine nullo.

— — Roxb. corom. 3. t. 254.

122. NOTELAEAE Venten. (1803.)

Cal. quadrifidus, inferus.*Cor.* tetrapetala, petalis ope filamentorum basi nexis.*Drupa* monosperma, putamine chartaceo. Albumen rugosum.*Jasmineae.* Venten. sel. 25. [*Rhyospermum* Gärtn. t. 224.]

123. LINOCIERA Sw. apud Schreb. (1791.)

Cal. quadridentatus, inferus.*Cor.* tetrapetala, petalis linearibus.*Stam.* Antherae elongatae sessiles, petala opposita basi nectentes.*Drupa* bilocularis, loculis monospermis.*Sem.* embryo inverso.*Jasmineae.* Sw. fl. ind. occ. 1. t. 2. Gärtn. t. 215.

124. AIDELUS * (1827.)

Cal. quadripartitus.*Cor.* petalis quatuor, calyce multo minoribus.*Stam.* brevissima.*Stigma* sessile.*Caps.* compressa, unilocularis, bivalvis: placenta libera centralis.*Caryophyllaceae?*

125. *ACAENA* Mutis. Linn. (1771.)

Cal. basi squamis duabus suffultus, tubulosus, glochidibus armatus.

Cor. petala quatuor aut. quinque, libera vel basi coalita, apici calycis inserta.

Stam. 2 3. 4.

Stigmata duo, penicilliformia.

Sem. 1. 2. in calycis tubo inclusa.

Rosaceae. Jacq. eclog. 1. t. 55. [Aucistrum Forst. Lam. ill. t. 22. f. 1.]

126. *LABICHEA* Gaudich. Cand. (1825.)

Cal. tripartitus, subpentasepalus, inaequalis.

Cor. petala quinque obovato-orbicularia.

Stam. Antherae subsessiles, altera longiore basi pollinifera, dein producta in acumen sterile, apice suboperculatum.

Stylus filiformis, subincurvus.

Germen stipitatum uniloculare, ovulis binis.

Leguminosae.

II. *Flores imperfecti.*127. *MARGYRICARPUS* Ruiz. et Pav. (1794.)

Cal. quadri-quinquepartitus superus, lacinis extus ad basin spinula dentiformi instructis.

Cor. 0.

Stigma multifido-plumosum.

Drupa nucleo monospermo. Embryo inversus.

Rosaceae. Flor. peruv. prodr. t. 33.

128. *DIALIUM* N. L. Burm. (1768.)

Cal. quinquepartitus aut pentasepalus, subaequalis, sepalis exterioribus paullo latioribus.

Stam. receptaculo inserta, antheris crassis bilocularibus.

Stylus simplex.

Legumen intus pulposum, 1—2 spermum.

Leguminosae. [Arouna Aubl. guian. t. 5.]

129. *LITHOPHILA* Sw. (1788.)

Cal. pentasepalus, tribracteatus, inferus: sepala duo interiora navicularia erecta.

Cor. 0.

Stam. simplicia; antheris oblongis, erectis.

Stigma obtusum.

Stylus simplex.

Caps. bilocularis.

Chenopodeae. Sw. fl. ind. occ. 1. t. 1.

130. *DYSPHANIA* R. Br. (1810.)

Flores polygami monoeci.

Calyx tripartitus, coloratus, lacinis cochleariformibus.

Stam. basi calycis inserta.

Stylus simplex.

Pericarpium turbinatum, semini adnatum, calyce aucto-
cinctum. Sem. albuminosum, embryone peripherico.

Chenopodeae.

131. *CYPSELEA* Turp. (1806.)

Cal. campanulatus, persistens, coloratus, quinquepartitus,
laciniis duabus minoribus.

Cor. 0.

Stam. 2. 3. calyci inserta.

Stylus brevissimus, vel 0. *Stigmata* duo.

Caps. unilocularis circumscissa, polysperma.

Sedeae. Annal. du mus. 7. t. 121. f. 5.

132. *SUFFRENTIA* Bellard. (1796.)

Cal. tubuloso-campanulatus, quadrifidus, dentibus quatuor
minimis ad sinus.

Cor. 0.

Stam. tubo calycis inclusa.

Stylus filiformis, stigmatibus capitato.

Caps. bivalvis, demum unilocularis, septo ad apicem
evanido. Sem. numerosa.

Salicariae. Act. acad. taurin. 7. t. 1. f. 1.

133. *PIMELEA* Banks et Soland. Gärtn. (1788.)

Perianthium infundibuliforme, limbo quadrifido.

Cor. 0.

Stam. faucibus inserta, laciniis exterioribus opposita.

Stylus lateralis, stigmatibus capitato.

Nucula corticata, monosperma. Embryo inversus.

Thymelaeae. Gärtn. t. 39. Smith. nov. holl. t. 11. Labill.
nov. holl. t. 3—7. Rudge in Linn. transact. 10. f. 13. 14.

[*Banksia* Forst. char. gen. t. 4.]

134. *REICHENBACHIA* * (1824.)

Cal. corollinus, tubulosus, quinquedentatus, inferus.

Stam. hypogyna, filamentis capillaribus, calycem ae-
quantibus.

Stylus simplex, stigmatibus pedicellato.

Utriculus monospermus, calycis basi inclusus.

Nyctagineae? Bullet. de la soc. philom. Paris, 1823. t. 11.

135. *FRAXINUS* Virgil.

Flores polygami.

Cal. quadripartitus, aut nullus.

Cor. tetrapetala, quadripartita, aut nulla.

Stam. brevia, antheris ovatis, bilocularibus, extus de-
hiscentibus.

Stigma subsessile, bifidum.

Fructus bilocularis, compressus, apice alatus, submo-
nospermus.

Jasmineae? *Acerinae?* Schk. Handb. t. 375.

136. *TRIADICA* Lour. (1790.)

Flores dioeci, amentacei.

Amenta filiformia, distincta tuberculis multifloris. *Perianthium* minimum, campanulatum trifidum.

Cor. 0.

Stam. filamentis brevissimis, planis: *Antheris* bilobis, lobis lateraliter dehiscentibus.

Stylus brevis, crassus, stigmatibus tribus oblongis.

Bacca triloba, trilocularis, monosperma.

137. *SALIX* Varr. Cat.

Flores amentacei, plerumque dioeci.

Squamae ut plurimum diandrae.

Stam. filamentis simplicibus, antheris erectis bilocularibus.

Stylus simplex, vel nullus. *Stigmata* duo simplicia vel divisa.

Glandula nectarifera basilaris.

Caps. bivalvis. *Sem.* basi pilosa, valvis adhaerentia, exalbuminosa. *Embryo* erectus.

Amentaceae. Schk. Handb. t. 317. Sturm. fl. germ. Gärtn. t. 90.

138. *PIPER* Horat. Cels.

Amenta subandrogyna.

Cal. nullus. *Squamae* appressae laterales, vel *Bracteae* peltatae aut lineares.

Stam. duo, vel incerti numeri: filamentis brevissimis, antheris bilocularibus.

Stylus unicus vel nullus.

Stigma 1, peltato-fimbriatum, vel 2. 3.

Bacca monosperma. *Sem.* albuminosum. *Embryo* inversus.

Pipereae. Humb. nov. gen. t. 3. Hook. exot. flor. t. 22. 23. 92. Gärtn. t. 92.

[*Peperomia* R. et P.]

139. *CLADIUM* P. Brown. (1756.)

Spiculae undique imbricatae, uni - biflorae, squamis infimis vacuis.

Stam. duo, tria: antheris erectis, bilocularibus.

Stylus capillaceus, stigmatibus binis hirsutis.

Nucula putamine duplici.

Cypereae. Sturm. fl. germ. Fl. dan. 1202.

140. *LEMNA* Dalechamp. (1587.)

Perianthium spathaceum, membranaceum.

Stam. filamentis simplicibus subcurvatis.

Antherae didymae, subglobosae, extus dehiscentes.

Stylus stigmatibus obtuso.

Utriculus monospermus.

Najades. Wolff comment. de Lemna. Erl. 1801. 4. cum tabula.

141. *CERATIOLOA* Mx. (1803.)

Flores dioeci e gemmulis imbricatis.

Squamae oppositae, appressae.

Cal. et *Cor.* 0.

Stam. filamentis planis, antheris bilocularibus.

Stylus crassiusculus, erectus. *Stigma* multipartitum, radiatum.

Bacca nuculis duabus monospermis.

Empetrum ~~ae.~~ Hook. in Bot. mag. 2758.

D I G Y N I A.

142. CRYPTA Nuttall. (1817.)

Cal. inferior, disepalus.

Cor. di-tripetala, clausa.

Styl. 0. *Stigmata* duo vel tria, minima.

Caps. bi-trivalvis, bi-trilocularis, loculis oligospermis.

Sem. subcylindrica, striata.

Salicariae. Journ. of acad. of Philad. 1. t. 6. f. 1.

143. GUNNERA L. (1767.)

Amentum verticillatum: *squamis* persistentibus.

Cal. bidentatus.

Stam. filamentis duobus brevissimis: antheris oblongis.

Styli breves.

Drupa cum calyce tumido coalita.

Urticeae. Lam. ill. t. 801. [Misandra Juss. Panke Feuill.]

144. PODOSTEMON Mx. (1803.)

Cal. disepalus, subsquamatus.

Stam. filamenta pedicellata: antheris bilocularibus.

Stigmata duo sessilia.

Caps. ovata, striata, bilocularis, bivalvis polysperma.

Podostemeae Rich. *Pluknet.* t. 138. f. 1. *Michaux flor.*
bor. americ. t. 44.

G R A M I N A. Cf. cl. III.

145. ANTHOXANTHUM L. (1745.)

Panicula spicaeformis.

Cal. bivalvis, triflorus.

Cor. duae inferiores neutrae univalves aristatae: media bivalvis perfecta, mutica, valvis inaequalibus acutis.

Stam. filamentis capillaribus: antheris bilocularibus.

Styli duo hirsuti.

Sturm. fl. germ.

146. CORYCARPUS Zea. (1806.)

Cal. bivalvis, rigidus, tri-sexflorus.

Cor. cartilaginea, bivalvis, valva inferiore superiorem conduplicatam involvente.

Sem. corolla indurata involutum.

Festuca diandra Mx. t. 10. *Diarrheua americana* Pal. Beauv.
Agrost. t. 25. f. 2. *Jacqu. ecl. gram.* t. 29.

147. SCHMIDTIA Trattin. (1819.)

Flores composito - corymbosi.

Cal. 0.*Cor.* bivalvis, valvis inaequalibus denticulatis.*Stam.* antheris ovalibus, quadrilocularibus.*Styli* hirsuti.*Semen* maximum liberum.*Sturm.* fl. germ. [Coleanthus Seidel. Röm. et Schult.]

148. REIMARIA Flügg. (1810.)

Spicae paniculatae.*Cal.* univalvis uniflorus.*Cor.* bivalvis acuminata.*Humb.* nov. gen. t. 21.

CLASSIS III. T R I A N D R I A.

CONSPECTUS

CLASSIS TERTIAE

M O N O G Y N A E

I. Fructibus inferis.

Onagreae, Valerianeae et Hydrocharideae n. 149—157.
Spathaceae et Coronariae 158—173.
Haemodoreae et Affines 174—176.
Irideae 177—182.
Santalea 183.

II. Fructibus superis.

Variae 184—189.
Commelineae 190—193.
Aliae 194—214.
Restieae 215—222.
Cypercae 223—246.
Gramineae 247—251.

D I G Y N A E.

G R A M I N A.

I. Calycibus unifloris.

A. Floribus spicatis 252—264.
B. Floribus racemosis 265—272.
C. Floribus paniculatis,
a. glomerato-fasciculatis 273—281.
b. vere paniculatis 282—297.

II. Calycibus sesquifloris 298—312.

III. Calycibus bi-trifloris.

A. Corolla altera perfecta, altera mascula, aut neutra bivalvi 313—321.
B. Corolla utraque, et, si tertia affuerit, et haec perfecta 322—334.

IV. Calycibus multifloris.

A. Floribus spicatis 339—349.
B. Floribus glomerato-paniculato-spicatis 350—353.
C. Floribus paniculatis 354—371.

T R I G Y N A E

Gramineae 372. 373.

Restieae 374—376.

Aliae 377—387.

M O N O G Y N I A.

I. *Fructus inferi.*

149. PRIEUREA Cand. (1828.)

Cal. tubo elongato, limbo tripartito persistente.

Cor. tripetala.

Stamina laciniis calycis alterna, filamentis brevibus.

Stylus brevis.

- Caps. trilocularis.*
Onagrac.
150. **VALERIANA** Brunsfels. (1532.)
Cal. brevissimus, margine integerrimo.
Cor. tubulosa, basi gibba: limbo regulari, quinquelobo (rarius trilobo).
Stam. fauci corollae inserta. Antherae biloculares.
Stylus simplex, stigmatate plerumque trifido.
Achenium, pappo plumoso.
Valerianeae. Sturm. fl. germ. [Phyllactis Pers. corollae limbo trilobo. Astrephia Dufresn.]
51. **FEDIA** Adans. (1763.)
Cal. dentatus, persistens.
Cor. tubulosa, limbo quinquelobo subregulari.
Stam. fauci corollae inserta. Antherae biloculares.
Stylus simplex, stigmatate trifido.
Caps. trilocularis, loculis duobus crassis, calyce aucto coronata.
Valerianeae. Reichenbach, icon. t. 60 — 70.
[Valerianella Vaill.]
152. **AXIA** Lour. (1790.)
Perianthium triphyllum, foliolis acutis, brevibus, inaequalibus, deciduis.
Cor. campanulata, minima, limbo decemlobo.
Stam. antheris didymis.
Stylus stigmatate crassiusculo.
Fructus monospermus, hirsutus.
Nyctagineae? Valerianeae?
153. **SONEBILA** Roxb. (1821.)
Cal. tri-quadrifidus, patens.
Cor. rotata, tri-quadrupartita.
Stam. (3. 4.) tubo corollae inserta, declinata, laciniis corollae alterna. Antherae longissimae, arcuatae, biloculares, rostratae, apice poris geminis dehiscentes, basi filamentis affixae.
Stylus capillaris, stigmatate capitato.
Germen tri-quadriloculare, ovulis numerosis farctum.
Melastomeae Lindl.
154. **BURMANNIA** L. (1737.)
Cal. corollinus tubulosus trialatus vel tricarinatus, limbo sexfido, laciniis alternis minimis.
Stam. Filamenta brevissima sub laciniis calycis minimis appressa planiuscula, utrinque antherae loculos seiunctos gerunt. (Hinc Hexandriae Linnaeus non sine veri specie annumeravit.)
Stylus stigmatibus tribus capitatis.
Caps. trilocularis, polysperma: placentae inter dissepimenta.
Hydrocharideae Mart. Roxb. coromand. 3. t. 242. Mart. nov. gen. 1. t. 5. [Tripterella Mx. Maburnia Thuars.]

155. *ELODEA* Mx. (1803.)

Spatha subbifida, minima. Flores diclini vel hermaphroditi.
Perianthium tenerum, tubulo gracili, limbo sexpartito,
 laciniis interioribus maioribus.

Stamina tria fertilia, laciniis exterioribus opposita, earum-
 que basi inserta. Accedunt in quibusdam stamina
 sterilia sena in masculis, terna in femineis floribus.

Stylus filiformis: stigmatibus tribus bifidis.

Fructus oblongus, unilocularis, gelatina repletus, oli-
 gospermus.

Sem. oblongo - cylindrica.

Hydrocharideae. Humb. plant. aequil. 2. t. 128,
 [Serpicula Mühlenb. Pursh. Udora Nuttall.]

156. *MELOTHRIA* L. (1737.)

Cal. cum corolla connatus, basi tubulosus, limbo quin-
 quedentato.

Cor. rotata, quinqueloba.

Stam. filamenta conica, tubo corollae adnata, duo an-
 theris didymis, tertium simplici.

Stylus cylindricus. Stigmata tria.

Bacca oblonga, trilocularis polysperma.

Cucurbitaceae. Plum. ic. t. 66. f. 2.

157. *BENINCASA* Sav. (1818.)

Cal. cum corolla connatus; limbo plicato quinquefido,
 laciniis bi-tridentatis.

Cor. rotata quinqueloba.

Stam. aequalia, libera; filamentis planis.

Stylus columnaris, stigmatibus trilobis.

Bacca trilocularis polysperma pilosa.

Cucurbitaceae. Sav. mém. cucurbit. p. 6. ic.

S P A T H A C E A E. CORONARIAE. Cf. Cl. VI.

158. *CROCUS* Virgil. Cels.

Spatha monophylla.

Cor. tubus longissimus, limbus regularis sexpartitus.

Stam. corollae inserta, antheris sagittatis erectis.

Stylus filiformis, stigmatibus tribus dilatatis, serratis aut
 incisis.

Caps. trilocularis, polysperma. *Sem.* subglobosa.

Sturm. fl. germ.

159. *WITSENIA* Thunb. (1781.)

Cor. tubulosa, limbo sexpartito, patente vel connivente.

Stam. filamentis brevibus, tubo superne inserta. Antherae
 oblongae, erectae.

Stylus filiformis, stigmatibus trifidis.

Caps. duriuscula, trifariam dehiscens. *Sem.* plura, an-
 gulate - compressa.

Lam. ill. t. 30. (Ixia) Forst. in comment. botting. 9. t. 2.

160. OVIEDA * (1817.)

Spatha bivalvis.*Cor.* tubulosa, tubo gracili, longissimo: limbo plano sexpartito, subregulari.*Stam.* prioris.*Stylus* filiformis; stigmatibus tribus, gracilibus, bipartitis, patentibus.*Caps.* membranacea, triangularis. Sem. numerosa, triquetra.(Ixia) *Andr.* repos. 35. *Jacqu.* ic. rar. 2. t. 268 — 270.

[Lapeyrousia Pourr. Gawler.]

161. ANOMATHECA Gawler. (Ker. 1805.)

Spatha bivalvis, herbacea.*Cor.* hypocrateriformis, tubo gracili, limbo sexlobo, subregulari.*Stigmata* tria, gracilia, bipartita.*Caps.* subrotunda, papilloso-pruinosa. Sem. subrotunda.(Gladiolus) *Andr.* repos. 66.

162. HESPERANTHA Gawl. (1805.)

Spatha bivalvis, valva altera tubum corollae subaequante vel superante.*Cor.* hypocrateriformis, limbo tubum aequante sexpartito regulari.*Stigmata* tria, elongata, ad tubum usque distincta.*Caps.* oblonga, membranacea. Sem. rugosa, corticata.(Ixia) *Andr.* repos. 44. 59. *Jacqu.* ic. rar. 2. t. 279.

163. GEISSORRHIZA Gawl. (1805.)

Spatha bivalvis, scariosa, lacera, tubum subaequans.*Cor.* subhypocrateriformis, limbo sexpartito, regulari.*Stylus* inclinatus. *Stigmata* latiuscula, crispula, patentia.*Caps.* membranacea, subtrigona. Sem. numerosa, minuta.(Bulbus fastigiatus, squamis crustaceis, imbricato-suggestis, ora infima dentato-fissili, inflexa, unde nomen: *γέλασος*, corona prominula.)(Ixia) *Bot. mag.* 672.

164. TRICHONEMA Gawl. (1805.)

Spatha subherbacea, valva altera elongata.*Cor.* tubus brevissimus; limbus, campanulatus, regularis, sexpartitus.*Stam.* tubi ori imposita; filamentis pilosis.*Stigmata* tria, bipartita, gracillima.*Caps.* membranacea. Sem. plurima, globosa.(Sisyrinchium) *Column.* ecphr. 1. p. 327.(Ixia) *Jacqu.* ic. rar. 2. t. 271. *Engl. bot.* t. 2549.

[Romulea Maratt. Sebast.]

165. SPARAXIS Gawl. (1805.)

Spatha bivalvis, scariosa, lacera, apice maculata.*Cor.* subregularis, campanulata, sexpartita.*Stam.* subinclinata.

Stigmata tria, recurva.

Caps. subrotunda. Sem. globosa.

Bot. reg. t. 258. (Ixia) Jacqu. ic. rar. 3. t. 273. 274.

166. Ixia Miller. (1731.)

Spatha bivalvis, mediocris, scariosa.

Cor. tubo gracili, limbo regulari, sexpartito.

Stam. filamentis abbreviatis.

Stigmata angusta, recurva.

Caps. membranacea. Sem. globosa.

Andr. repos. 50. Bot. reg. 530.

167. Diasia Cand. (1802.)

Spatha bivalvis, subherbacea, patens.

Cor. hexapetaloidea, aequalis, patens, laciniis cuspidatis.

Stam. ascendentia.

Stigmata recurva.

Caps. turbinata. Sem. pauca, globosa.

Redout. lil. t. 54. (Gladiolus) *Andr.* rep. 62. [Aglaea Pers.
Melasphaerula Gawler.]

168. Gladiolus Plin.

Spatha bivalvis subherbacea.

Cor. sexpartita, subringens.

Stam. ascendentia, antheris supra basin dorso affixis.

Stigmata tria latiuscula, recurva.

Caps. oblonga, trigona. Sem. alata.

Reichenb. ic. t. 598 — 600.

169. Antholyza L. (1741.)

Spatha bivalvis, subconvoluta.

Cor. tubus gracilis; limbus ringens, labio superiori elongato, inferiori e laciniis quinque, quarum laterales refractae.

Stigmata indivisa.

Caps. globosa, coriacea. Sem. globosa, corticata.

Andr. rep. t. 32.

170. Babiana Gawler. (1805.)

Spatha subtrivalvis, valva intima partita.

Cor. tubulosa, limbo sexpartito irregulari.

Stigmata patentia, latiuscula.

Caps. coriacea. Sem. baccata.

171. Tritonia Gawler. (1805.)

Spatha bivalvis, dentata.

Corolla campanulata, regularis vel subbilabiata, saepe resupinata: saepe etiam tubo elongato gracili, limbo rotato.

Stam. ascendentia.

Stigmata patentia, nonnunquam bifida.

Caps. subrotunda. Sem. globulosa.

Bot. reg. 135. 747.

172. *WATSONIA* Gawler. (1805.)

Flores spicati.

Spatha bivalvis, appressa, apice maculata.

Cor. basi tubulosa, limbo sexpartito regulari, vel subbilabiato.

Stam. ascendentia.

Stigmata bipartita.

Caps. cartilaginea. Sem. subalata.

Bot. mag. 553. (Ixia) Andr. repos. 177.

173. *ARISTEA* Ait. (1789.)

Spathae fasciculatae scariosae, raro herbaceae.

Cor. hexapetaloidica, regularis, planiuscula, tubo brevissimo.

Stam. recta, germini inserta.

Stylus cylindricus, stigmatibus tribus capitatis, vel latiusculis.

Caps. subprismatica. Sem. compressa.

Andr. bot. rep. 10.

174. *HAEMODORUM* Sm. (1798.)

Cal. corollinus sexpartitus, persistens.

Stam. basi sepalorum interiorum opposita.

Stylus filiformis, stigmate simplici.

Caps. trilocularis, loculis dispermis.

Sem. peltata, compressa, marginata.

175. *DILATIS* Berg. (1767.)

Cal. corollinus, hexapetalus, extus hirsutus, regularis.

Stamina sex, basi calycis inserta; tria sterilia breviora, tria longiora laciniis involuta. Antherae tres bisulcae, tertia maior.

Stylus filiformis, stigma simplex.

Caps. trilocularis, placenta triquetra libera. Sem. tria aut sex, peltata.

Smith exot. 1. t. 16.

176. *LACHNANTHES* Elliott. (1817.)

Cal. corollinus sexpartitus, extus lanatus, regularis, laciniis alternis angustioribus.

Stamina tria perfecta, basi calycis inserta, exserta. Antherae aequales.

Stylus filiformis, stigmate minuto, subtrilobo.

Caps. trilocularis, valvis introflexis placentiferis. Sem. numerosa, peltata.

[Heritiera Mx. fl. bor. amer. t. 4.]

I R I D E A E.

177. *IRIS* Theophr.

Spatha bivalvis, persistens.

Cor. basi tubulosa, nectarifera: limbo sexpartito, laciniis alternis saepe barbatis.

Pistillum petaloideum, trifidum: stigmatibus plicas in
exteriore laciniarum facie referentibus.

Caps. trilocularis, polysperma. Sem. angulata.

Schk. Handb. t. 5. a. b. *Gärtz.* t. 13.

178. *MORAEA* Mill. (1731.)

Spathae elongatae, subimbricatae.

Cor. sexpartita patens, laciniis interioribus minoribus, post
anthesin convolutis.

Stam. laciniis pistilli opposita: filamentis subconnatis.

Pist. } *Iris.*
Caps. }

Andr. rep. 83. *Bot. reg.* 312.

179. *MARICA* Schreb. (1789.)

Spathae elongatae, imbricatae.

Cor. sexpartita, subrotata, laciniis aequalibus, vel alternis
minoribus.

Stam. filamentis subconnatis; antherae cum stigmatibus al-
ternantes.

Stylus brevis, triqueter, stigmatibus petaloideis.

Caps. *Iris.*

[*Cipura* Aubl.] *Bot. reg.* 713. *Redout.* ill. 56. 172.

180. *PARDANTHUS* Gawler. (1805.)

Spatha bivalvis.

Cor. hexapetaloidea, rotata, laciniis aequalibus, unguiculatis.

Stam. filamentis capillaribus, antheris erectis.

Stigmata erecta, subcucullato-dilatata.

Caps. trilocularis: semina baccata globosa, placentae
liberae columnari affixa.

(Ixia) *Bot. mag.* 174.

181. *LIBERTIA* * (1825.)

Spatha bivalvis, remota, decidua, multiflora.

Cor. sexpartita, rotata, laciniis exterioribus minoribus an-
gustioribus.

Stam. filamentis capillaribus, antheris incumbentibus.

Stigmata tria simplicia.

Caps. subglobosa.

Reichenb. hort. 2. t. 157.

182. *OSYRIS* L. (1741.)

Flores dioeci vel polygami.

Cal. trifidus, turbinatus, persistens. *Annulus* nectarifer
trigonus.

Stam. filamentis brevissimis, antheris subrotundis.

Stylus simplex, stigmate tripartito.

Drupa globosa, monosperma.

Schk. Handb. t. 319. *Gärtn.* t. 216. *Santaleae.*

II. *Fructus superi.*

183. *EBENXYLON* Lour. (1790.)

Flores dioeci.

- Cal.* nullus.
Cor. tripetala.
Nectarium stellatum.
Stylus brevis.
Bacca trisperma.
184. **WACHENDORFIA** J. Burm. (1757.)
Cor. subsexpartita irregularis, lacinia suprema basi nectarifera, noctu reliquas obvolvens.
Stam. quinque aut sex, quorum duo vel tria sterilia.
Antherae incumbentes.
Stylus declinatus, stigmatē simplici.
Caps. trilocularis, trivalvis. Sem. solitaria, setosa.
J. Burmann Wachendorfia. Amstel. 1757. fol. Cürtn. t. 15.
185. **HAGENBACHIA** Nees. (1823.)
Spathae remotae, multiliorae.
Cor. sexpartita, regularis, lacinii alternis minoribus.
Stam. in apice laciniarum corollae interiorum. Filamenta brevissima. *Antherae* erectae, biloculares.
Stylus filiformis, stigmatē subcapitato.
Caps. trilocularis, loculis dispermis,
Nov. act. nat. cur. vol. 11. t. 2.
186. **BRODIAEA** Sm. (1811.)
Cal. corollinus, tubulosus, sexfidus, persistens.
Squamae petaloideae, ad faucem, cum staminibus alternantes.
Stam. Filamenta cum tubo connata. *Antherae* oblongae, utrinque dehiscentes.
Stylus filiformis, stigmatē trilobo.
Caps. obovata, trilocularis, polysperma. Semina in quovis loculo biserialia, axi centrali affixa.
Sm. Linn. transact. 10. t. 1. *Salisb.* parad. londin. t. 98.
Bot. reg. t. 1183. *Bot. mag.* t. 2877.
187. **XIPHIDIUM** Löffling. (1758.)
Perianthium hexapetalum, duplici petalorum serie: externa calycina maiora, interna corollina minora.
Stamina basi petalorum internorum inserta. *Antherae* oblongae.
Stylus filiformis, stigmatē trilobo.
Caps. baccata, trilocularis.
188. **HETERANTHERA** Ruiz. et Pav. (1794.)
Cor. tubus gracilis, limbus sexpartitus subaequalis.
Stamina fauci corollae inserta. *Antherae* inaequales.
Stylus simplex, stigmatē subtubuloso fimbriato.
Caps. trivalvis, polysperma: semina placentis ternis filiformibus valvarum affixis.
[Leptanthus Michaux fl. bor. am. t. 5, Hook. exot. fl. 2. t. 94.] Pontederaceae.

189. GILLIESIA Lindl. (1826.)

Perianthium e bracteis quinque imbricatis calyciformibus
Corollae rudimentum carnosum, antice labelliforme.

Urceolus germen cingens, staminifer.

Stam. sex, quorum anteriora tria fertilia, tria posteriora sterilia dentiformia.

Stylus filiformis, stigmate capitato, trigono.

Caps. trilocularis, polysperma.

Bot. reg. 992.

COMMELINEAE.

190. COMMELINA Dillen. (1732.)

Involucrum conduplicatum, cucullatum.

Cal. trisepalus.

Cor. tripetala, petalo uno aut altero saepe abortiente.

Stam. tria fertilia, tria sterilia glandulosa, cruciata.

Stigma simplex.

Caps. trilocularis, trivalvis, valvis medio septiferis. Semen in singulis loculis subbina.

Schmidel ic. t. 41. 49. *Lam.* ill. t. 35.

191. ANEILEMA R. Br. (1810.)

Involucrum loco bractee remotae.

Cal. trisepalus.

Cor. tripetala.

Stam.

Stigm. } Commelinae.

Caps. }

Bauer ill. nov. holl. t. 6. Bot. reg. t. 659.

192. CALLISIA Löffl. (1758.)

Cal. trisepalus.

Cor. tripetala.

Stam. Filamenta exserta, apice dilatata. Antherae globosae geminae, lateri interiori apicis filamentorum affixae.

Stylus capillaris. Stigmata tria penicilliformia.

Caps. bilocularis, disperma.

[*Hapalanthus* Jacqu. amer. t. 11.] *Lam.* ill. t. 35.

193. SYENA Schreb. (1789.)

Cal. trisepalus, persistens.

Cor. tripetala.

Stam. Filamenta capillaria. Antherae oblongae biloculares.

Stylus filiformis. Stigma trifidum.

Caps. globosa, trivalvis, hexasperma, valvis placentiferis.

[*Mayaca* Aubl. t. 15.]

194. WILLICHIA Mutis. L. (1771.)

Cal. quadrifidus, persistens, laciniis ovatis patentibus.

Cor. rotata, tri-quadriloba.

Stam. angulis limbi corollae inserta. Antherae subrotundae, erectae, biloculares.

Stylus declinatus. Stigma obtusum.

Caps. bilocularis, bivalvis, dissepimento valvis contrario.

Sem. plurima, subglobosa, minuta. Placenta globosa in loco utroque.

Personatae.

195. *HYDRANTHELIUM* Kunth. (1825.)

Cal. profunde quadripartitus, laciniis alternis obtusis trinerviis, alternis lanceolatis acutis.

Cor. infundibuliformis, limbo tritido, lobo tertio emarginato.

Stam. tria, fauci corollae inserta, cum lobis alternantia.

Antherae oblongae, erectae, biloculares.

Stylus filiformis. Stigma subbilidum.

Caps. bilocularis, polysperma, dissepimento utrinque placentaifero. *Sem.* cylindrica, striata.

Personatae. Humb. nov. gen. t. 646.

196. *MACROLOBIUM* Schreb. (1789.)

Cal. quadri-quinquefidus, basi bracteis duabus suffultus.

Cor. petalum unicum, unguiculatum, planum aut undulato-plicatum. Petalorum quatuor quandoque rudimenta parva.

Stam. Filamenta elongata, quartum interdum sterile.

Antherae tetragonae.

Germen pedicellatum, stylo filiformi superatum.

Legumen monospermum.

Leguminosae. Colebrook. in Linn. transact. 12. t. 17. [Outea et Vouapa Aubl. t. 7. 9.]

197. *CNEORUM* L. (1737.)

Cal. tri-quadridentatus, persistens.

Cor. tri-tetrapetala.

Receptaculum hemisphaericum.

Stylus prismaticus. Stigma trifidum.

Drupae baccatae tres, coadunatae: putamine biloculari, loculis monospermis. Embryo conduplicatus.

Terebinthaceae. Lam. ill. t. 27. [Chamaelea Tourn. Gärtn. t. 70.]

198. *RUMPHIA* L. (1767.)

Cal. tubulosus, trifidus.

Cor. tripetala, petalis oblongis, patentibus.

Stam. exserta.

Drupa turbinata, nuce triloculari, trisperma.

Terebinthaceae? Lam. ill. t. 25.

199. *COMOCLADIA* P. Brown. (1756.)

Cal. tripartitus.

Cor. tripetala, petalis elongatis.

Stam. brevia.

Stylus nullus. Stigma simplex, sessile.

Drupa monosperma.

Terebintheae. Lam. ill. t. 27. [Dodonaea Plum. t. 118. f. 1.]

200. OLAX L. (1747.)

Cal. integer, post anthesin ampliatus.

Cor. penta-hexapetala: petalis quatuor, per paria filamentorum iunctis, seu tribus semibifidis, quinto bero, aut petalis sex eodem modo iunctis. Appendiculi filiformes, simplices aut bifidae, petalis mediis insertae.

Stam. tria, quorum duo petalis coalitis adnata, tertium liberum. Antherae erectae biloculares,

Stylus filiformis.

Drupa monosperma, calyce cincta.

Olacinae Mirb. Gärtn. t. 201. [Spermaxyron Labill. holl. t. 233. Fissilia Commers. Lam. ill. t. 28.]

201. HIPPOCRATEA L. (1737.)

Cal. quinquepartitus, coloratus, deciduus.

Cor. pentapetala, petalis apice foveolatis.

Urceolus perigynus, staminifer.

Stam. tria, filamentis capillaribus, urceolo insertis: antheris transversim dehiscentibus.

Stylus simplex, stigmatibus subtrilobis.

Capsulae tres, bivalves, samaroideae. Sem. deorsum alati funiculo umbilicali latissime expanso.

Hippocrateae Juss. ann. mus. 18. p. 486. [Familia non satis certo stabilita.] Lam. ill. t. 28. Rudg. guian. t. 2. Roxb. corom. 3. t. 205.

202. SALACIA L. (1771.)

Cal. quinquelobus aut quinquepartitus.

Cor. pentapetala, petalis apice planis.

Urceolus perigynus, extra stamina.

Stam. filamentis basi dilatatis; antheris peltatis, transversim dehiscentibus.

Stylus brevissimus. Stigma subtrigonum.

Bacca subtrilocularis, loculis oligo-vel polyspermis.

Hippocrateae Juss. [Tontelea Aubl. t. 10. Lam. ill. t. 3. Anthodon Ruiz. et Pav. Kunth. in Humb. nov. t. 443. Calypso Thuars. veg. afr. t. 6. Tonsella Schreb. Anthodus Mart. Raddisia Leandr. in act. acad. monach. t. 15.]

† 203. JOHNSIA Roxb. (1821.)

Cal. quinquefidus.

Cor. pentapetala, petalis unguiculatis.

Urceolus perigynus.

Stam. Antherae sessiles in margine urceoli.

Bacca monosperma.

Hippocrateae.

14. **BOLDOA** Cav. (1802.)

Cal. corollinus, tubulosus, limbo plicato, subquadridentato.

Stamina annulo perigyno inserta, unilateralia.

Stylus filiformis, stigmatē simplici.

Achenium calyce persistenti inclusum.

Nyctagineae. *Car.* hort. matrit. t. 7. [*Salpianthus*

Bompl. in Humb. pl. aequin. t. 41.]

15. **OXYBAPHUS** Herit. (1790.)

Involutrum quinquefidum.

Cal. corollinus infundibuliformis, brevissimus.

Achenium, calyce explanato scarioso cinctum.

Nyctagineae. [*Mirabilis* *Cur.* ic. t. 19. 379. 437. *Calyx-hymenia* *R. et P.* flor. peruv. t. 75. *Orteg.* dec. 5. t. 1. 8. t. 11.]

106. **ANTHOBOLUS** R. Br. (1810.)

Flores dioeci.

Cal. trisepalus, deciduus.

Stam. basi sepalorum inserta.

Stigma sessile, trilobum.

Drupa monosperma.

Santaleis affinis.

107. **PSEUDANTHUS** Sieber. (1825.)

Flores monoeci, masculi terminales glomerati; feminei axillares solitarii.

Cal. sexpartitus, foliaceus, persistens.

Cor. 0.

Antherae sessiles subglobosae in basi calycis.

Nux sexcostata, monosperma.

Santaleis affinis.

108. **ENDIANDRA** R. Br. (1810.)

Cal. corollinus sexfidus, fauce glandulosa.

Cor. 0.

Stam. perigyna: antheris adnatis posticis bifocularibus, a basi ad apicem dehiscentibus.

Stylus simplex.

Drupa monosperma.

Laurinae Venten.

109. **LÖFLINGIA** L. (1758.)

Cal. quinquepartitus, lacinii exterioribus basi bidentatis.

Cor. pentapetala, petalis minimis conniventibus.

Stam. antheris didymis.

Stylus simplex, stigmatē trifido.

Caps. unilocularis, trivalvis, polysperma. *Sem.* placentae centrali affixa.

Paronychieae Juss. *Löfling.* it. t. 1. *Cav.* ic. t. 94.

Lam. ill. t. 49.

110. **ORTEGIA** Löfl. (1758.)

Cal. pentapetalus.

Cor. 0.

Stamina tria fertilia: duo sterilia plerumque squamis formia.

Stylus simplex, stigmatē capitato vel trifido.

Caps. unilocularis, apice trifariam dehiscens. *Sem.* fun-
culis umbilicalibus capillaribus ad fundum capsul-
haerentia.

Paronychieae. Lam. ill. t. 29. Cav. ic. t. 47. Gärtn. t. 41.

211. *ROTALA* L. (1771.)

Cal. tubulosus, membranaceus, tridentatus.

Cor. 0.

Stamina tria, medio calycis tubo inserta.

Stylus stigmatē trifido.

Caps. calyce tecta, trilocularis, polysperma.

Salicariae. Lam. ill. t. 435.

212. *POLYCNUM* L. (1737.)

Cal. corollinus pentasepalus, bracteis duabus suffultus.

Stam. tria aut quinque; filamentis basi subcoadunatis; an-
theris bilocularibus.

Stylus brevissimus, stigmatē bifido.

Utriculus monospermus, apice marginatus.

Chenopodeae. Schk. Handb. t. 5. Sturm. fl. germ. Gärtn.
t. 128. [Hemichroa R. Br.]

213. *THRELKELDIA* R. Br. (1810.)

Cal. urceolatus, intra marginem truncatum squamis tribus
membranaceis.

Stamina squamis opposita.

Utriculus calyce baccato inclusus. Semen ovatum albu-
minosum. Embryo periphericus.

Chenopodeae.

214. *POSIDONIA* König. (1806.)

Perianthium nullum, nisi *Filamenta* dilatata calycina
persistentia, basi extus antherifera.

Antherae adnatae, lobis basi solutis.

Stigma sessile villosum.

Bacca monosperma.

Najades. Ann. of bot. 2. t. 6. [Zostera. L. Kerner & W.
Caulinia R. Br. Ruppia Labill.]

R E S T I E A E.

*Familia, junceis finitima, numero partium ternario (duplicato),
culmo saepe enodi, vaginis aphyllis obsesso, perianthio
tripartito aut hexaphyllo bracteato. Sed differt: staminibus
cum sepalis externis alternantibus, floribus declinatis et embryone
ad basin albuminis sito. Plantae hemisphaerio australi pec-
uliāres.*

215. *XYRIS* Gronov. (1743.)

Flores capitati, squamis unifloris, undique imbricatis, in-
ferioribus vacuis.

Calyx corollinus tripartitus.

Stamina fertilia tria, laciniis calycis inserta; sterilia totidem apice penicillata, cum laciniis calycis alterna.

Stylus trifidus, stigmatibus capitatis.

Caps. unilocularis, trivalvis, polysperma.

Lam. ill. t. 36. *Transact. of Linn. soc.* 10. t. 15.

216 ABOLBODA Humb. Bonpl. (1812.)

Flores capitati, squamis unilioris, inferioribus vacuis.

Calyx corollinus, tubo cylindrico, limbo hypocrateriformi trilobo.

Stamina omnia fertilia, fauci calycis imposita.

Stylus trifidus, stigmatibus bi-trifidis.

Caps. unilocularis, trivalvis, polysperma.

Hamb. plant. aequinoct. 2. t. 114.

217. RESTIO Rottböll. (1772.)

Flores dioeci, amentaceo - spicati.

Perianthium ebracteatum, sexvalve, duplici serie, valvis interioribus acutiusculis.

Stam. 2 3. *Antherae* simplices peltatae.

Stylus 2 - 3 - partitus.

Caps. bi - trilocularis, angulis dehiscens. Sem. solitaria.

Rottböll. ic. rar. t. 2. *Labill. nov. holl.* t. 226. 227.

218. LEPTOCARPUS R. Br. (1810.)

Flores dioeci, amentaceo - spicati.

Perianth. sexvalve.

Stam. tria, antheris simplicibus peltatis.

Stylus simplex, stigmatibus duobus aut tribus.

Nucula monosperma, basi styli coronata.

[*Schoenodum* ♀ *Labill. nov. holl.* t. 229]

219. THAMNOCHORTUS Berg. (1767.)

Flores dioeci, amentaceo - spicati.

Perianth. sexvalve.

Stam. tria, antheris oblongis, incumbentibus.

Stylus simplex, stigmate indiviso.

Nucula perianthio tecta.

[*Restio* Thunb. diss. ic. t. 5. 6. *Berg. cap.* t. 5. f. 8.]

220. WILDENOVIA Thunb. (1790.)

Flores dioeci, amentaceo - spicati.

Perianth. sexvalve.

Stam. tria, antheris compressis oblongis, utrinque marginatis, ope rimarum duarum longitudinalium dehiscentibus.

Stylus simplex, basi orbiculo sexlobo cinctus. Stigmata duo aut tria.

Drupa monosperma.

Act. acad. holm. 1790. t. 2. f. 1 — 3.

221. LYGINIA R. Br. (1810.)

Flores dioeci subfasciculato - spicati.

Perianth. sexvalve.

Stam. filamentis coalitis, antheris didymis, utrinque bifido
Stylus tripartitus.

Caps. trilocularis, angulis dehiscens. Sem. solitaria.
 [Schoenodum ♂ Labill. nov. holl. t. 229.]

222. *HYPOCHLAENA* R. Br. (1810.)

Flores dioeci, masculi amentacei.

Perianth. sexvalve.

Stam. filamentis distinctis capillaribus, antheris simpliciter peltatis.

Stylus bi-tripartitus, deciduus.

Nux monosperma, perianthio cincta, spicam unifloram terminans.

C Y P E R E A E.

Plantae gramineae, culmo enodi, foliis culmum saepe imitantitereti-subulatis. Flores simplicissimi, squamis solis insigsaepe loco perianthii interni setae hypogynae. Fructus simplicissimi, caryopses, nuculae. Semen omnino albuminosum embryo basilari haud evoluto.

223. *ELYNIA* Schrad. (1806.)

Squamae undique imbricatae, uniflorae.

Perianth. duplex, utrumque univalve: exterius maius interius angustius genitalia involvens.

Stylus capillaris, stigmatibus tribus.

Caryopsis nuda.

[Cobresia W.] (Carex Bellardi Schk. Handb. t. 286. D.)

224. *SCHOENUS* L. (1737.)

Squamae in spicam congestae, uniflorae, inferiores vacuae

Perianth. 0.

Stylus simplex, deciduus, stigmatibus duobus aut tribus

Caryopsis nuda, quandoque basi setis suffulta.

(Chaetasporea R. Br. Streblidia Link.)

Sturm. fl. germ.

225. *LEPIDOSPERMA* Labill. (1804.)

Spiculae androgynae, subuniflorae, squamis undique imbricatis, plerisque vacuis.

Stylus simplex, deciduus, stigmatibus tribus.

Squamae hypogynae sex, basi connatae,

Nux ventricosa, obtusa.

Labill. nov. holl. t. 11 — 17.

226. *DULICHIMUM* Pers. (1805.)

Squamae in spicas distichas compositas axillares coagmutatae, vacuae subnullae.

Perianth. 0.

Stylus bifidus longissimus.

Setae hypogynae retrorsum asperae.

Elliott. carolin. austr. t. 2. f. 3.

27. *REYNCHOSPORA* Vahl. (1805.)

Squamae in spicam imbricatae, inferiores vacuae.

Perianth. 0.

Stylus basi persistens, caryopsis late coronans. Stigmata duo.

Setae hypogynae.

(*Schoenus albus* et *fuscus* Sturm. fl. germ. Fl. dan. 320. 1562. *Schoenus surinamensis* Rottb. ic. rar. t. 21. f. 1.)

28. *CAUSTIS* R. Br. (1810.)

Squamae in spicam unifloram congestae, plurimae vacuae.

Perianth. 0.

Stylus basi dilatatus. Stigmata tria.

Setae hypogynae nullae.

Nux ventricosa, basi stylis turgida coronata.

29. *ARTHROSTYLIS* R. Br. (1810.)

Squamae in spicam unifloram congestae, inferiores vacuae.

Flores capitati, involucrati.

Perianth. 0.

Stylus subulatus, trigonus, cum germine articulatus, deciduus. Stigmata tria.

Setae hypogynae nullae.

Nux trigona.

30. *FIMBRISTYLIS* Vahl. (1805.)

Squamae in spicam undique imbricatae, omnes fertiles, dein caducae.

Perianth. 0.

Stylus bifidus, basi compressus, ciliatus.

Setae hypogynae nullae.

Caryopsis basi styli coronata.

Delil. fl. aeg. t. 6. f. 3. Labill. nov. caled. t. 16. [Isoplepis R. Br. *Echinolytrum* Desv.]

31. *DICHROMENA* Mx. (1803.)

Squamae in spicam imbricatae: spicae subcapitatae, involucre lato elongato suffultae.

Perianth. 0.

Stylus bifidus, basi dilatatus.

Setae hypogynae nullae.

Caryopsis lenticularis rugulosa.

(*Schoenus* Radg. guian. t. 16.)

32. *SCIROPUS* Trag. (1551.)

Squamae in spicam undique imbricatae, omnes floriferae.

Perianth. 0.

Stylus bi-trifidus, subdeciduus.

Setae hypogynae aut nullae.

Sturm. fl. germ. [Eleocharis R. Br. *Blysmus* et *Holoschoenus* Link.]

33. *TRICHOPHORUM* Pers. (1805.)

Squamae undique in spicam congestae.

Perianth. 0.

Stylus deciduus.

Setae hypogynae senae capillares, demum elongatae.

Elliott. carol. austr. t. 3. f. 4.

234. *ERIOPHORUM* L. (1737.)

Squamae uniflorae, undique in spicam congestae.

Perianth. 0.

Stylus deciduus.

Lana longissima semen cingens.

Sturm. fl. germ.

235. *ABILDGAARDIA* Vahl. (1805.)

Valvae uniflorae, compressae, mucronatae, in spicam disticham dispositae.

Stylus basi pyramidato-bulbosus. *Stigmata* tria.

Setae nullae.

Caryopsis pyriformis, apice basi styli bulboso-pyramdata coronata.

(*Cyperus Rottb. ic. rar.* t. 13. f. 3.)

236. *CYPERUS* Theophr.

Valvae uniflorae in spicam disticham dispositae. Inferiori nonnunquam vacuae.

Stylus deciduus, bi-vel trifidus.

Setae hypogynae nullae.

Caryopsis nuda, triquetra aut compressa.

Schk. Handb. t. 7. *Engl. bot.* 1309.

237. *PAPYRUS* Theoph. Thuars. (1809.)

Valvae uniflorae in spicam disticham dispositae.

Squamae duae membranaceae, liberae, valvis oppositae.

Setae hypogynae nullae.

Stylus trifidus.

Caryopsis triquetra.

(*Cyperus Rudg. guian.* t. 20.)

238. *MACHAERINA* Vahl. (1805.)

Squamae laxae imbricatae, in spicam paucifloram dispositae.

Cal. bivalvis.

Setae perigynae.

Caryopsis triangularis.

239. *HYPELYTRUM* Rich. (1803.)

Squamae uniflorae, confertissime imbricatae.

Cal. bivalvis, valvis inaequalibus diaphanis.

Setae hypogynae nullae.

Stylus bifidus, trifidus.

Caryopsis trigona.

Pal. Beauv. fl. owar. 2. t. 67. [*Hypaelytrum* Vahl.]

240. *MARISCUS* Vahl. (1805.)

Cal. bivalvis, bi-vel triflorus, spiculam teretiusculam constituens.

Cor. univalves, inferior superiorem inferne amplectens.

Setae hypogynae nullae, vel brevissimae.

Stylus trifidus.

Humb. nov. gen. t. 65 — 67.

241. KYLLINGA Rottböll. (1773.)

Spiculae uniflorae, capitato-congestae, involucratae.

Cal. bivalvis.

Cor. bivalvis.

Stylus bifidus.

Rottb. ic. t. 4. [Thryocephalum Forst. char. gen. 65.]

242. MELANCRANIS Vahl. (1805.)

Squamae in spicam imbricatae. Spicula quinque-septemflora sub singula squama, compressa disticha.

Cal. bivalvis.

Cor. univalvis.

Setae hypogynae nullae.

Stylus bifidus.

243. PUIRENA Rottböll. (1773.)

Squamae undique imbricatae, uniflorae, in spicam dispositae.

Cor. trivalvis, valvis petaloideis unguiculatis patentibus.

Setae hypogynae, petalis interpositae, vel nullae.

Stylus trifidus.

Rottb. ic. t. 19. f. 3. Lam. ill. t. 39. [Vaginaris Rich.]

244. OREOBOLUS R. Br. (1810.)

Spatha bivalvis, uniflora.

Perianthium sexpartitum, cartilagineum, post lapsum nucis persistens.

Stylus deciduus. Stigmata tria.

Nux crustacea.

245. MAPANIA Aubl. (1775.)

Flores capitato-glomerati. Involucrum amplissimum, triphyllum.

Perianthium sexvalve, imbricatum, persistens.

Stylus simplex, stigmatibus tribus.

Aubl. guian. t. 17.

246. REMIREA Aubl. (1775.)

Flores paniculato-congesti.

Cal. bivalvis.

Cor. bivalvis: valva superior post anthesin incrassata, suberosa, nuculam includens.

Aubl. guian. t. 16. [Miegia Schreb.]

GRAMINEAE.

Familia magis perfecta, quam Cypereae, praesertim culmo nodoso, foliis latioribus, duplici perianthio, (externum calycem, internum corollam vocamus). Tertium perianthium, haud constans, lodiculae Palisoto, squamae hypogynae nobis, cum R. Brownio dicitur. Plurimumque fructus est ostryopsis, sed

saepe etiam achenium et nucula. Embryo semper basi est ad albumen, ope scutelli cotyledonei agglutinatus. Dist hic, in systemate artificiali, artificialis.

247. **LYGEUM** Löffling. (1758.)

Flos spathaceus.

Cal. bivalvis, biflorus, inferne cum corollarum basi tubum concrescens.

Cor. univalvis, calyce longior.

Achenium, basi calycis coriacea longe pilosa inclusum.

Rich. in ann. of bot. 2. t. 15. *Schreb.* gram. t. 52. 55

248. **NARDUS** Lobel. (1570.)

Flores spicati unilaterales in rhachi excavata.

Cal. nullus.

Cor. bivalvis, involuta, cum semine connata.

Schreb. gram. t. 7. *Schkuhr.* Handh. t. 11.

249. **LUDOLFIA** Willd. (1808.)

Flores spicati vel paniculati, polygami.

Cal. bivalvis, multiflorus, flosculis brevior.

Cor. herbacea, bivalvis: valva inferior acutissima, ^{superio-} bidentata.

Squamae hypogynae tres.

Stylus tripartitus.

Sem. liberum, maximum.

[*Arundinaria* Mx. *Pal.* *Beauv.* agrost. t. 25. f. 7. *Elliot* carolin. austr. t. 4. f. 4. *Miegia* Pers.]

† 250. **CHUSQUEA** Kunth. (1825.)

Spiculae lanceolato-cylindricae, uniflorae.

Valvae calycinae plures, distiche imbricatae, ^{superio-} florem hermaphroditum involvens.

Squamae hypogynae duae.

Stylus bipartitus.

251. **SPINIFEX** L. (1771.)

Flores polygami dioeci, spicato-fasciculati, involucri feminei in capitulum echinatum congesti.

Cal. biflorus, bivalvis.

Cor. bivalvis, mutica, aecedente saepe neutra univalvi.

Stylus bipartitus.

Pal. *Beauv.* agrost. t. 25. f. 1. *Labill.* nov. holl. t. 230. 231

DIGYNIA.

GRAMINA.

I. CALYCIBUS UNIFLORIS.

A. Flores spicati.

252. **HORDEUM** Plin.

Flores seriatim spicati, saepe polygami, laterales masculae

Involucrum e setis duabus.

- Cal. bivalvis, valva infer. longe aristata.*
Cor. 0.
Squamae hypogynae.
Semen corticatum vel nudum.
E. Viborg von der Gerste. Kopenh. 1802. tab. 1—4.
- 253. PLEURAPHIS** Torr. (1825.)
Flores spicati, polygami, laterales masculi.
Involucrum villosum.
Cal. centralis uniflorus bivalvis, valva altera tri-altera
biaristata.
Cor. bivalvis, valva altera uniaristata.
Calyces laterales biflori, valva infer. dorso aristulata.
Ann. of the lyc. of New-York, 1. t. 10. [Hymenothecium
Lag.]
- 254. OROPETIUM** Trin. (1820.)
Flores spicati in rhacheos scrobiculis.
Cal. univalvis, cartilagineus.
Cor. bivalvis hyalina, pilis stipata; valva inferior ventricosa,
superior plana.
Sem. corolla tectum.
Trin. fund. agrost. t. 1. f. 3. [Monermae spec. Pal. Beauv.
agrost. t. 20. f. 10.]
- 255. PEROTIS** Ait. (1789.)
Flores spicati.
Cal. bivalvis, longissime aristatus.
Cor. bivalvis, minima, mutica.
Squamulae duae hypogynae.
Semen cylindricum, calyce inclusum.
[Xystidium Trin. fund. agrost. t. 1. f. 2.]
- 256. PASPALUM** L. (1759.)
Flosculi subspicati unilaterales ad rhachin planiusculam.
Cal. bivalvis, cartilaginea, mutica vel submucronata: valva
interior convexa exteriorem planiorem amplexens.
Cor. bivalvis, cartilaginea, mutica vel submucronata: valva
interior convexa exteriorem planiorem amplexens.
Squamae hypogynae truncatae.
Semen corticatum.
Humb. nov. gen. t. 23—28. Trin. ic. gram. t. 97—107.
[Axonopodis spec. Pal. Beauv. Cabrera Lag.]
- 257. MICROCHLOA** R. Br. (1810.)
Flosculi spicati unilaterales.
Cal. bivalvis, membranaceus, muticus.
Cor. bivalvis, mutica, villosa.
Humb. nov. gen. t. 22.
- 258. ERIOCHLOA** Kunth. (1815.)
Flosculi spicati subunilaterales. Involucella stellato-pilosa.
Cal. bivalvis, valva inferior subulata.
Cor. bivalvis, calyce minor.

Sem. corolla persistente tectum.

Humb. nov. gen. t. 30.

259. SPARTINA Schreb. (1789.)

Spicae geminae pluresve alternae. Flosculi unilaterales lanceolati, rhachi triquetrae appressi.

Cal. bivalvis, acutissimus, valvis inaequalibus carinatis.

Cor. bivalvis, calyce minor.

Styli connati.

Pal. Beauv. agrost. t. 7. f. 6. Jacqu. ecl. gramm. v. 27.
[*Dactylis* Engl. bot. 380. *Trachynotia* Mx. *Limnetis* Pers.]

260. GYMNOPOGON Pal. Beauv. (1812.)

Flosculi spicati, remoti, alterni: spicae compositae.

Cal. bivalvis, corolla longior.

Cor. bivalvis, valva altera apice bifida, dorso aristata, altera apice integra basi aristata.

Semen cylindricum, nudum.

Pal. Beauv. agrost. t. 9. f. 3.

261. DIGITARIA Heist. Adans. (1762.)

Spicae fasciculato-digitatae, flosculi unilaterales.

Cal. bivalvis, corolla minor.

Cor. bivalvis, oblonga, chartaceo-coriacea: valva inferior mucronata vel setigera superiorem angustiore inhi dens. Rudimentum pedicellatum subclavatum corollae alterius.

Squamae hypogynae.

Semen liberum, vel corticatum.

Pal. Beauv. agrost. t. 10. f. 12. [Cynodon Rich. Syntherisma Walt.]

262. MEROSTACHYS * (1824.)

Flores polygami spicati, unilaterales.

Cal. bivalvis, inaequalis subcoriaceus, valva altera subaristata.

Cor. bivalvis mutica; valva altera villosa alteram involvens. *Antherae* elongatae sessiles.

Squamae hypogynae tres, truncatae.

263. AMPHIPOGON R. Br. (1810.)

Flores spicati, flosculis extremis quandoque sterilibus.

Cal. bivalvis, subaequalis.

Cor. bivalvis: valva utraque aristata, altera tri-altera bifida. *Styli* basi connati.

264. DIPLOPOGON R. Br. (1810.)

Flores spicato-capitati, flosculis extremis sterilibus involucrantibus.

Cal. bivalvis, membrenaceus, aristatus.

Cor. bivalvis: valva exterior apice triaristata, arista intermedia tortili, interior biaristata.

B. Flores racemosi.

265. ZOYSIA Willd. (1801.)

Racemus spicaeformis simplicissimus.

Cal. univalvis, cartilagineus, corollam includens.

Cor. univalvis, membranacea, persistens.

Pal. Beauv. agrost. t. 4. f. 1. [Matrella et Zoysia Pers.

Epiphytis Trin. agrost. t. 1. f. 1.

266. PLEUROPOGON R. Br. (1824.)

Racemus spicaeformis simplicissimus.

Cal. bivalvis, valvis inaequalibus, muticis.

Cor. bivalvis; valva infer. mutica, apice scariosa, superior nervo utrinque lateraliter biseto.

Squamae hypogynae.

Caryopsis libera.

Parry it. app. t. D.

267. KNAFFIA Sm. (1803.)

Racemus spicaeformis simplicissimus.

Cal. bivalvis, truncatus.

Cor. bivalvis pilosa; valva altera latiore apice denticulata, altera angusta, seu obsoleta.

Engl. bot. 1127. Trin. ic. gram. t. 17. [Sturmia Sturm. fl. germ. Chamagrostis Borkhaus. Mibora Adans. Pal. Beauv. agrost. t. 8. f. 4.]

268. SCHIZOPOGON Reichenb. (1827.)

Racemus spicatus simplex.

Cal. 0.

Cor. basi barbata, bivalvis: valvae alterius arista subbasilari tortili.

Rudimentum corollae alterius aristae rectae, medio incrassatae, forma, liberum ad basin corollae perfectae.

269. LYCURUS Kunth. (1815.)

Racemus spicatus. Flores polygami.

Calyces gemini: alter hermaphroditus pedicellatus, alter masculus vel neuter subsessilis. Valva inf. bi- aut tri-superior uniaristata.

Cor. bivalvis, valva infer. aristata.

Sem. liberum.

Humb. nov. gen. t. 45.

270. AEGOPOGON Humb. Bonpl. Willd. (1809.)

Racemus simplex. Flores gemini aut terni, laterales saepe masculi.

Cal. bivalvis, valvis bifidis, medio aristatis.

Cor. bivalvis: valvae apice bidentato-emarginatae: infer. tri-superior biaristata.

Humb. nov. gen. t. 42. 43. Pal. Beauv. agrost. t. 22. f. 3.

271. TRIAENA Kunth. (1815.)

Racemus simplex.

Cal. bivalvis: valva inferior duplo minor, basi aristata.

Cor. bivalvis: valvae inaequales, muticae.
 Rudimentum corollae alterius, arista tricuspis.

Humb. nov. gen. t. 61.

272. *LAPPAGO* Schreb. (1789.)

Racemus subcompositus. Flosculi inversi.

Cal. bivalvis, valvis inaequalibus, superior coriacea, un-
 nato - muricata.

Cor. bivalvis: valvae inaequales, inferiore superiorem te-
 riorem involvente.

Squamae hypogynae.

Caryopsis corolla calyceque tecta.

Schreb. gram. t. 4. Sturm. fl. germ. [Tragus Hall.]

Beauv. agrost. t. 6. f. 13.]

C. Flores paniculati:

a. glomerato - fasciculati.

273. *CORNUCOPIAE* L. (1737.)

Fasciculi florum involucri cyathiformi immersi.

Cal. bivalvis, valvis basi connatis, apice truncatis, muticis.

Cor. univalvis, truncata, mutica, semen obtegens.

Schreb. gram. t. 41.

274. *CRYPsis* Ait. (1789.)

Fasciculi florum spicati vel capitati.

Cal. bivalvis: valvis carinatis, inaequalibus, muticis.

Cor. bivalvis: valvis hyalinis, inaequalibus, muticis.

Sem. liberum.

Sturm. fl. germ. [Schoenus Schreb. gram. t. 32. Heleod.]

Host. gram. 1. t. 29. 30. Antitragus Gärtn.]

275. *ECHINOPOGON* Pal. Beauv. (1812.)

Fasciculi florum capitati.

Cal. bivalvis, valvis lanceolatis, acutis, subaequalibus.

Cor. bivalvis: valva infer. sub apice integro aristata, super.
 apice bidentata.

Rudimentum corollae alterius pedicellatum clavatum.

Germen apice barbatum.

Pal. Beauv. agrost. t. 9. f. 5.

276. *LAGURUS* L. (1737.)

Fasciculi florum subcapitati.

Cal. bivalvis, acuminato - cuspidatus, plumosus.

Cor. bivalvis, calyce minor: valva altera biseta, dorso aris-

Rudimentum alterius corollae cylindricum.

Eugl. bot. 1334. Pal. Beauv. agrost. t. 8. f. 12.

277. *ERIOCHRYSIS* Pal. Beauv. (1812.)

Fasciculi florum paniculato - spicati.

Cal. bivalvis, villosus, coriaceus, muticus.

Cor. calyce minor, bivalvis: valvis inaequalibus mutici

Squamae hypogynae tridentatae.

—en rostratum.

id. Beauv. agrost. t. 4. f. 11.

278. GASTRIDIVM Pal. Beauv. (1812.)

Fasciculi florum paniculato-spicati.

Cal. bivalvis, basi ventricosus, corolla maior.*Cor.* callo insidens, pilis stipata, bivalvis, coriacea: valva inf. tri-quadridentata, sub apice aristata, superior bifido-dentata.*Pal. Beauv.* agrost. t. 6. f. 6. [*Milium lendigerum* Schreb. gram. t. 23. f. 3. Engl. bot. 1107. *Sturm.* fl. germ.]**279. ALOPECURUS** L. (1735.)

Fasciculi florum spicati.

Cal. bivalvis, corollam aequans: valvae subconnatae, muticae vel subulatae.*Cor.* subunivalvis, infra medium aristata.*Styli* connati.*Squamae* hypogynae nullae.*Sturm.* fl. germ. [*Tozzetia* Sav. *Colobachne* Pal. Beauv.]**280. PHLEUM** L. (1735.)

Fasciculi florum spicati.

Cal. bivalvis, corollam superans: valvae naviculares, truncato-mucronatae.*Cor.* bivalvis, truncata, mutica.*Squamae* hypogynae.

Nonnunquam et filum tenuissimum ad basin corollae, quod rudimentum corollae alterius.

Sturm. fl. germ. [*Chilochloa* et *Achnodontium* Pal. Beauv.]**281. PHALARIS** Dioscor.

Fasciculi florum spicati.

Cal. bivalvis, corollam superans: valvae carinatae acutiusculae.*Cor.* bivalvis coriacea.

Rudimentum corollae alterius vel subbivalve, vel penicilli-forme, vel calli specie.

Caryopsis corolla tecta.*Pal. Beauv.* agrost. t. 7. f. 1. *Schrad.* fl. germ. t. 6. f. 5. [*Digraphis* Trin. *Baldingera* fl. wetter.]b. *Flores vere paniculati.***282. POLYPOGON** Desfont. (1800.)

Panicula pyramidalis subcontracta.

Cal. bivalvis, membranaceus, corolla maior: valva utraque apice aristata.*Cor.* bivalvis: valva altera e dorso aristata.*Squamae* hypogynae nullae.*Caryopsis* nuda.*Pal. Beauv.* agrost. t. 6. f. 8. [*Agrostis panicea* Engl. bot. 1704. *Phleum crinitum* Schreb. gram. t. 20. f. 3.]**283. CHAETURUS** Link. (1799.)

Panicula pyramidalis subcontracta, ramis articulatis.

Cal. bivalvis, corolla maior: valva inferior longe subulata.
Cor. bivalvis, subhyalina, mutica: valva superior tripl.
 minor.

Caryopsis libera.

Pal. Beauv. agrost. t. 5. f. 7. [*Polypogon subspicatus* *Panx.*
Ideen, t. 9. f. 1.]

284. *AGROSTIS* L. (1735.)

Panicula varia.

Cal. bivalvis, acutiusculus, muticus, corollam aequans
 vel superans.

Cor. bivalvis, callo saepe insidens aut basi pilis brevibus
 suffulta; valva altera saepe obsoleta aut nulla, altera
 aristata vel mutica.

Squamae hypogynae minimae.

Caryopsis libera.

a. *Vilfa* Adans. (*Sporobolus* R. Br.) corolla mutica bivalvi.

aa. *Colpodium* Trin. R. Br., valvae corollinae obtusae
 scariosae, altera trinervia, nervo medio in setulam
 subexsertam abeunte, lateralibus parallelis. [*Vilfa*
 Trin. ic. gram. t. 57.]

ab. *Phippsia* Trin. R. Br., valvae corollinae obtusae,
 nervis alterius sursum divergentibus. [*Agrostis algida*
Wahlenb. fl. lappon. t. 1. Fl. dan. 1505.]

β. *Trichodium* Mx., valva corollae altera minutissima vel
 nulla, altera mutica. *Pal. Beauv.* agrost. t. 4. f. 8.

γ. *Agraulos* *Pal. Beauv.*, valva corollae altera minutissima
 vel nulla, altera infra medium aristata. *Pal. Beauv.*
 agrost. t. 4. f. 7.

δ. *Apera* Adans. *Pal. Beauv.*, valva corollina altera sub apice
 aristata, altera apice bidentata. Rudimentum saepe co-
 rollae alterius fili aut aristae forma. *Jacqu.* ecl. gram.
 t. 30. *Pal. Beauv.* agrost. t. 7. f. 11. [*Anemagrostis* et
Lachnagrostis Trin.]

285. *PODOSAEUM* DesvauX. (1812.)

Panicula varia.

Cal. bivalvis, corolla subminor; valvae muticae aut mu-
 cronatae.

Cor. bivalvis, subcoriacea: valvae subaequales, inferior
 bifido-dentata, inter dentes aristata.

Pal. Beauv. agrost. t. 8. f. 1—3. *Humb.* nov. gen. 7. t. 681—
 683. [*Tosagris*, *Trichochloa* et *Clomena* *Pal. Beauv.*]

286. *MÜHLENBERGIA* Schreb. (1789.)

Panicula coarctata.

Cal. minutus: valvae inaequales obtusae, dentatae, vel
 acutae integrae.

Cor. callo insidens, basi subpilosa: valva altera ex apice
 aristata, alteram apice bidentatam involvens.

Semen liberum cylindricum apice rostratum.

Schreb. gram. t. 50. 51. *Pal. Beauv.* agrost. t. 7. f. 9. t. 9. f. 3.
 [*Brachyelytrum* *Pal. Beauv.* differt rudimento alterius co-
 rollae pedicelliformi, squamis etiam hypogynis obtusis basi
 ventricosis.]

287. *MILIUM* Varr. L. (1735.)

Panicula varia.

Cal. bivalvis, herbaceus, corolla maior.

Cor. bivalvis, demum cartilaginea, mutica, vel valva altera aristata.

Caryopsis corolla tecta.

Fl. dan. 1143. Schreb. gram. t. 18. f. 2. Host. gram. 3. t. 45. [*Piptatherum* Pal. Beauv. *Urachne* Trin. aristam corollae caducam habet. *Caryochloa* * cur. post., arista incurva; ad hanc pertinent *Milium capense* L., altera e Montevideo, tertia et quarta e Chili, quas reportavit Pöppig. *Helopus* Trin. s. *Oedipachne* Link. pedicellis clavatis insignis.

288. *STREPTACHNE* R. Br. (1810.)

Panicula varia.

Cal. bivalvis, laxus, muticus, vel valvis mucronatis.

Cor. bivalvis: valvae inaequales, altera coriaceo-indurata, apice continuo in aristam, basi subtortilem abiecis, alteram involvens.

Caryopsis oblecta.

Humb. nov. gen. t. 40.

289. *STIPA* L. (1735.)

Panicula ut plurimum effusa.

Cal. bivalvis, membranaceus, herbaceus, corollam aequans aut superans.

Cor. callo insidens, bivalvis: valva altera demum indurata, alteram involvens, apice in aristam basi articulataam tortilem, plerumque longissimam, abiens.

Caryopsis corolla oblecta.

Schk. Handb. t. 16. Broter. phytogr. lus. t. 7. 8. Host. gram. t. 4. f. 34.

Arista in plerisque longissima, vel pilosa aut plumosa, vel nuda. In nonnullis tamen brevior, licet semper flores excedat, basi articulata sit, ac tortilis: e.g. *Stipa Aristella* L., sibirica Lam., *Calamagrostis* Wahlenb., (*Lasiagrostis* Link.) *sinubriata* Kunth., *Jarava* P. B. Huc et *Oryzopsis asperifolia* Mx. (Pal. Beauv. agrost. t. 6. f. 5.) et *O. melanocarpa* Mühlenb. gram. p. 79. seu *Milium racemosum* Sm. in Rees cyclop. cum qua *Stipa virescens* Kunth. in Humb. nov. gen. t. 680. omnino congruit.

290. *CALAMAGROSTIS* Diosc. Adans. (1762.)

Panicula effusa.

Cal. bivalvis, corollam aequans vel superans.

Cor. bivalvis, basi pilis cincta: valvae apice dentatae, altera sub apice aristata.

Caryopsis libera.

Pal. Beauv. agrost. t. 5. f. 9. [Arundo Schrad. fl. germ. t. 4.]

291. *DEYEUXIA* Clarion. Pal. Beauv. (1812.)

Panicula varia.

Cal. bivalvis, membranaceus, corolla longior: valvae subaequales.

Cor. bivalvis, basi pilis cincta: valvae apice dentatae: a tera dorso vel inferne aristata.

Rudimentum corollae alterius filii aut aristae forma, pl-
mosum.

Caryopsis libera.

Pal. Beauv. agrost. t. 9. f. 9. 10. *Humb.* nov. gen. t. 46.

292. *LEERSIA* Sw. (1788.)

Panicula laxa, pedicellis sub flore articulatis.

Cal. herbaceo - coriaceus, compressus, valva infer. nav-
icularis, super. angusta.

Cor. 0.

Squamae hypogynae.

Caryopsis calyce tecta, compressa.

Fl. dan. 1744. [*Asprella* Schreb. *Pal. Beauv.* agrost. t. 4.
f. 2. *Phalaris* Schreb. gram. t. 22. *Homalocenchrus* Mieg.
Hall. Pollich.]

293. *LIMNAS* Trin. (1820.)

Panicula rara, subracemosa, pedicellis sub flore articulata.

Cal. bivalvis, herbaceo - coriaceus, deciduus, corollam
aequans, muticus, vel mucronatus.

Cor. in calycis basi sessilis, bivalvis: valvae inaequales,
inferior latior, e basi aristam tortilem emittens.

Styli connati.

Squamae hypogynae lanceolatae.

Trin. ic. gram. t. 18. [*Pleuroplitis* Trin. fund. agrost. f. 16.]

294. *CYRTOPOGON* Pal. Beauv. (1812.)

Panicula racemosa stricta.

Cal. bivalvis, corollam aequans: valvae acutissimae, sub
apice setigerae.

Cor. bivalvis: valva altera apice bifida: arista inter lacinulas
curvata, geniculata.

Pal. Beauv. agrost. t. 8. f. 7.

295. *ARISTIDA* L. (1741.)

Panicula varia.

Cal. bivalvis, corollam subaequans: valvae acutissimae aut
mucronatae.

Cor. bivalvis: valva infer. demum indurata, apice triari-
stata, arista media nuda vel plumosa: vel arista solitaria
tortili apice tripartita.

Pal. Beauv. agrost. t. 8. f. 10. *Cav.* ic. 5. t. 468 — 471.

Desfont. fl. atlant. t. 21. 35. *Delil.* flor. aegypt. t. 13.

Labill. nov. Caled. t. 17. [*Chaetaria* Pal. Beauv. agrost.
t. 8. f. 5. 6. *Arthratherum* Pal. Beauv. agrost. t. 8. f. 8. 9.]

296. *PENTAPOGON* R. Br. (1810.)

Panicula racemosa.

Cal. bivalvis, valvis aequalibus muticis.

Cor. pedicellata, bivalvis: valva inf. apice quadriseta, inter
setas dentata, arista media tortili, longiore: valva su-
perior minor, apice bidentata.

Stigmata sessilia.

Pal. Beauv. agrost. t. 8. f. 11. [Agrostis quadrifida Labill. nov. holl. 1. t. 22.]

297. *ANISOPOGON* R. Br. (1810.)

Panicula laxa.

Cal. bivalvis, corolla maior.

Cor. bivalvis: valva inferior apice triaristata, media longiore, tortili: valva superior angustior, bifido-dentata.

Rudimentum corollae alterius pedicellatum clavulatum, clavula setigera.

Pal. Beauv. agrost. t. 9. f. 8. [Danthonia Trin. ic. gram. t. 61.]

II. CALYCIBUS SESQUIFLORIS.

Praeter corollam perfectam, altera univalvis neutra.

298. *MELINIS* Pal. Beauv. (1810.)

Panicula effusa.

Cal. bivalvis: valvae inaequales, infer. minutissima integra, superior emarginata, mucronata.

Cor. perfecta bivalvis, mutica, subcoriacea: valva inferior subbidentata.

Cor. imperfecta univalvis, herbacea, apice bidentata, arista inter dentes longissima.

Pal. Beauv. agrost. t. 11. f. 4. [Tristegis Nees in hor. physic. herol. t. 7. Suardia Schrank. hort. monac. 2. t. 58.]

299. *SACCHARUM* L. (1737.)

Panicula effusa, articulis bifloris, flosculo altero sessili, altero pedicellato.

Cal. lanugine involucratus, bivalvis.

Cor. perfecta bivalvis mutica; valva inf. interdum obsoleta, nonnunquam aristata.

Cor. imperfecta univalvis hyalina.

Squamae hypogynae.

Caryopsis libera.

Trin. fundam. agrost. f. 15. Jacq. ecl. gram. t. 5. Roxb. corom. 3. t. 232. Tussac fl. antill. 1. t. 23. [Imperata Cyrill. ic. rar. 2. t. 11. Erianthus Mx. Ripidium Trin.]

300. *ANDROPOGON* L. (1737.)

Flores spicati, racemosi vel paniculati; polygami.

Cal. ♂ uniflorus, bivalvis, muticus, basi pilosus, pedicellatus, hermaphrodito adstans: corolla bivalvis mutica.

Cal. ♀ biflorus, sessilis, muticus, basi pilosus.

Cor. perfecta interior bivalvis, neutra exterior univalvis, utraque aristata.

Squamae hypogynae dentatae.

Caryopsis libera.

Schkuhr Handb. t. 342. [Heteropogon Pal. Beauv. agrost. t. 23. f. 8. Chrysopogon, Centrophorum Trin. fund. agrost. f. 5. Rhaphis Lour. Trin. ic. gram. t. 8. 9.]

301. *ANATHERUM* Pal. Beauv. (1812.)

Flores paniculati, vel spicati, polygami.

Cal. ♂ uniflorus, bivalvis, pedicellatus, hermaphrodito astantans: corolla bivalvis mutica.*Cal.* ♀ biflorus, sessilis.*Cor.* perfecta interior bivalvis, neutra exterior univalvis utraque mutica.*Squamae* hypogynae truncatae, denticulatae.*Pal. Beauv.* agrost. t. 22. f. 10. 11. [*Elionurus* Kunth. t. 11. f. 7. Humb. nov. gen. t. 62. 63.]302. *POLLINIA* * (1815.)

Flores paniculati vel subspicati, polygami.

Cal. ♂ uniflorus, bivalvis, aristatus, basi pilosus. *Cor.* bivalvis mutica.*Cal.* ♀ biflorus, sessilis, aristatus, basi pilosus.*Cor.* perfecta bivalvis, valva altera sub apice aristata. *Cor.* neutra univalvis mutica.[*Andropogon* Schk. Handb. t. 342. a. *Host.* gram. 2. t. 1. *Pogonatherum* Pal. Beauv. agrost. t. 11. f. 7. *Diectoma* Kunth. in Humb. nov. gen. t. 64.]303. *CYMBOPOGON* * (1815.)

Flores subpaniculati polygami. Involucrum foliaceum flores verticillato - aggregatos, hermaphroditos et masculos, colligens.

Cal. ♂ uniflorus, bivalvis, muticus, basi pilosus. *Cor.* uni-bivalvis mutica.*Cal.* ♀ biflorus, bivalvis, muticus, basi pilosus. *Cor.* perfecta bivalvis, valva altera aristata. *Cor.* neutra univalvis mutica.[*Andropogon cymbarius* in act. acad. petropol. a. 1807 et 1808. tom. 2. t. 9.]304. *ANTHESTERIA* L. fil. (1779.)Fasciculi florum paniculati, involucrati. Flores quaterni aut seni, verticillati, masculi perfectum ambeuntes, sessiles vel subpedicellati. *Cal.* univalvis, uniflorus. *Cor.* bivalvis, mutica, hyalina.

Flos perfectus centralis stipitatus.

Cal. bivalvis, biflorus, chartaceus.*Cor.* perfecta hyalina, bivalvis: valva inferior plerumque apice longissime aristata, rarius mutica. *Cor.* neutra, univalvis, membranacea.*Pal. Beauv.* agrost. t. 23. f. 7. *Desfont.* fl. atlant. t. 254. *Gärtn.* t. 175.305. *SORGUM* Michel. (1729.)

Flores paniculati, polygami.

Cal. ♂ pedicellatus, uniflorus, bivalvis. *Cor.* bivalvis, mutica.*Cal.* ♀ cartilagineus, clausus, bivalvis, biflorus.

Cor. perfecta bivalvis, valva altera aristata. *Cor. neutra* univalvis mutica.

[*Holcus Arduin.* act. acad. patav. 1. t. 1—4. *Sorghum Jacq.* fil. ecl. gram. t. 18.]

306. *DIMERIA* R. Br. (1810.)

Flores racemoso-spicati.

Cal. subcoriaceus, bivalvis, muticus.

Cor. perfecta valva infer. bifida, arista media tortili elongata.

Cor. neutra univalvis mutica.

Caryopsis cylindrica.

Trin. fund. agrost. f. 11. [*Arthraxon Pal. Beauv. agrost. t. 11. f. 6.*]

307. *THRASSYA* Kunth. (1815.)

Flores spicati unilaterales in rhachi dilatata.

Cal. biflorus, flosculos superans, bivalvis: valva inferior bipartita, infra apicem arista incurva.

Cor. perfecta bivalvis mutica.

Cor. imperfecta univalvis, infra apicem aristata.

Humb. nov. gen. t. 30.

308. *PANICUM* Plin.

Flores racemoso-spicati vel paniculati.

Cal. biflorus, bivalvis, valva altera plerumque minore.

Cor. perfecta bivalvis mutica, demum indurata.

Cor. neutra univalvis, pariter mutica.

Sem. corticatum.

Pal. Beauv. agrost. t. 10. f. 1—5. Host. gram. 2. t. 20.

[*Isachne*, *Monachno Pal. Beauv. Streptostachys Desv.*

Thalassium * cur. post.]

309. *SETARIA* Pal. Beauv. (1812.)

Panicula spicata. Involucrum bi-vel multisetum.

Cal. biflorus, bivalvis, valva altera plerumque minore.

Cor. perfecta bivalvis mutica, persistens.

Cor. neutra uni-bivalvis mutica.

Pal. Beauv. agrost. t. 13. f. 3. [*Panicum Engl. bot. t. 871.*

875. *Urochloa Pal. Beauv. agrost. t. 11. f. 1.*]

310. *ORTHOPOGON* R. Br. (1810.)

Spica composita.

Cal. biflorus, bivalvis: valvae subaequales, inaequaliter aristatae.

Cor. perfecta bivalvis, mutica, demum indurata.

Cor. imperfecta uni-bivalvis, mutica.

Caryopsis corticata.

[*Panicum Fl. dan. 1564. Echinochloa Pal. Beauv. agrost.*

t. 11. f. 2. Oplismenus Ej. t. 11. f. 3.]

311. *PENNISSETUM* Rich. Pers. (1801.)

Flores spicati aut spicato-racemosi. Involucrum multisetum, uni-biflorum.

Cal. biflorus, bivalvis, muticus.

Cor. perfecta bivalvis, mutica, demum indurata.

Cor. imperfecta uni-bivalvis, mutica.

Caryopsis corticata.

Pal. Beauv. agrost. t. 13. f. 5. *Humb.* nov. gen. t. 3.
 [*Gymnothrix Pal. Beauv.* agrost. t. 13. f. 6. *Humb.* nov.
 gen. t. 678. *Penicillaria Sw.* Willd. *Pal. Beauv.* t. 13. f. 1.
Jacqu. ecl. gram. t. 17.]

312. *CENCHRUS* L. (1737.)

Racemus spicatus. Involucrum connivens, multipartitum
 plerumque spinosum, demum induratum, uni-quadriflorum.

Cal. chartaceus vel coriaceus, biflorus, bivalvis, mutica.

Cor. perfecta bivalvis, mutica, demum indurata.

Cor. imperfecta univalvis, mutica.

Styli basi connati.

Achenium calyce tectum.

Pal. Beauv. agrost. t. 13. f. 7. *Schreb.* gram. t. 23. f. 1. t. 34.
 [*Antheophora Schreb.* gram. t. 44. *Colladoa Pers.*]

III. CALYCIBUS BI-TRIFLORIS.

A. *Corolla altera perfecta, altera mascula aut neutra bivalvi.*313. *CTENIUM* Panz. (1813.)

Flores spicati unilaterales, biseriales.

Cal. bivalvis, subtriflorus: valva altera minima, altera dorso setam patentem rigidam gerens.

Cor. perfectae valvae hyalinae, nervosae, altera apice fissa, altera apice aristata.

Cor. mascula bivalvis mutica, cum rudimento tertiae corollae pedicellatae, clavatae.

Act. acad. monach. tom. 4. t. 13. [*Chloris monostachya Mx.*
Campulosus Pal. Beauv. agr. t. 13. f. 1. *Monocera aromatica Elliott.*]

314. *CHLORIS* Sw. (1788.)

Flores spicati unilaterales.

Cal. bivalvis, bi-triflorus: valva exterior paullo maior, mutica vel aristata.

Cor. perfecta sessilis inferior: valva inferior obovata a ephippiiformis, aristata vel mutica, superiorem fovens.

Cor. imperfecta mascula vel neutra pedicellata, truncata, uni-bivalvis, mutica vel aristata. Saepe praeter masculam corollam bivalvem tertia adest neutra corolla univalvis.

Schk. Handb. t. 343. Panzer in act. acad. monach. 4. t. 10. f. 2. [*Rhabdochloa et Leptochloa Pal. Beauv.* *Eustachys Desv.* *Tetrapogon Desf. fl. atl. t. 255.*]

315. *ATHEROPOGON* Mühlenb. Willd. (1809.)

Flores spicati vel racemosi.

Cal. biflorus, bivalvis; valvae inaequales muticae.

Cor. perfecta bivalvis: valva inferior apice bidentata: aut triaristata, arista media longiore: superior bidentata.

Cor. imperfecta, vel mascula bivalvis, valva altera trisetâ, vel neutra univalvis pedicellata aristata, vel demum eius rudimentum, arista divisa.

Caryopsis libera subrostrata.

Jacqu. ecl. gram. t. 7. 12. [Dinebra Kunth. triplicem corollae imperfectae formam habet: masculae bivalvis, valva altera trisetâ, in *D. repente*, *Humb.* nov. gen. t. 52.: tum neutrius univalvem trisetam, in *D. bromoide*, ib. t. 51.: tum solam aristam tridentem, in *D. chondrosioides*, ib. t. 53. et in *D. aristidoide*, ib. t. 695. Haec aristae tridentis forma in *Chondrosio* Pal. Beauv. recurrit, sed valva altera corollae perfectae proxime sub arista dentes binos et paulo inferius itidem dentes aristaeformes, itaque quaternos habet dentes. *Humb.* l. c. t. 56 — 59. Nec multum distat *Pentarrhaphis* Kunth. ib. t. 60. Corollae masculae enim valva altera, praeter aristam terminalem duas laterales habet et dentes aristae adstantes fissos. *Heterostegia* Desv. Kunth. l. c. t. 54. cum *Chondrosio* iterum sub eodem genere comprehendenda est. Nec absimilis est *Polyodon* Kunth. l. c. t. 55., licet hic arista tridens deficiat. *Tristhera* Desv. Pal. Beauv. agr. t. 9. f. 4. ad *Dinebrae* aristoidis et chondrosioidis formas proxime accedit: nisi quod arista haud tridens, sed ternae aristae basi iunctae adaint. *Bouteloua* Lag. *Dinebrae* primam et secundam formam ostendit.]

116. *THUAREA* Pers. (1805.)

Spica vaginata, secunda: rhachis dilatata coriacea.

Cal. bivalvis, valvis subaequalibus, muticis.

Flores polygami et masculi.

Cor. utraque bivalvis, mutica.

Achenium corolla et rhachi indurata obtectum.

Pal. Beauv. agrost. t. 22. f. 9.

17. *ECHINOCHLAENA* Desf. (1813.)

Spica unilateralis.

Cal. bivalvis, coriaceus, valvis inaequalibus acuminato-subulatis.

Cor. perfecta coriacea, valvis obtusis.

Cor. imperfecta mascula bivalvis, valvis acutis membranaceis.

Humb. nov. gen. t. 38. [*Echinolaena polystachya* Kunth. l. c. t. 679. non huius loci, sed potius ad *Coelachnen* R. Br. pertinere videtur.]

118. *ROTTBÖLLA* L. fil. (1779.)

Spica articulata, articulis uni-bifloris: flosculi scrobiculis rhacheos immersis.

Cal. biflorus, saepe geminus, masculus et androgynus, uni-aut bivalvis.

Cor. perfecta bivalvis, hyalina, mutica.

Cor. mascula vel neutra uni-aut bivalvis, hyalina, mutica.

Caryopsis rostrata.

Host. gram. 1. t. 24. Fl. dan. 938. *Delil.* fl. aeg. t. 14. f. 1.

Labill. nov. Caled. t. 20. [*Ophiurus* Gärtn. R. Br. arti-

bulis unifloris, Lepturus R. Br. praeter articulos unius
 etiam calyce univalvi, Hemarthria R. Br. spica comprae-
 semiarticulata; Lodicularia Pal. Beauv. agrost. t. 21.
 squamis hypogynis trilobis insignibus gaudet. Phlois
 Trin. quandoque corolla mascula destituitur.]

319. XEROCHLOA R. Br. (1810.)

Spiculae binae, quaternae, abbreviatae, pauciflorae,
 vaginis conditae,
Cal. bivalvis, biflorus, scrobiculis rhacheos semiimmersa
 valva exteriore minore.
Cor. exterior ♂, interior ♀, utraque exserta, bivalvis
 mutica, subulata.
Styli basi connati.
Squamae hypogynae nullae.
Caryopsis valva interiore corollae inclusa.

320. ISCHAEUM L. (1764.)

Spica rhachi articulata.
Cal. gemini, pedicellatus alter masculus vel neuter, al-
 sessilis biflorus, valvis coriaceis.
Cor. perfecta bivalvis, subhyalina, valva inferior bidentata
 aristata.
Cor. mascula bivalvis, hyalina, mutica.
Caryopsis oblecta.

Pal. Beauv. agrost. t. 21. f. 4. *Lam.* ill. t. 839. f. 4. *Salis-*
ic. t. 1. [*Colladoa distachya* Cav. *ic.* t. 460. *Meoschium* P.B.]

321. THELEPOGON Roth. (1821.)

Spica rhachi articulata. Involucrum monophyllum appres-
 sum, cartilagineum.
Cal. bivalvis, triflorus, muticus: valvae inaequales papil-
 losae, exterior apice bidentata, interior scrobiculo rha-
 cheos immersa.
Cor. perfecta, mutica, hyalina.
Cor. duae laterales masculae, valvis exterioribus infra me-
 dium arista geniculata tortili instructis.

322. TRIPSACUM Mill. (1738.)

Spica rhachi articulata, floribus monoecis.
Calyces ♂ superiores bivalves, biflori.
Cor. utraque ♂ bivalvis mutica.
Calyces ♀ inferiores bivalves biflori, rhacheos articuli
 sinuatis incrassatis immersi.
Cor. ♀ bivalvis, mutica: neutra univalvis.
Squamae hypogynae hyalinae.
Achenium.

Pal. Beauv. agrost. t. 22. f. 1. *Willd.* hort. berol. t. 1.

323. MANISURIS L. fil. (1779.)

Flores spicati.
Calyces coriacei, bivalves, imbricati, perfectorum et mascu-
 lorum florum sibi oppositi. Calycis perfecti valva exterio-

marginata: masculi biflori valva exterior apice dorsi appendicula membranacea aucta.

Cor. utraque hyalia, bivalvis, mutica: ad masculam accedit corolla secunda neutra.

Schreb. gram. t. 43. Palis. Beauv. agrost. t. 21. f. 10. [Peltophorus Pal. Beauv. agrost. t. 21. f. 11.]

21. ICHNANTHUS Pal. Beauv. (1812.)

Flores paniculati.

Cal. bivalvis subtriflorus: valva inferior latior apice bidentata, inter dentes mucronata.

Cor. perfecta bivalvis, cartilaginea, reliquis contraria.

Cor. mascula altera bivalvis, neutra altera univalvis utraque mutica.

Squamae hypogynae truncatae, emarginatae.

Pal. Beauv. agrost. t. 12. f. 1.

22. HIEROCHLOE J. G. Gmelin. (1747.)

Flores paniculati.

Cal. bivalvis, subtriflorus, muticus, corollas aequans aut superans: valvae ovatae nervosae.

Cor. perfecta media, valvis ovatis.

Cor. masculae binae laterales, perfectae similes, sed valva infer. apice mucronata vel inferne aristata, valva super. bifida.

Squamae hypogynae suborbiculatae, vel lanceolatae.

[Holcus Schk. Handb. t. 314. Fl. dan. 963. 1508. Sturm fl. germ. Disarrhenum Labill. nov. holl. t. 232. Toresia H. et P. Pal. Beauv.]

23. HOLCUS L. (1737.)

Flores paniculati.

Cal. bivalvis, biflorus, muticus.

Cor. perfecta bivalvis, mutica.

Cor. mascula bivalvis, sub apice aut dorso aristata.

Caryopsis libera.

Engl. bot. 813. 1169. 1170. Utramque corollam perfectam observavit Pal. Beauv. Equidem in H. molli saepe vidi corollam perfectam aristatam, masculam muticam.

27. ACRATHERUM Link. (1827.)

Panicula.

Cal. biflorus, bivalvis, valvis nervosis inaequalibus.

Cor. perfecta basi pilosa, bivalvis, valva infer. arista torta.

Cor. neutra bivalvis, mutica.

[Goldbachia Trin.]

28. AFLUDA L. (1742.)

Flores semiverticillati, paniculati, polygami. Spiculae binae spatha involucretae, sessilis altera; altera crasse pedunculata.

Spiculae sessilis involucrellum cartilagineum, bipartitum, lacinia altera apice bidentata. Flores subhorizontales.

Cal. bivalvis, biflorus. *Cor.* ♂ bivalvis, valva inf. apice bifido-aristata. *Cor.* ♂ bivalvis, mutica.

Spiculae pedicellatae involucellum cartilagineum, bivalv uniflorum. *Cal.* bivalvis, muticus. *Cor.* ♂ similis.

Caryopsis tecta.

Schreb. gram. t. 42. [Calamina Pal. Beauv. t. 23. f. 1.]

329. *CHRYSURUS* Pers. (1805.)

Panicula racemosa secunda.

Spiculae fertiles subbinæ cinctae spiculis sterilibus (masculis aut neutris) multifloris.

Cal. ♂ membranaceus, biflorus.

Cor. Perfecta pedicellata, bivalvis, valva infer. aristata.

Cor. mascula vel neutra, longius pedicellata, bivalvis (interdum univalvis) aristata.

Spiculae steriles corollae bivalves muticae obtusae, vel aristatae, vel cuspidatae.

Caryopsis tecta.

Pal. Beauv. t. 22. f. 5. Cynosurus Host. gram. 3. t. 4. Eng bot. t. 1333.

330. *MELICA* L. (1737.)

Panicula racemosa.

Cal. laxus, bivalvis, coloratus vel scariosus, flosculi aequans, superans, vel parum brevior, subtriflorus.

Cor. perfecta una aut altera, valvis muticis, membranaceo-cartilagineis, muticis.

Cor. abortiens terminalis, una aut altera, bivalvis, aut univalvis, mutica, subpedicellata.

Caryopsis corolla tecta.

Pal. Beauv. agrost. t. 14. f. 4. 5. Host. gram. 2. t. 10—1

331. *COELACHNE* R. Br. (1810.)

Panicula angusta.

Cal. biflorus, bivalvis; valvis subaequalibus obtusissimi muticis.

Cor. perfecta inferior, bivalvis, valvis muticis, exte-
ventricosa.

Cor. superior pedicellata, feminea.

Caryopsis libera.

B. *Corolla utraque, ac, si tertia affuerit, et haec perfecta.*

332. *AIRA* L. (1735.)

Panicula.

Cal. bivalvis, biflorus.

Cor. bivalvis, mutica vel aristata.

Caryopsis libera.

α. *Airopsis* Desv. calyce flosculos superante, valva corollae infer. apice tridentata. *Pal. Beauv. t. 14. f. 8. Aira glabra Desv. in Journ. de bot. 1. t. 7.*

β. *Catabrosa* Pal. Beauv. calyce flosculis breviori, valva corollae infer. truncata, apice erosa. *Pal. Beauv. t. 19. f.*

7. *Dupontia* R. Br. calyce corollisque scariosis, muticis.
8. *Periballia* Trin. ramis paniculae inferioribus sterilibus verticillatis coloratis, flosculis inferioribus muticis, superioribus aristatis, arista dorsali recta (Aira involucreta Cav. ic. 1. t. 44.)
9. *Deschampsia* Pal. Beauv. valva corollae infer. basi aristata, arista recta. *Pal. Beauv. agrost. t. 18. f. 3.* [Campella Link.]
10. *Uralpis* Nutt. calyce flosculos superante, valva corollae infer. apice tricuspidata, brevissime aristata. [Airochloa Link.]
11. *Corynephorus* P. B. arista corollae basilari clavata, medio articulata, barbata. *Pal. Beauv. agrost. t. 18. f. 2.* (Aira canescens Schk. Handb. t. 12.)

ERIACHNE R. Br. (1810.)

Flores racemosi, vel paniculati.

Cal. bivalvis, valvis cuspidatis flosculos subaequantibus, biflorus.

Corollae basi barbatae aut undique villosae, valva exteriore vel mutica vel ex apice aristata.

Caryopsis corolla tecta.

PENTAMERIS Pal. Beauv. (1812.)

Flores paniculati.

Cal. bivalvis, flosculos superans.

Cor. bivalvis, valva inferior nervosa, apice in quatuor dentes setigeros et aristam tortilem abiciens: valva superior multo minor, bidentata.

Squamae hypogynae emarginatae.

Caryopsis libera, umbilicata, stella lanuginosa coronata.

Pal. Beauv. t. 18. f. 8.

MOLINIA Schrank: (1789.)

Flores paniculati.

Cal. bivalvis, subtriflorus, flosculis multo brevior.

Cor. bivalvis, valvis inaequalibus acutis.

Squamae hypogynae obovatae.

Caryopsis libera, apice binucronulata.

Palis. Beauv. agr. t. 14. f. 6. [Melica Host. gram. 2. t. 8. Enodium Gaud.]

BECKMANNIA Host. (1805.)

Flores racemoso-paniculati unilaterales.

Cal. biflorus, valvis navicularibus obtusis.

Cor. bivalvis, valva infer. rostrato-mucronata.

Squamae hypogynae lanceolatae.

Caryopsis libera.

Palis. Beauv. agr. t. 19. f. 6. Host. gram. 3. t. 6. [Joachimia Tenor. fl. neap. t. 5.]

SECALE Plin.

Flores spicati.

Cal. bivalvis, biflorus.

Cor. bivalvis: valva infer. aristata.

Caryopsis libera.

Pal. Beauv. agr. t. 20. f. 6. Host. gram. 2. t. 48. 4. f. 12.

338. DINEBA Delil (1814.)

Spicae unilaterales racemosae.

Cal. bi-triflorus, bivalvis, flosculis longior: valvae aminatae.

Cor. bivalvis: valva inf. apice emarginata, breviss. mucronata.

Caryopsis libera.

Delile fl. aegypt. t. 11. f. 3. Pal. Beauv. agr. t. 16. f. 1. [Cynosurus Link. Cav. ic. 1. t. 91. Wangenheimia Moench]

IV. CALYCIBUS MULTIFLORIS.

A. Flores spicati.

339. TRITICUM Varr.

Spica vel racemus spicatus.

Cal. bi-multiflorus: valvae muticae vel mucronatae.

Cor. bivalvis: valvae muticae vel aristatae.

Caryopsis nuda vel Achenium corolla et calyce indur.

Metzger cereal. t. 1 — 34. [Agropyron Pal. Beauv., Elygia Desv., Arthrostachys Link: comprehendit Trib. agrestia, rhachi articulata. Brachypodium Pal. Beauv. (Trachynia, Catopodium, Brizopyrum Link.) eas species quae spicas gerunt compositas subracemosas.]

340. LOLIUM Virg.

Flores spicati.

Cal. univalvis, multiflorus.

Cor. bivalvis; valva inferior mutica vel aristata.

Achenium corolla involutum.

Schk. Handb. t. 18.

341. ELYMUS Theophr. L. (1735.)

Flores spicati.

Calyces bivalves, plerumque gemini, interdum subtriflori, raro fasciculati, bi- aut multiflori, involucrum mentes: valvae flosculos aequantes aut superantes; muticae subulatae aut aristatae.

Cor. bivalvis, mutica vel aristata.

Achenium corolla obvolutum.

Pal. Beauv. agrost. t. 20. f. 7. Jacq. ecl. gram. t. 37. [Asprella Cav., Gymnostichum Schreb. gram. t. 43. Elymus Hystrix L., ubi plerumque calyces abortiunt, callos abeuntes, sed in solo fertili vidi iterum evanescentes et calyces prodire.]

342. AEGILOPS Theophr. L. (1735.)

Flores spicati, rhachi insculpti.

Cal. triflorus, bivalvis, cartilagineus: valvae apice quadriaristatae.

Cor. bivalvis: valva infer. mutica aut aristis 1. 2. 3, superior conduplicata. Corollae summae saepe masculae.

Achenium corolla tectum.

Schk. Handb. t. 344. Hist. gram. 2. t. 5—7.

341. POMMEREULA L. fil. (1779.)

Spica vaginata.

Calyces multiflori, alterni, bivalves: valvae cuspidatae.

Cor. bivalvis: valva inferior apice quadricidentata, dentibus setigeris, arista intermedia terminali flexuosa.

Styli connati.

Roxb. corom. 2. t. 131.

342. HILARIA Kunth. (1815.)

Flores spicati polygami.

Calyx (s. involucreum) irregulariter sexpartitum, laciniis obtusis muticis et aristatis, lanceolatis etiam apiceque bidentatis.

Spiculae ternae, laterales ♂ sexflorae, intermedia ♀, 1-florae.

Cor. (s. calyx) ♂ bivalvis: valvae subaequales, oblongae, obtusae muticae.

Cor. ♀ bivalvis: valva altera linearis subspathulata obtusa, altera linearis acuta, utraque mutica.

Caryopsis involucreo inclusa.

Humb. nov. gen. t. 37.

343. GAUDINIA Pal. Beauv. (1812.)

Spica laxa, spiculis alternis.

Cal. bivalvis, multiflorus, valvis obtusiusculis muticis.

Cor. bivalvis: valva inf. bidentata, dorso supra medium aristata, arista subtorta: valv. super. apice bi-quadridentata.

Achenium corolla involutum.

Pal. Beauv. agrost. t. 19. f. 5.

344. ELEUSINE Gärtn. (1788.)

Spica simplex vel composita, vel racemoso-paniculata.

Spiculae subunilaterales, 3-10-florae.

Cal. bivalvis, muticus vel mucronatus aut subulatus.

Cor. bivalvis: valva infer. plerumque bidentata, aristata vel mutica.

Caryopsis corolla laxè tecta.

Panzer in act. acad. monach. 4. t. 8. f. 1. Gärtn. t. 1.

[*Rhachochloa* et *Leptochloa* Pal. Beauv. agr. t. 15. f. 1.

t. 17. f. 3. *Dactyloctenium* Willd. *Oxyadenia* Nuttall.]

47. CYNOSURUS L. (1735.)

Spica racemosa, seu racemus spicatus. *Spiculae* subsecundae, involucrelo pinnato suffultae.

Cal. muticus, bivalvis.

Cor. bivalvis, valva inferior longior, cuspidata.

Caryopsis nuda, subrostrata.

Sturm. fl. germ.

348. *TRICHAETA* Pal. Beauv. (1812.)

Spica capitata.

Cal. subtriflorus, bivalvis: valvae hispidae, acutae.

Cor. bivalvis: valva infer. bifida, sub apice seta
cato-flexuosa: valva super. bifido-dentata.

*Pal. Beauv. agr. t. 17. f. 8. [Bromus ovatus Cav.
t. 591. f. 2.]*

349. *STREPTOGYNE* Pal. Beauv. (1812.)

Spica subracemosa stricta.

Cal. bivalvis: valvae inaequales, superior apice mucr

Cor. bivalvis: valva infer. bidentata, inter dentes se
super. subtruncata emarginata.

Germen apice barbatum.

Stigmata retrorsum hispida, siccitate tortilia.

Pal. Beauv. agrost. t. 16. f. 8.

B. Flores glomerato-paniculato-spicati.

350. *KÖLERIA* Pers. (1805.)

Panícula glomerato-spicata.

Cal. bivalvis, 3-6-florus: valvae inaequales acut
mucronatae.

Cor. bivalvis: valva infer. sub apice integro se
super. bifida.

Pal. Beauv. agr. t. 17. f. 1. [Aeluropus Trin.]

351. *SESSLERIA* Scop. (1760.)

Flöres glomerato-spicati vel subcapitati.

Cal. bivalvis, membranaceus, bi-quinqueflorus:
acuminatae.

Cor. bivalvis: valva infer. apice tri-quinquedentata,
intermedio subaristato.

Styli basi coaliti.

Caryopsis tecta.

*Pal. Beauv. agr. t. 16. f. 7. Host gram. 2. t. 76. 97
vol. 4. t. 22. [Echinaria Desf. Pal. Beauv. agr. t.
Psilathera et Oreochloa Link.]*

352. *ELYTROPHORUS* Pal. Beauv. (1812.)

Glomeruli florum spicati. *Involucrum* glomeruloru
lyphyllum, foliolis lanceolatis.

Cal. tri-sexflorus, bivalvis: valvae acuminatae.

Cor. bivalvis: valva inf. navicularis, subulata, super
emarginata, mucronato-denticulata.

Pal. Beauv. agr. t. 14. f. 2. [Echinalysium Trin.]

353. *PAPPOPHORUM* Schreb. (1789.)

Panícula spicata.

Cal. bivalvis, membranaceus, subtriflorus, floscul
quans: valvae inaequales, muticae.

Cor. bivalvis: valva inferior latior, ventricosa, apice aristis parallelis subdenis obsessa.

Caryopsis libera.

Pal Beauv. t. 16. f. 11. [Enneapogon Desvauz.]

C. Flores paniculati.

84. TRIBHAPHS R. Br. (1810.)

Panicula effusa vel coarctata.

Cal. bivalvis, multiflorus: valvae aequales muticae.

Corollae inferiores perfectae, superiores masculae vel neutrae. Valva exterior apice triaristata, aristis rectis, interior mutica.

Squamulae hypogynae.

85. ECTROSLA R. Br. (1810.)

Panicula subcoarctata.

Cal. bivalvis, multiflorus: valvae aequales muticae.

Cor. infima perfecta, reliquae masculae vel neutrae.

Valva exterior arista terminali simplici, quae longior in masculis neutrisque.

86. ZEUGITES Patr. Brown. (1756.)

Panicula rara.

Cal. (s. involucrem) bivalvis triflorus: valva infer. latior truncata.

Cor. duae masculae pedicellatae, altera supra alteram: valvae aequales, obtusiusculae, muticae.

Cor. feminea sessilis inferior, univalvis: valva scariosa aristata.

P. Brown jamaic. t. 4. f. 3.

87. AVENA Varr.

Panicula effusa aut racemosa.

Cal. bivalvis, membranaceus, longitudine varius, bi-multiflorus.

Cor. plerumque basi barbata; valva inferior apice bifida, dentibus saepe setigeris, arista media tortili.

Squaminae hypogynae.

Caryopsis tecta, vel nuda.

Hist. gram. 2. t. 56. 58. 59. vol. 3. t. 43. 44.

a. Danthonia Cand. calycibus flosculos aequantibus vel superantibus, arista terminali, dentibus lateralibus setigeris aut aristatis. *Jos. Monti prodr. bonon. t. 2. f. 1. Trin. ic. gram. t. 49 — 54. 62 — 68.*

β. Trisetum Pers. valva infer. apice biseta, arista supra medium dorso affixa, tortili aut flexuosa. *Pal. Beauv. agrost. t. 18. f. 1.*

γ. Rostraria Trin. floribus minutis, calyce flosculis breviori, arista dorsali minus tortili. [*Avena Sav. fl. pis. t. 1. f. 4.*]

δ. Triplasis Pal. Beauv. valvae infer. aristis lateralibus terminalibus mediaque basilari rectiusculis. *Pal. Beauv. t. 16. f. 10.*

358. ARUNDO Varr.

Panicula patula.

Cal. multiflorus, bivalvis, valvis subaequalibus flosculo aequantibus.

Cor. bivalvis, callo insidens, basi pilis cincta: valva infer apice mucronato - dentata, sub apice setigera.

Caryopsis tecta.

Host. gram. 4. t. 38. [Donax *Pal. Beauv.*, Trichoon *Host*
Scolochloa Mert. et Koch., Phragmites et Ampelodesmos
Link]

359. GYNERIUM Bonpl. Pal. Beauv. (1812.)

Gramen dioecum, paniculatum.

Cal. bivalvis, uni - bi - multiflorus. Valvae inaequales in mascula flosculis breviores, in feminea longiores.

Cor. basi longe pilosa, in bifloris altera sessilis, altera pedicellata. Valvae inaequales, superior apice bidentata

Squamae hypogynae truncatae, dentato - ciliatae.

Caryopsis bicornis.

Pal. Beauv. agrost. t. 24. f. 6. *Humb.* pl. aequin. 2. t. 115.

360. BROMUS Theophr. L. (1737.)

Panicula varia, spiculis compressis.

Cal. bivalvis multiflorus.

Cor. bivalvis; valva infer. apice emarginata aut bidentata, arista sub apice prodiens; valva super. ciliata.

Host. gram. 1. t. 16 — 19. [Libertia *Lejeun.* in nov. act.
nat. cur. 12. t. 65.]

361. FESTUCA Dodon. (1557.)

Panicula varia, spiculis compressis.

Cal. bivalvis multiflorus.

Cor. bivalvis: valva infer. apice aristata.

Host. gram. 2. t. 78. 81 — 91. vol. 3. t. 20. vol. 4. t. 60.

[*Dactylis L.*, *Schedonorus Pal. Beauv.*, *Schoenodorus et Vulpia Link.*]

362. CALOTHECA Pal. Beauv. (1812.)

Panicula simpliciuscula.

Cal. bivalvis, multiflorus, flosculis brevior.

Cor. bivalvis: valva inf. margine dilatato, sub apice mucronata vel aristulata; superior apice appendiculata.

Achenium corolla tectum.

Pal. Beauv. agrost. t. 17. f. 6. 7. [*Chasacolytrum Desv.*]

363. CENTOTHECA Desv. (1812.)

Panicula varia.

Cal. bivalvis, flosculis brevior, tri - quadriflorus.

Cor. bivalvis: valva infer. mucronata, marginibus muricato - setosa.

Pal. Beauv. agr. t. 14. f. 7.

364. CERATOCHLOA Pal. Beauv. (1812.)

Panicula effusa, spiculis compressis.

Cal. bivalvis, multiflorus.

Cor. bivalvis: valva infer. apice bidentata, inter dentes mucronata.

Achenium corolla tectum, bi-tricorne.

Pal. Beauv. t. 15. f. 7.

365. DIPLAGHNE Pal. Beauv. (1812.)

Panicula effusa.

Cal. bivalvis, multiflorus: valva super. apice mucronata.

Cor. bivalvis: valva infer. apice bidentata, inter dentes setigera, superior subtruncata, emarginata.

Caryopsis libera.

Pal. Beauv. agr. t. 16. f. 9.

366. TRIODIA R. Br. (1810.)

Panicula varia.

Cal. bivalvis, tri-quinqueflorus, flosculos subaequans.

Cor. bivalvis: valva inf. apice tridentata.

Caryopsis libera.

Jacqu. ecl. gram. t. 16. [Graphephorina Desv. Pal. Beauv.

Windsoria Nuttall., Trideus Link.]

367. BRIZA Theophr. L. (1737.)

Panicula varia, spiculis cordato-ovatis.

Cal. multiflorus, valvis obtusis.

Cor. valva inf. ventricosa cordata, super. subreversa, utraque mutica.

Sturm. fl. germ.

368. POA L. (1737.)

Panicula varia, spiculis ovatis vel lanceolatis.

Cal. tri-multiflorus, bivalvis, flosculis brevior.

Cor. bivalvis, mutica.

Caryopsis libera.

Sturm. fl. germ.

a. *Sclerochloa* Pal. Beauv. t. 19. f. 4. flosculis obtusis.

β. *Sphanopus* Trin. pedicellis cuneiformibus. (Poa divaricata Gouan. ill. t. 2. f. 1.)

γ. *Hydrochloa* Hartw. valvis corollae multinerviis. (Poa aquatica Host. gram. 2. t. 60. Fl. dan. 920.)

δ. *Eragrostis* Pal. Beauv. agr. t. 14. f. 11. valva corollae super. reflexa ciliata, stylis in caryopse persistentibus.

ε. *Megastachya* Pal. Beauv. agr. t. 15. f. 5. valva corollae infer. emarginata, submucronata.

ζ. *Orthoclada* Pal. Beauv. agr. t. 14. f. 9. ramis paniculae strictis apice floriferis, flosculis acutis, germinibus rostellatis.

369. GLYCERIA R. Br. (1810.)

Panicula effusa, spiculis teretibus.

Cal. bivalvis, flosculis longe brevior; valvis obtusis, margine hyalinis.

Cor. bivalvis: valva infer. sub-croca, utraque obtusa, margine hyalina.

Squamae hypogynae obcordatae coalitae.

Caryopsis libera.

Pal. Beauv. agr. t. 19. f. 7.

370. **SCHISMUS** Pal. Beauv. (1812.)

Panicula coarctata.

Cal. bivalvis, flosculos subaequans, valvis membranaceis obtusis.

Cor. bivalvis, valva inf. apice obcordato emarginata, setula intermedia, margine hyalina.

Pal. Beauv. t. 15. f. 4. [Electra Panzer in act. acad. monach. 4. t. 9. f. 3.]

371. **UNFOLA** L. (1737.)

Panicula varia, spiculis ancipitibus.

Cal. bivalvis, muticus, multiflorus.

Cor. infima univalvis abortiens. Reliquae perfectae, valvis navicularibus, acutis, mucronatis.

Caryopsis stylorum basi persistente coronata.

Pal. Beauv. agr. t. 15. f. 6. [Chasmanthium Link.]

TRIGYNIA.

GRAMINEAE.

372. **PSAMMA** Pal. Beauv. (1812.)

Flores fasciculato-spicati.

Cal. bivalvis, uniflorus, corolla longior: valvae carinatae, inaequales.

Cor. basi pilosa, bivalvis: valvae sub apice aristolatae.

Squamae hypogynae lanceolatae, germine longiores.

Pal. Beauv. agr. t. 6. f. 1. [Ammophila Host. grain. 4. t. 41.]

373. **CHAMAERHAPHIS** R. Br. (1810.)

Flores spicati. Rhachis articulata: aristae longissimae rectae infra flosculos ex ipsa rhachi prodeunt.

Cal. bivalvis, biflorus; valvae inaequales.

Cor. bivalvis, exterior ♂, interior ♀, utraque conformis mutica.

Squamae hypogynae.

Achenium corolla cartilaginea tectum.

RESTIEAE.

374. **ELEGIA** L. (1771.)

Flores dioeci, amantacel. Amenta bracteis s. spathis suffulta, ♂ laxa subsimplicia, ♀ in racemo composito.

Perianthium hexaphyllum: foliola ♂ setacea, ♀ lanceolata.

Caps. trilocularis.

[Rest. o Rottb. ic. t. 3. f. 4.]

75. LEPYRODIA R. Br. (1810.)

Flores dioeci, vel hermaphroditi.

Perianthium bracteatum, sexvalve, utriusque sexui conforme.

Antherae simplices, peltatae.

Caps. trilocularis, trisperma.

[*Calorophus* Labill. nov. holl. t. 278.]

76. ANARTHRIA R. B. (1810.)

Flores dioeci in spica composita, bracteati.

Perianthium sexvalve subaequale.

Antherae didymae, utrinque bifidae.

Caps. trilocularis, trisperma.

† 77. GALURUS * (1825.)

Flores dioeci, spicati.

Perianthium tripartitum, utriusque sexus conforme.

Styli tres, laciniato-multifidi.

Caps. trilocca.

Tricoccae. [*Acalypha* N. L. Burm. ind. t. 61. f. 2. *Caturus* L. mant. 1. 1767. Forte ad *Acalypham* pertinet.]

78. HOLOSTEUM Dioscor.

Cal. pentasepalus.

Cor. pentapetala, petalis denticulatis.

Caps. unilocularis, polysperma, apice in sex dentes dehiscens.

Caryophylleae. Schk. Handb. t. 20. Gärtn. t. 130.

79. LECHEA L. (1751.)

Cal. trisepalus, bracteis binis suffultus.

Pet. tria lanceolata.

Stamina numero varia, ternarium tamen vel duplicatum servantia.

Caps. trivalvis, valvis medio septiferis. Sem. pauca.

Cisteae. Gärtn. t. 129. [*Gaura* Lam. ill. t. 281. f. 3.]

80. MONTIA Michel. (1729.)

Cal. disepalus.

Cor. pentapetaloides, lobis duobus minoribus.

Stam. epipetala.

Stigmata tria sessilia.

Caps. trivalvis, trisperma.

Portulacaeae. Michel. nov. gen. t. 13. Fl. dan. 131. 1926.

Engl. bot. 1208. Schk. Handb. t. 20. Sturm. fl. germ.

81. POLYCARPON Löffl. (1758.)

Cal. pentasepalus, sepalis mucronatis, margine membranaceis.

Petala quinque emarginata.

Stamina 3. 5.

Styli brevissimi.

Caps. unilocularis, trivalvis polysperma.

Paronychieae. Engl. bot. 1031. Gärtn. t. 129.

382. MINUARTA Löff. (1758.)
Cal. pentasepalus.
Pet. quinque, calyce multo minora.
Stamina ex annulo perigyno, tria, quinque.
Styli tres, filiformes.
Caps. unilocularis, trivalvis, polysperma.
Paronychieae? *Caryophylleae?* Cav. anhl. de
nat. 2. t. 21.
383. TILLAEA Michel. (1729.)
Cal. tripartitus.
Pet. tria.
Filamenta tria sterilia hypogyna.
Capsulae tres dispermae, intus dehiscentes.
Sedeac. Michel. gen. t. 20. Reichenb. ic. t. 191.
384. KÖNIGIA L. (1767.)
Cal. tripartitus corollinus.
Stam. filamentis capillaribus, calyce brevioribus, an
subrotundis.
Stigmata tria sessilia.
Caryopsis nuda, in fundo calycis.
Fl. dan. 418. Gärtn. t. 128.
385. DONATIA Forst. (1775.)
Cal. trisepalus, sepalis acuminatis.
Petala octona aut dena, oblongo-linearia, calyce
longiora.
Filamenta tria subulata. *Antherae* didymae.
Styli tres filiformes.
Caps. infera trilocularis?
Saxifrageae? Forst. char. gen. t. 5.
386. PROSERPINACA L. (1741.)
Cal. superus tubulosus, limbo tripartito.
Cor. 0.
Stigmata tria sessilia.
Nux triquetra, trilocularis, trisperma. Embryo et
linearis in albumine.
Haloragaeae? R. Br. Lam. ill. t. 50. Juss. in ann. s
t. 30. f. 1. [Trixis Gärtn. t. 24.]
387. HOUTTUYNIA Thunb. (1784.)
Spatha tetraphylla, patula, corollam referens.
Spadix oblongus, genitalibus nudis tectus.
Stamina angulis germinis inserta. *Antherae* obli-
biloculares.
Stigmata tria sessilia.
Caps. unilocularis, polysperma. Placentatio pari
Embryo basilaris, non evolutus.
Hook. in Bot. magaz. t. 2731.

CLASSIS IV.
T E T R A N D R I A.

CONSPECTUS

CLASSIS QUARTAE.

M O N O G Y N A E.

† Floribus perfectis.

I. Corolla monopetala.

A. Fructibus superis n. 388 — 428.

B. Fructibus inferis.

Vellutina et Aggregatae 429 — 436.

Rubrae.

a. Fructibus mono-dispermis, 437 — 466.

b. Fructibus tetraspermis, 467 — 470.

c. Fructibus polyspermis 471 — 481.

II. Corolla pleiopetala.

A. Fructibus superis, 482 — 498.

B. Fructibus inferis, 499 — 508.

†† Floribus imperfectis.

I. Fructibus superis.

509 — 542.

543 — 561.

II. Fructibus inferis, 562 — 570.

D I G Y N A E.

I. Floribus perfectis, 571 — 578.

II. Floribus imperfectis 579 — 582.

T R I G Y N A E.

584 — 586.

T E T R A G Y N A E.

587 — 590.

M O N O G Y N I A.

† Flores perfecti.

I. COROLLA MONOPETALA.

A. *Fructus superi.*

GLOBULARIA Clus. (1601.)

Flores aggregati in capitulo globoso. Receptaculum
paleaceum. Involucrum imbricatum.

Cal. quinquefidus persistens.

Cor. quinquefida irregularis.

Stigma simplex.

Caryopsis calyce laxo obvoluta. Embryo inversus.

Globulariaceae. Ann. des scienc. nat. 9. t. 40. 41.
ill. t. 56. Gärtn. t. 44. Sturm. fl. germ.

389. *STILBE* Berg. (1767.)

Flores polygami, capitati, spicati. Receptaculum paucum. Involucrum nullum.

Calyx duplex, exterior paleis receptaculis ternis quemvis flosculum constituitur: interior monophyllus, rigidus, quinquefidus.

Cor. infundibuliformis, fauce villis clausa, limbo quinquepartito erecto, laciniis linearibus.

Stamina intra lacinias corollae inserta, subulata, antheris didymis.

Stigma simplex.

Caryopsis calyce interno tecta.

Globulariaceae. Berg. fl. cap. t. 4. f. 1. Lam. ill. t. 4.

390. *MATTUSCHKEA* Schreb. (1789.)

Flores fasciculato-capitati. Receptaculum subpaleis e bracteis cuivis pedicello adstantibus.

Cal. simplex, quadripartitus, persistens.

Cor. subinfundibuliformis; tubo gracili, limbo quadrilobato aequali.

Stamina intra lacinias corollae inserta; antheris didymis. *Stylus* filiformis, stigmate simplici.

Caryopses binae aut quaternae intra calycem.

Globulariaceae. [Perama Aubl. guian. t. 18.]

391. *ALLIONIA* Löff. (1758.)

Involucrum subtriflorum, mono- aut triphyllum.

Calyx corollinus tubulosus, limbo quadrilobato, basi persistens.

Nectarium urceolatum staminiferum.

Nucula embryone peripherico, albumine centrali, calyce tecta.

Nyctagineae. Lam. ill. t. 58. Herit. stirp. t. 31. Gd. t. 214.

V E R B E N E A E.

392. *CALLICARPA* L. (1741.)

Cal. quadrifidus.

Cor. subcampanulata limbo, quadrilobato.

Stamina exserta.

Bacca tetrasperma.

Lam. ill. t. 69. Gärtn. t. 94. Hook. exot. flor. t. 133. [Triphyra Lour.]

393. *AEGIPHILA* Jacq. (1767.)

Cal. quadridentatus.

Cor. tubulosa, subhypocrateriformis, limbo quadrilobato.

Stamina exserta.

Stylus bifidus.

Drupa quadrilocularis, tetrasperma.

Lam. ill. t. 70. Vahl. eclog. t. 10. [Manaboa Aubl. guian. t. 23—25.]

184. *PEITITIA* Jacq. (1763.)

Cal. quadridentatus.

Cor. infundibuliformis, limbo quadrifido.

Stamina exserta.

Stylus simplex, stigmate capitato.

Drupa monoppyrena, nucleo biloculari.

Jacqu. amer. t. 182. f. 6.

185. *WALLENIA* Sw. (1788.)

Cal. quadrilobus.

Cor. tubulosa, limbo quadrilobo erecto subconnivente.

Stam. filamentis e fundo corollae, exserta, antheris erectis.

Stylus subulatus, brevis.

Bacca monosperma.

Swartz. fl. ind. occid. 1. t. 6.

186. *PELYOZUS* Lour. (1790.)

Cal. tubulosus quadridentatus, demum cum fructu coalescens.

Cor. tubo brevi, limbo quadrifido reflexo.

Stamina fauci inserta, corolla breviora, antheris erectis.

Stylus filiformis, corolla longior, stigmate emarginato.

Bacca disperma.

187. *BUCHIA* Kunth. (1815.)

Cal. bipartitus.

Cor. subinfundibuliformis, limbo quadripartito aequali.

Stam. fauci corollae inserta, filamentis brevissimis, antheris bilocularibus.

Stylus filiformis, stigmate tripartito.

Fructus chartaceus trilocularis, loculis monospermis.

Humb. nov. gen. t. 132.

188. *CHILIANTHUS* Burchell (1824.)

Cal. quadrifidus.

Cor. tubulosa, quadrifida.

Stam. corollae inserta.

Caps. unilocularis, quadrivalvis, tetrasperma.

[*Scoparia arborea* Thuub.]

189. *LABATIA* Sw. (1788.)

Cal. quadripartitus, laciniis exterioribus maioribus.

Cor. tubuloso-urceolata, limbo erecto quadrilobo. Squamulae inter lobulos.

Stamina fundo corollae inserta, eius laciniis opposita, inclusa: antheris bilocularibus.

Stylus subulatus, stigmate simplici.

Fructus baccatus quadrilocularis, tetraspermus. *Semina* testa ossea: cotyledones crassae, albumen subnolle.
Sapotaceae. Sw. fl. ind. occ. 1. t. 6. Mart. nov. gen. t. 161. 162.

400. *POUTERIA* Aubl. (1775.)

Cal. quadripartitus aequalis, persistens.

Cor. campanulato-urceolata, limbo quadridentato; *petala* setae inter denticulos.

Stamina fundo corollae inserta, inclusa, antheris didymis, dorso adnatis.

Stylus quadrangularis, stigmate quadripartito.

Caps. unilocularis, quadrialvis, tetrasperma. *Semina* angulata, membrana tenui obducta.

Stryaceae. Aubl. Guian. t. 33.

401. *PHELLINE* Labill. (1824.)

Cal. quadridentatus, persistens.

Cor. rotata, quadripartita, patula.

Stamina ad basin corollae inserta, corollam subaequant filamentis brevibus; antheris oblongis didymis, erectis.

Stylus brevis subtubulosus, ore quadridentato.

Capsula corticata, quadrilocularis, loculis suberosis, tetraspermis.

Stryaceae? Labill. nov. Galed. t. 38.

402. *NÜXIA* Commers. Juss. (1789.)

Cal. subcampanulatus, quadrifidus, persistens.

Cor. tubulosa, limbo quadrifido, reflexo.

Stamina inter lacinias limbi inserta, exserta, antheris didymis.

Stylus simplex, stigmate truncato, simplicissimo.

Caps. polysperma.

Personatae. Lam. ill. t. 71.

403. *BUDDLEA* L. (1737.)

Cal. quadrifidus, aequalis.

Cor. campanulata, tubulosa, limbo quadrifido regulari.

Stam. fauci corollae inserta, inclusa.

Stigma clavatum.

Caps. bilocularis, bivalvis, valvis bifidis: placenta centrali demum libera, polysperma.

Personatae. Gärtn. t. 49. Humb. nov. gen. t. 182—1 Hook. in Bot. mag. 2713.

404. *SCOPARIA* L. (1742.)

Cal. quadripartitus aequalis.

Cor. rotata, quadripartita: faux pilis erectis coronata.

Stigma obtusum.

Caps. bilocularis, bivalvis, valvis subbifidis, placenta centrali demum libera.

Personatae. Lam. ill. t. 85.

93. *CENTUNCULUS* Dillen. (1719.)

Cal. quadripartitus, persistens.

Cor. supera, calyce minor, subrotata, quadrifida, persistens.

Stamina tubo corollae inserta.

Stylus brevis, stigmate truncato.

Caps. globosa, circumscissa, polysperma. Placenta centralis globosa.

Primulaeae. Fl. dan. 177. 1567. *Schk.* Handb. t. 24.

94. *PLANTAGO* Plin.

Flores spicati, capitati.

Cal. quadrifidus.

Cor. supera, rotata, quadripartita, persistens.

Stam. elongata.

Stylus simplicissimus.

Caps. circumscissa, bilocularis, di-tetrasperma.

Plantagineae. Lam. ill. t. 85. *Sturm.* fl. germ. *Gärtn.* t. 51.

95. *SANGUISORBA* Fuchs. (1542.)

Flores spicati.

Cal. caepalus.

Cor. supera, subrotata, quadrifida, persistens.

Stam. corollam aequantia vel superantia, antheris didymis.

Stylus brevis, stigmate quadrilido.

Drupa exsucca, quadrangula, submonosperma.

Rosaceae. Engl. bot. 1312. [*Pimpinella* *Gärtn.* t. 32.]

96. *LEUCONOTIS* Jack. (1823.)

Cal. quadripartitus.

Cor. tubulosa, basi gibba, sursum coarctata, limbo quadrilobo.

Stam. tubo corollae inserta, inclusa, antheris sagittatis.

Stylus brevis, stigmate annulato.

Becca mono-trisperma.

Contortae.

97. *LASIOSTOMA* Schreb. (1789.)

Cal. quinquepartitus, brevissimus, basi squamulis suffultus.

Cor. infundibuliformis, intus villosa, limbo quadrifido.

Stamina tubo inserta, antheris oblongis erectis.

Stylus corolla longior, stigmate obtuso.

Caps. corticata unilocularis, disperma.

Contortae. [Rouhamon *Aubl.* guian. t. 36.]

G E N T I A N E A E.

10. *EXACUM* L. (1747.)

Cal. quadripartitus, persistens.

Cor. tubo globoso, limbo quadrifido.

Stam. tubo inserta, longitudine limbi. Antherae subrotundae, longitudinaliter dehiscentes.

Stylus declinatus, stigmate simplici.

Caps. compressa, bilocularis, polysperma. *Pl.*
utrinque dissepimento adnata, demum libera.

Smith, ic. pict. 3. t. 18. *Gärtn.* t. 114. [*Gentiana* Fl. dan

411. *ORTHOSTEMON* R. Br. (1810.)

Cal. tubulosus quadridentatus.

Cor. limbo brevi quadripartito, fauce nuda.

Stam. corollae inserta: antherae longitudinaliter dehisc

Stylus simplex, stigmatibus binis capitatis.

Caps. semibilocularis, polysperma.

412. *CALLOPISMA* Mart. (1826.)

Cal. campanulatus, quadripartitus, laciniis carinatis.

Cor. hypocrateriformis, tubo aequali cylindrico, fauce
limbo quadripartito.

Stam. tubo inserta. Antherae birimosae, apice
duplici.

Stigma exsertum, bilobum.

Caps. unilocularis, polysperma, bipartibilis. *Valva*
flexae placentiferae.

Mart. nov. gen. 2. t. 183. 184. [*Deianira Schlecht.*]

413. *COUTOUBEA* Aubl. (1775.)

Cal. subcampanulatus, quadripartitus, laciniis acumi

Cor. hypocrateriformis, limbo patente quadrifido.

Stamina tubo inserta, filamentis ad basin appendic

Antherae longitudinaliter dehiscentes.

Stylus simplex, stigmate bilobo.

Caps. semibilocularis, polysperma.

Aubl. guian. t. 27. 28. *Mart.* nov. gen. 2. t. 185. [*Pi*
Schreb.]

414. *PLADERA* Roxb. (1821.)

Cal. cylindricus, aut quadrangularis, inaequaliter qu
dentatus.

Cor. infundibuliformis, limbo irregulari.

Stam. unicum maximum: Antherae longitudinaliter
hiscentes, post anthesin non mutatae.

Stylus stigmate bilobo.

Caps. semibilocularis, polysperma.

[*Hoppea* Willd. *Hayne* term. bot. t. 30. f. 3. *Ca*
Lam.]

415. *MITRASACME* Labill. (1804.)

Cal. angulatus, quadri- (raro bi-) fidus.

Cor. tubo angulato, limbo quadripartito, aequali.

Stam. aequalia, plerumque inclusa, antheris sagit
posticis.

Stylus basi bifidus, apice simplex, stigmate capitati

Caps. semibilocularis, inter styli brachiola dehiscens.

Labill. nov. holl. t. 49.

STYBAEA Soland. R. Br. (1810.)*Cal.* ventricosus, quadripartitus, carinatus, angulis alatis.*Cor.* quadrifida, marcescens.*Stam.* exserta. Antherae longitudinaliter dehiscentes, post anthesin apice calloso recurvae, vel poro dehiscentes.*Stylus* stigmatibus duobus.*Caps.* semibilocularis, valvis margine inflexis: placenta demum libera.

[Schultesia Mart. nov. gen. 2. t. 180 — 182.]

HOUSTONIA Gronov. L. (1737.)*Cal.* quadrifidus, persistens.*Cor.* infundibuliformis, tubo elongato, limbo quadrilobo.*Stam.* tubo corollae inserta, brevia.*Stylus* brevissimus, stigmatibus bifido acuto.*Caps.* didyma, bilocularis, valvis dissepimento contrariis: placenta in dissepimento: semina pauca.

Lam. Ill. t. 79. Bot. mag. t. 370. 2822. Gärtn. t. 49.

POLYPREMUM L. (1741.)*Cal.* tetrasepalus.*Cor.* rotata, quadrifida: fauce barbata.*Stam.* brevissima.*Stylus* brevis truncatus.*Caps.* bilocularis, polysperma: dissepimento valvis contrariis. Placenta demum libera.

Gärt. t. 62.

ANDREWSIA * (1817.)*Cal.* quadrifidus appressus.*Cor.* campanulata, quadrifida, persistens.*Stigma* incrassatum, bilobum.*Caps.* unilocularis bivalvis, valvis inflexis placentiferis.

[Centaurella Mx. fl. bor. amer. t. 42. Bartonis Mühlenb.]

FRASERA Walt. (1788.)*Cal.* quadripartitus.*Cor.* quadripartita patula. Lacinias medias occupat glandula orbicularis barbata.*Caps.* unilocularis, bivalvis, valvis inflexis placentiferis.

Semina alata, imbricata.

SALVADORA Garcin. (1749.)*Cal.* minutus, quadrilobus.*Cor.* quadrifida, laciniis revolutis.*Stam.* cum laciniis corollae alterna.*Stigma* simplex.*Bacca* monosperma.

Chenopodeae. Fahl. symb. 1. t. 4. Roxb. corom. 1. t. 26

BLAIRIA L. (1737.)*Cal.* quadrifidus.*Cor.* subcampanulata, quadrifida.

Stam. receptaculo imposita, plerumque exserta: an-
tibilocularibus muticis (rarius aristatis).

Stylus simplex, exsertus, stigmate obtuso.

Caps. quadrilocularis, quadrangula, angulis dehiscens
Ericaceae. Wendl. collect. 1. t. 25. 31. 2. t. 38. 43. 49

† 423. *SYMPLEZA* Lichtenst. (1817.)

Cal. turbinatus, compressus, bifidus.

Cor. tubulosa, subbifida.

Stam. tubo inserta, exserta, antheris bilocularibus,
ticis.

Stylus simplex, stigmate obtuso.

Ericaceae. [Blaeria bracteata Wendl. collect. 2. t. 37.]

424. *GARDNERIA* Wallich. (1821.)

Cal. quadrifidus.

Cor. subrotata, coriacea, quadrifida; laciniae altern
cum calycis laciniiis.

Stam. brevissima, basi corollae inserta; antherae
culares.

Stylus filiformis.

Bacca bilocularis. *Semina* duo urceolata. *Embryo*
vatus evolutus in albumine corneo.

425. *PENAEA* L. (1737.)

Cal. disepalus, seu bracteae deciduae coloratae.

Cor. campanulata, limbo brevi, quadrifido.

Stam. summae corollae inter lacinias inserta: filam
brevissima: antherae biloculares.

Stylus quadrangulus, stigmate quadrilobo.

Caps. quadrilocularis, quadrivalvis: valvae medio s
ferae: loculi dispermi.

— — *Hook.* in Bot. mag. 2809. *Gärtn.* t. 225. T
in Berl. mag. 1. t. 3.

426. *ROUSSEA* Sm. (1789.)

Cal. quadri-(quinque-) partitus, coriaceus, persistens

Cor. subampullaeformis, limbo quadri-(quinque-) pa
reflexo.

Stamina (4, 5) conniventia, exserta: antherae eri
sagittatae.

Stylus tetra-(penta-)gonus, stigmate peltato.

Bacca ovato-pyramidata, in stylum attenuata, qu
(quinque-)locularis. *Placentae* crassae, e centr
euntes. *Sem.* numerosa.

Solanaceae? Smith ic. ined, 1. t. 6. *Lam.* ill. t. 75. C
t. 212.

427. *CYLINDRIA* Lour. (1790.)

Cal. tubulosus, brevis, persistens, quadrifidus.

Cor. tubulosa, quadrifida, laciniiis cucullatis tubum
tinuantibus.

Stam. Antherae biloculares, cucullis laciniarum corollae inclusae. Filamenta subnulla.

Stylus brevissimus: stigma quadrifidum.

Bacca exsucca, monosperma. Sem. lanuginosum.

— — (*Blimbingum* *Humph. umbon.* 4. t. 73.)

48. *Asagoa* Humb. (1818.)

Cal. tetra-pentasepalus, persistens.

Cr. hypocrateriformis, fauce villosa, limbo quadrifido regulari.

Stam. corollae inter lacinias inserta, erecta, exserta: antherae cordatae.

Stylus simplex, stigmatе globoso.

Caps. bilocularis, quadrivalvis, loculis tetraspermis. Sem. superposita, peltata, alata. Embryo rectus in albumine.

— — *Humb. nov. gen.* 3. t. 216. 217.

B. *Fructus inferi.*

V A L E R I A N E A E.

49. *Patrinia* Juss. (1810.)

Cal. quadri-quinquedentatus, persistens, bracteae ovali membranaceae laterali adnatus.

Cr. campanulata, regularis, quinqueloba.

Caps. trilocularis, loculis duobus abortientibus, calyce coronata, bractea laterali adnata.

Reichenb. hort. 1. t. 20. [*Fedia sibirica* *Gärtn.* t. 86.]

50. *Opercularia* Gärtn. (1788.)

Involucrum monophyllum, campanulatum, tri-sexflorum, dentatum.

Receptaculum supra planum, inferne in dissepimenta angulata divisum.

Cr. supera, infundibuliformis, quadri-quinqueloba.

Stamina receptaculo inserta.

Stylus filiformis, stigmatе bilobo.

Semina receptaculo immersa, angulata, dissepimentis sciuncta.

Gärtn. t. 24. *Annal. du mus.* 4. t. 70. 71. *Labill. nov. holl.* t. 46 — 48.

A G G R E G A T A E.

Familia Compositis finitima, floribus in eodem receptaculo congestis, involucri seu anthodii, staminibus epipetalis et fructu, caryopse vel achenio potius pappo coronatis. Sed differunt numero quaternario et libero positu staminum. maxime etiam situ embryonis inverso, albumineque consumito.

II. *Dipsacus* Plin.

Involucrum polyphyllum persistens.

- Receptaculum* conicum, paleis subpungentibus obscuris.
Cal. proprius obsoletus.
Cor. tubulosa, quadrifida.
Achenium coronula urceolata terminatum.
Schk. Handb. t. 21. *Jacqu.* ecl. 2. t. 106. 107. *Gärtn.* t. 8
432. **SCABIOSA** Seb. Vaill. (1722.)
Involucrum subduplici serie.
Receptaculum paleaceo-setaceum.
Cor. tubulosa, quadrifida, subradians.
Achenium tetraëdram, paleis simplicibus aut setis
 ronatum.
Tournef. inst. t. 263. Fl. dan. 447.
433. **SUCCISA** Seb. Vaill. (1722.)
Involucrum imbricatum, in paleas receptaculi continuae
Cor. tubulosa, quadrifida, aequalis aut radians.
Achenium corona paleaceo-setosa terminatum.
 [Scabiosa leucantha *Gärtn.* t. 86. *Kitab.* pl. hung. 1. t.
 tom. 3. t. 230.]
434. **ASTEROCEPHALUS** Seb. Vaill. (1722.)
Involucrum simplex, multipartitum vel polyphyllum.
Receptaculum paleaceum.
Corolla tubulosa, quinquefida, radians.
Achenium duplici corona, scariosa externa et setis
 quinis interioribus composita.
Vaillant in act. acad. paris. 1722. tab. 13. f. 18. 20. *Stu-*
ranus Entwickel. des Embr. t. 5. f. 82—86.
435. **PTEROCEPHALUS** Seb. Vaill. (1722.)
Involucrum simplex, multipartitum vel polyphyllum.
Receptaculum setosum.
Achenium corona plumosa.
Vaill. l. c. f. 23. 24. *Gärtn.* t. 86.
436. **KNAUTIA** L. (1737.)
Involucrum simplex, polyphyllum, erectum, subdece
 florum.
Receptaculum nudiusculum.
Cor. inaequalis, quadrifida, lacinia exteriore maiore.
Achenium corona paleaceo-setacea.
Schk. Handb. t. 22. *Gärtn.* t. 86.

RUBIACEAE.

*Plantae herbaceae, fruticosae, arboreae, foliis oppositis aut
 ticiatis, stipulis intra- aut extrapetiolaribus, corollis
 rumque monopetalis, fructu semper infero, semine albumin
 embryo erecto.*

a. *Fructibus mono-dispermis.*

437. **CEPHALANTHUS** L. (1737.)
Receptaculum commune subglobosum, setosum, flos
 los plures in capitulum colligens.

Involucrum nullum.

Cal. proprius quadrifidus.

Cor. infundibuliformis, aequalis, quadrifida.

Stam. tubo inserta, corolla breviora.

Stylus simplex, corolla longior, stigmate globoso.

Cap. calyce persistente coronata, inverse pyramidata, quadrilocularis, loculis duobus abortientibus. *Semina* angula angulata, inferne attenuata, apice apophysi callosa.

Scht. Handb. t. 21. Gärtn. t. 86.

40. **MITCHELLA** L. (1751.)

Flores bini iidem germi insidentes.

Cal. quadridentatus, brevis.

Cor. infundibuliformis: limbo quadripartito, aequali, intus limbo.

Stam. intra tubum, filamentis brevissimis.

Stylus exsertus, stigmate quadripartito.

Baccas binae coadunatae, depressae, calyce coronatae.

Sam. quaterna.

Lam. ill. t. 63. Gärtn. t. 192.

41. **ACRODEYON** * (1825.)

Receptaculum commune subglobosum villosum, flosculos plures in capitulum colligens.

Involucrum nullum.

Cal. proprius quadrifidus.

Cor. infundibuliformis quadrifida.

Stam. brevissima, ad incisuras corollae.

Bacculae monospermae in baccam compositam succulentam confluentes.

[*Cephalanthus occidentalis* et *angustifolius* Lour.]

42. **DECLIEUXIA** Humb. Kunth. (1818.)

Cal. quadrifidus.

Cor. infundibuliformis, limbo quadrifido regulari, fauce barbata.

Samina fauci inserta, aut inter divisuras limbi posita.

Antherae lineares, versatiles, dorso affixae.

Stylus stigmate bifido acuto.

Drupa dipyrena, compressa: pyrenis subcoriaceis monospermis.

Humb. nov. gen. 3. t. 281.

43. **ERNODEA** Sw. (1788.)

Cal. quadripartitus, parvus.

Cor. hypocrateriformis, tubo elongato tetragono, fauce nuda, limbi lacinis revolutis.

Stam. tubo inserta, exserta. *Antherae* versatiles.

Stylus elongatus, filiformis, stigmate obtuso, emarginato.

Bacca bilocularis, loculis monospermis.

Sw. fl. ind. occid. 1. t. 4. [Pavetta foetidissima Cyrill. nesp. 1. t. 1. Asperula calabrica Herit. stirp. 4. t. 32. Putoria Pers.]

442. *CUNCEA* Hamilt. Don. (1825.)*Cal.* minimus, quadridentatus.*Cor.* tubo brevi, fauce villis clausa, limbo quadrilobo.*Antherae* lineares, sessiles, inclusae.*Stigma* bifidum.*Bacca* bilocularis, disperma.443. *BACONIA* Cand. (1807.)*Cal.* quadrilobus, liber.*Cor.* infundibuliformis, fauce villosa, limbo quadrilobo.*Stamina* fauci inserta, antheris linearibus.*Bacca* exsucca, bilocularis, disperma.*Cand. in ann. mus. 9. p. 219.*444. *FRÖLICHIA* Vahl (1796.)*Cal.* quadridentatus.*Cor.* tubulosa, limbi laciniis apice incrassatis triquetris.*Stigmata* duo, obtusiuscula.*Bacca* exsucca monosperma, seminis integumento laxo.*Vahl. ecl. t. 10.*445. *PAVETTA* L. (1747.)*Cal.* minimus, campanulatus, obsolete quadridentatus.*Cor.* infundibuliformis, tubo elongato gracili; limbo quadripartito, laciniis lanceolatis.*Stamina:* *Antherae* subsagittato-lineares, subsessiles extra faucem.*Stylus* filiformis, elongatus, stigmate oblongo, crassiusculus.*Bacca* bilocularis, altero loculo abortiente, ut monosperma fiat.*Bot. reg. 198. Gärtn. t. 25.*446. *TETRAMERIUM* Gärtn. fil. (1805.)*Cal.* parvus, obtuse quadridentatus.*Cor.* infundibuliformis, limbo quadripartito, aequali.*Stam.* filamentis brevissimis tubo inserta, antheris brevibus.*Stylus* simplex, stigmate bifido.*Bacca* drupacea, depressa, putaminis vertice quadripartito. Embryo in foveola albuminis sub chalaza.*Gärtn. t. 196. Humb. nov. gen. 3. t. 287. [Pavetta P. Brodiaeae t. 6. f. 1. Faramea Aubl. guian. t. 40.]*447. *IXORA* L. (1747.)*Cal.* exiguus, quadrifidus.*Cor.* tubo gracili elongato, limbo regulari, quadrilobo patentissimo.*Stam.* *Antherae* subsessiles ad faucem limbi, eiusque laciniis alternae.*Stylus* longe exsertus, stigmatibus binis erectis crassiusculis.*Bacca* bilocularis, disperma. Embryo dorsalis.*Gärtn. t. 25. 95. Bot. reg. t. 100. 154. 513. 540. 648. 782.*

40. HYDROPHYTUM Jack. (1825.)

Cal. integer.*Cor.* tubulosa, fauce villosa, limbo quadrifido.*Sam.* Antherae subsessiles, fauci corollae eiusque laciniis insertae.*Stylus* exsertus, stigmatibus bifido.*Locca* disperma. Semina integumento tenaci. Embryo nullus.

41. SIDERODENDRON Schreb. (1789.)

Cal. quadridentatus, minimus.*Cor.* tubo cylindrico, elongato, incurvo, limbo quadrilobo, subreflexo.*Stam.* filamentis brevissimis fauci insertis, antheris oblongis erectis.*Stylus* filiformis, stigmatibus obtuso.*Locca* bilocularis, dissepimento contrario. Sem. solitaria, dissepimento affixa.*Gärtn.* t. 197. [*Sideroxyloides Jacqu. amer.* t. 175. f. 9.]

42. HYDROPHYLAX L. fil. (1781.)

Cal. quadripartitus, laciniis subcarnosis.*Cor.* infundibuliformis, fauce barbata, laciniis limbi revolutis.*Stam.* filamentis tubo impositis, exserta. Antherae subhastatae.*Stylus* filiformis, stigmatibus bifido.*Drupa* exsucca, sulcato-angulata, bilocularis, dissepimento contrario. Sem. solitaria triquetra.*Roxb. corom.* 3. t. 233. [*Sarissus Gärtn.* t. 25.]

43. CHOMELIA Jacqu. (1763.)

Cal. tubulosus, limbo quadrifido, recurvo.*Cor.* tubo cylindrico longissimo; limbo quadrifido, patentissimo, laciniis acuminato-cuspidatis.*Stam.* filamentis brevissimis, tubo insertis; antherae lineares subprominulae.*Stylus* filiformis, stigmatibus duobus patentibus oblongis.*Drupa* ovalis, nuce quadrisulca, biloculari, loculis monospermis.*Jacqu. amer.* t. 13.

44. SCOLOSANTHUS Vahl. (1796.)

Cal. quadrifidus.*Cor.* tubulosa, subtetragona, limbo revoluta.*Stam.* filamentis basi subcoactis.*Drupa* monosperma.*Vahl. eclog.* 1. t. 10.

45. CUNNINGHAMIA Schreb. (1789.)

Cal. quadridentatus, exiguus.*Cor.* urceolata, hypocrateriformis, limbo quadrifido.

Discus perigynus glandulosus.

Stamina filamentis fauci insertis; antheris subrotundis.

Stylus filiformis, stigmatibus duobus obtusis.

Drupa nuce biloculari, disperma.

[*Malanea* *Aubl.* *guian.* t. 41. *Lam.* *ill.* t. 66. *Antirhea* J.

454. *COUSSAREA* *Aubl.* (1775.)

Cal. quadridentatus.

Cor. tubo brevissimo, limbi laciniis patentissimis, ac

Stam. filamentis cum corollae laciniis alternantibus, insertis. Antherae oblongae.

Stylus filiformis, stigmate quadrifido.

Bacca unilocularis monosperma.

Aubl. *guian.* t. 38.

455. *PETESIA* *P. Brown.* (1756.)

Cal. quadridentatus, deciduus.

Cor. infundibuliformis, limbo quadripartito.

Stam. filamentis brevibus tubo insertis, subinclusa.

Stylus simplex, stigmate bifido.

Drupa dipyrena, pyrenis monospermis. Placenta ciliolata. Sem. meniscoideo-umbilicata.

P. Brown. *jam.* t. 2. *Gärtn.* t. 192.

456. *RUBIA* *Plin.*

Cal. quadridentatus.

Cor. campanulato-rotata, quadri-(tri-quinque-)fida.

Styli saepe duo, plerumque simplex bifidus, stigmatibus globosis.

Bacca monosperma didyma. Embryo curvatus dorsum in albumine cartilagineo.

Sturm. *fl. germ.* *Gärtn.* t. 195.

457. *ANTHOSPERMUM* *L.* (1737.)

Flores dioeci.

Cal. minimus quadridentatus.

Cor. tubo brevissimo, limbo quadripartito.

Stam. filamentis tubo insertis, antheris oblongis exsertis.

Stylus brevis, stigmatibus binis.

Caryopses geminae ovals.

Linn. *hort. Cliff.* t. 27. *Gärtn.* t. 195.

458. *GALIMUM* *Diosc.*

Cal. quadridentatus.

Cor. subrotata, quadripartita; (interdum tripartita, triandra).

Stam. filamentis tubo insertis, antheris bilocularibus.

Stylus apice bipartitus, stigmatibus globosis.

Caryopsis geminae (glabrae, asperae aut hispidae, coronatae.)

Schk. *Handb.* t. 23. *Lam.* *ill.* t. 60.

459. *ASPERULA* *Dodon.* (1559.)

Cal. quadridentatus.

Cor. campanulata, vel infundibuliformis, quadrifida.

- Stam.* tubo inserta.
Stylus filiformis, stigmate bifido.
Caryopsis geminae (non coronatae.)
Schk. Handb. t. 23. *Reichenb.* ic. t. 93.
141. **VALANTIA** Tournef. (1706.)
Cal. quadridentatus persistens.
Cr. rotata: masculae binae germiui insidentes.
Achenium calyce denticulato coronatum.
Lam. ill. t. 361. *Michel.* nov. gen. t. 7.
142. **SHERARDIA** Dillen. (1719.)
Cal. quadrifidus.
Cor. infundibuliformis, quadrifido.
Stam. tubo corollae inserta.
Stylus filiformis, stigmatibus binis globulosis.
Caryopsis geminae, calycis denticulis coronatae.
Schk. t. 22. *Gärtn.* t. 24.
143. **CRUCIANELLA** L. (1737.)
Involutum uniflorum (abortientibus corollis lateralibus),
palaceum, di-triphyllum.
Cal. nullus.
Cor. tubo filiformi, limbo quadrifido.
Stam. tubo adnata.
Stylus inaequaliter bifidus, stigmatibus globulosis.
Caryopsis duae.
Schmidel. ic. t. 23. 24. *Lam.* ill. t. 61. *Reichenb.* ic. 3. t. 262.
144. **SPERMACOE** Dillen. (1732.)
Cal. quadridentatus persistens.
Cor. infundibuliformis, quadrifida.
Caps. bilocularis, calyce coronata, non dehiscens, loculo
 altero clauso. Semen solitarium.
Lam. ill. t. 62. *Gärtn.* t. 25.
145. **BORRERA** Meyer. (1818.)
Involutum polyphyllum, foliis setaceis.
Cal. bi-quadrifidus, laciniis subulatis, interiectis saepe
 totidem setis.
Cor. infundibuliformis, limbo quadrifido.
Caps. bilocularis, bipartibilis; dissepimento imperfecto,
 e marginibus valvarum inflexis orto. Sem. solitaria.
Meyer fl. essequib. t. 1. [Octodon *Schumach.* guin. p. 74.]
146. **DIODIA** Gronov. (1743.)
Cal. persistens, disepalus, vel tetrasepalus, sepalis duo-
 bus longioribus.
Cor. infundibuliformis, limbo quadripartito.
Stam. tubo inserta.
Stylus filiformis, *Stigma* bifidum.
Caps. ovata tetragona, bilocularis, bivalvis. Sem. hinc
 convexa, inde sulcata.
Lam. ill. t. 63. *Gärtn.* t. 25. [Stavrospermum *Thonning* apud
Schumach. guin. p. 73.]

466. *KNOXIA* L. (1747.)*Cal.* tetrasepalus, inaequalis, deciduus.*Cor.* infundibuliformis, tubo filiformi, limbo quadri-*Stamina* fauci corollae inserta.*Stylus* filiformis, stigmatibus duobus capitatis.*Caps.* bipartibilis, disperma. Placenta media fi
(sicut in Umbelliferis.)*N. L. Burm. fl. ind. t. 13. f. 2. Gärtn. t. 25.*b. *Fructibus tetraspermis.*467. *LASIANTHUS* Jack. (1825.)

Flores axillares, bracteis magnis oppositis.

Cal. quadri-(interdum quinque-)partitus, lacinis
ribus.*Cor.* infundibuliformis, utrinque pilosa, limbo qui
(interdum quinquefido.)*Bacca* pilosa, calyce coronata, tetrasperma. F
cylindricus, in axi albuminis.468. *MYRMECODIA* Jack. (1825.)

Flores basibus petiolorum semitecti.

Cal. subinteger.*Cor.* tubulosa, quadrifida: annulo barbato tubi inf
minum insertionem.*Stam.* corolla breviora.*Stylus* stigmatibus simplicibus.*Bacca* ovata, quadrilocularis, tetrasperma.469. *DUNALIA* * (1815.)*Cal.* quadridentatus, dentibus bifidis.*Cor.* regularis, quadrifida, calycem aequans.*Stam.* corolla breviora.*Stylus* bifidus, stigmatibus subulatis.*Caps.* globulosa, quadrivalvis, bilocularis, loculis
mis. Sem. globosa.[Peplis tetrandra Jacqu. amer. t. 180. f. 29, Hedyot
rosa Sw. obs. bot. t. 1. f. 2.]470. *MYONIMA* Commers. Juss. (1789.)*Cal.* minimus, subinteger.*Cor.* tubo brevi, limbo quadrilobo.*Stam.* exserta.*Stigma* crassiusculum.*Bacca* sicca, depresso, non coronata, nuce quat
lari, tetrasperma. Sem. hinc convexa, inde con*Lam. ill. t. 68.*c. *Fructibus polyspermis.*471. *PYROSTRIA* Commers. Juss. (1789.)*Cal.* minimus, subquadridentatus.

Cor. subcampanulata, semiquinquefida, fauce tomentosa.
Stigma capitatum.

Fructus pyriformis, octostriatus: nucibus octo monospermis.

Lam. ill. t. 68.

EVESMIA Bonpl. (1814.)

Cal. parvus quadridentatus.

Cor. subrotata, quadrifida.

Stam. fauci inserta, exserta.

Stigma incrassatum.

Fructus ovatus, calyce coronatus, quadrilocularis, loculis polyspermis.

Humb. plant. aequinoct. 2. t. 131. [O-higginsia Ruiz. et Pav. fl. peruv. 1. t. 83. 85.]

GONZALEA Pers. (1803.)

Cal. uccolatus quadrifidus, persistens.

Cor. infundibuliformis, tubo elongato, limbo quadripartito.

Stigma inclusa.

Stylus filiformis, stigmatibus quadrifido.

Drupa depresso-globosa, tetrapyrena, pyrenis polyspermis.

Humb. plant. aequinoct. 1. t. 64. [Gonzalagunia Ruiz. et Pav. flor. peruv. 1. t. 86. Buena Cur. in anal. de cienc. 2. t. 23.]

COCCOCYPSELUM P. Brown. (1756.)

Cal. quadripartitus.

Cor. infundibuliformis, limbo quadripartito erecto.

Stam. tubo adnata, inclusa.

Stylus apice bifidus, stigmatibus oblongis.

Bacca subrotunda, bilocularis: loculis polyspermis.

P. Brown. jam. t. 6. f. 4. Swartz. fl. ind. occid. 1. t. 5. [Condalia repens Ruiz. et Pav. fl. peruv. 1. t. 84. Tontanea guianensis Aubl. guian. 1. t. 42. Bellardia W. Fernelia Lam. ill. t. 67. Gärtn. t. 191.]

HOFFMANNIA Sw. (1788.)

Cal. parvus, quadridentatus.

Cor. hypocrateriformis, tubo brevissimo, limbo quadripartito.

Antherae sessiles in fauce, lanceolato-lineares, acutae, conniventes.

Stylus filiformis, stigmatibus subemarginato.

Bacca subtetragona, bilocularis: placenta distincta ovata in utroque loculo.

Swartz. flor. ind. occid. 1. t. 5.

HELOSPORA Jack. (1825.)

Cal. quadridentatus, persistens.

Cor. tubulosa, fauce nuda, limbo quadripartito.

Antherae sessiles, lineares, inclusae.

Stylus quadrisulcus, stigmatibus quatuor.

Bacca calyce coronata, polysperma, seminibus dasyserie cruciatim dispositis, nidulantibus, linearibus.

477. *CATESBAEA* Gronov. (1743.)

Cal. quadridentatus seu quadrifidus, persistens.

Cor. tubo longissimo, infundibuliformis, limbo quadri-

Stam. filamentis basi corollae insertis, longissimis, exsertis.

Stylus filiformis, stigmate bidentato.

Bacca calyce coronata, bilocularis; dissepimento utrisque placentifero. Sem. imbricata, squamata.

Lam. ill. t. 67. *Gärtn.* t. 192. *Bot. reg.* t. 858.

478. *CARPHELEA* Juss. (1789.)

Cal. turbinatus, quadrifidus, lacinis spathulatis seorsum persistentibus.

Cor. tubo filiformi elongato; fauce ventricosa, interiori limbo quadripartito, lacinis angustis.

Antherae subsessiles oblongae, fauci insertae.

Caps. calyce coronata, bilocularis, bivalvis: dissepimento valvis opposito, bipartibili.

Lam. ill. t. 59.

479. *MANETTIA* Mutis. L. (1771.)

Cal. quadri-vel octo- (interdum quinque-vel decem-) partitus.

Cor. tubulosa, fauce coarctata, limbo quadripartito hirsuto.

Stamina fauci inserta, inclusa.

Stylus simplex, stigmate bifido.

Caps. bivalvis, valvis introflexis dissepimentum interius: placentae duae liberae. Sem. orbiculata, membranacea.

Bot. reg. 693. *Ruiz. et Pav.* flor. peruv. 1. t. 89. 90. *G.* t. 197. [*Lygistum* P. Brown. *jam.* t. 3. f. 2. *Nam.* Aubl. t. 37.]

480. *BOUVARDIA* Salisb. (1805.)

Cal. quadripartitus, dentibus interiectis.

Cor. tubulosa, limbo quadrifido.

Stamina filamentis tubo adnatis; antherae tubum superantes.

Stylus simplex, stigmate bilobo.

Caps. bilocularis. Sem. marginata.

Salisb. parad. lond. t. 88. *Bot. reg.* t. 107. *Humb.* *gen.* 3. t. 288. [*Houstonia* Andr. *repos.* t. 106. *Las* Jacqu. *hort. schönbrunn.* 3. t. 257. *Aeginetia* Cav. *ic.* t. 305.]

481. *HEDYOTIS* L. (1747.)

Cal. quadripartitus.

Cor. tubulosa, quadrifida.

Stam. subexserta.

Stylus simplex, stigmate bifido.

Caps. didyma, bilocularis: apice inter dentes calycis dehiscentis, placentis columnaribus in utroque loculo.

Cav. ic. 6. t. 573. 574. Ruiz. et Pav. fl. peruv. 1. t. 87. 88.

Humb. nov. gen. 3. t. 289. Roxb. coroin. 1. t. 3. Bot. mag. 2840. Gärtn. t. 30.

II. COROLLA PLEIOPETALA.

A. *Fructus superi.*

482. EPIMEDIUM (Diosc.) Dodon. (1559.)

Cal. tetrasepalus, coloratus, basi bracteolis suffultus.

Petala quatuor ovalia.

Nectaria petaliformia, cucullata.

Antherae biloculares, a basi inde dehiscentes.

Folliculus bivalvis.

Berberideae. Schk. Handb. t. 24. Sturm. fl. germ.

483. PTELEA L. (1751.)

Flores dioeci.

Cal. brevis quadripartitus.

Petala quatuor longiora patentissima.

Stam. Filamenta basi hirsuta. Antherae erectae biloculares.

Stylus brevis. Stigma bilobum.

Samara bilocularis. Sem. oblonga, embryo inverso.

Acera. Schmidt Baumz. 2. t. 76. Gärtn. t. 49.

484. SERINGIA * (1818.)

Cal. quadrilobus.

Petala quatuor ovata, patentia.

Discus perigynus staminifer.

Stam. filamentis brevibus, antheris bilobis.

Stylus subnullus, stigma minimum.

Samara bilocularis, loculo altero abortiente. Embryo erectus.

Acera. [Ptelidium Aubert du Pet. Thouars ins. austr. Afric. t. 2.]

485. PHILAGONIA Blum. (1825.)

Flores dioeci.

Cal. quadridentatus.

Petala quatuor.

Stam. petalis alterna. Antherae biloculares dorso affixae.

Discus perigynus annularis obsoletus.

Stigma peltatum magnum.

Caps. quadrilocularis, loculis dispermis.

Terebinthaceae.

486. MAYEPEA Aubl. (1775.)

Cal. quadripartitus.

Petala quatuor, basi concava, apice in filamentum longum abeuntia.

Stam. Filamenta brevissima. Antherae oblongae biloculares, in cavitate petalorum reconditae.

Stylus nullus. Stigma crassiusculum.

Drupa olivaeformis, nuce lignosa monosperma.

Rhamneae. Aubl. guian. t. 31. [Ceranthus Schreb.]

487. *SAMARA* L. (1771.)

Cal. quadripartitus exiguus.

Petala quatuor ovata, basi lacuna longitudinali.

Stam. Filamenta subulata elongata, lacunis petalorum inserta. Antherae subcordatae.

Stylus cylindricus. Stigma infundibuliforme.

Drupa monosperma.

Rhamneae. [Cornus zeylanica. Burm. zeyl. t. 31. ? Elaeagnus Forst. char. gen. t. 6.]

488. *EVODIA* Forst. (1775.)

Cal. quadripartitus, persistens.

Petala quatuor, calycem subaequantia.

Discus perigynus urceolatus quadrilobus.

Stam. extra discum, filamentis erectis, antheris subcordatis

Stylus quadrisulcus. Stigma quadrilobum.

Caps. quadricocca, coccis bivalvibus, monosperma.

Rutaceae. Forst. char. gen. t. 7. Adr. Juss. in Mém. du Mus. t. 22. f. 28. Labill. nov. Caled. t. 74.

489. *ZIERIA* Sm. (1798.)

Cal. quadripartitus.

Petala quatuor.

Discus perigynus petala promit et glandulas quatuor staminiferas.

Stam. subulata. Antherae cordatae.

Stylus quadrisulcus, stigmate quadrilobo.

Cocci quatuor bivalves.

Rutaceae. Adr. Juss. in Mém. du Mus. t. 22. f. 2 Bot. mag. t. 1395.

490. *MONETIA* Herit. (1784.)

Cal. campanulatus quadrifidus.

Petala quatuor linearia subreflexa.

Stam. receptaculo inserta, antheris oblongis bilocularibus

Stylus simplex, stigmate acuto.

Bacca monosperma.

— — Herit. stirp. t. 1.

491. *SKIMMIA* Thunb. (1784.)

Cal. quadripartitus, minimus, persistens.

Petala quatuor minuta, concava.

Stam. filamentis brevissimis.

Bacca obsolete quadrisulca, tetrasperma.

— — Thunb. jap. p. 4. 62.

† 492. *LEPTA* Lour. (1790.)

Cal. quadripartitus.

Petala quatuor ovata.

Stam. Filamenta brevissima. basi petalorum inserta. Antherae didymae, quadrisulcae.

Bacca quadriloba, monosperma.

[Oihera Thunb.]

† 493. *ORIXA* Thunb. (1784.)

Cal. quadripartitus, brevissimus.

Petala quatuor lanceolata.

Stam. Filamenta petalis breviora. Antherae globosae.

Stylus simplex, stigmate capitato.

94. *CURTISIA* Ait. (1789.)

Cal. quadripartitus.

Petala quatuor ovata, obtusa.

Stam. Filamenta receptaculo inserta. Antherae ovatae.

Stylus simplex, stigmate quadrifido.

Drupa subglobosa, nuce quadriloculari, loculis monospermis.

Celastrinae Cand. [Sideroxylon Burm. afr. t. 82.]

495. *BOTRYCERAS* Willd. (1811.)

Involucrum multifidum, laciniis bifidis curvatis: flores de-
mm secundi reflexi.

Cal. quadripartitus persistens.

Petala quatuor linearia persistentia.

Stam. Filamenta brevissima. Antherae oblongae, erectae.

Stylus arcuatus, stigmate capitato.

Nux?

— — Berl. Mag. 5. t. 10. f. 10.

496. *CISSUS* L. (1747.)

Cal. minimus, integer s. obsolete dentatus.

Petala quatuor persistentia.

Discus perigynus staminifer.

Stamina corollam aequantia. Antherae subrotundae.

Stylus simplex, stigmate acuto.

Bacca monosperma.

Ampelideae Lam. ill. t. 81

497. *ALLASIA* Lour. (1790.)

Cal. quinquefidus inferus.

Petala quatuor supera, subrotunda, concava, pilosissima.

Stam. Filamenta crassa acuta. Antherae bilobae, lobis bilocularibus.

Stylus subulatus.

Bacca unilocularis, polysperma.

498. *SIPHANTHERA* Pohl. (1827.)

Flores aggregati. *Involucrum* polyphyllum.

Cal. campanulatus, quadridentatus, interjectis foliolis longioribus persistentibus.

Pet. quatuor orbiculata, dentibus calycis adnata.
Stam. cum petalis alternantia, calycis margini inser
 Filamenta filiformia. Antherae basi processu obcorda
 apice rostrata, poro dehiscentia.

Styl. clavatus.

Caps. bilocularis polysperma.

Melastomeae. Pohl. plant. bras. t. 84. 85.

B. *Fructus inferi.*

499. CORNUS Virgil.

Cal. quadridentatus.

Petala quatuor.

Stamina receptaculo inserta, cum petalis alternantia.

Stylus simplex, stigmate obtuso.

Drupa disperma. Embryo inversus in axi albuminis.

Caprifolia. Schk. t. 24. Gärtn. t. 26.

500. AUCUBA Thunb. (1784.)

Flores monoeci.

Cal. truncatus, persistens, obsolete quadridentatus.

Petala quatuor ovata, patentia.

Discus perigynus (et in masculis floribus circa rudime
 tum styli) planus, media fovea insignis.

Stam. receptaculo inserta, cum petalis alternis. Fi
 menta crassa, brevissima. Antherae ovatae, didymae
 quadrisulcae.

Stylus brevis, persistens. Stigma capitatum.

Bacca monosperma.

— — Bot. Mag. 1197.

501. GLOSSOMA Schreb. (1789.)

Cal. turbinatus quadridentatus, persistens.

Petala quatuor.

Stam. Antherae subsessiles oblongae, subcoalitae, apice
 ligulam membranaceam productae, intus dehiscentes.

Stylus antherarum cono tectus. Stigmata quatuor.

Drupa monosperma.

— — [Votomita Aubl. guian. t. 35.

502. MONTINIA L. fil. (1781.)

Flores dioeci.

Cal. quadridentatus.

Petala quatuor, obtusissima, patentia.

Stam. cum petalis alterna: sterilia in ♀.

Stylus bifidus, stigmatibus peltatis.

Caps. ovalis, bilocularis, bivalvis. Placenta centralis. Se
 compressa, foliacea, imbricata.

Onagrae. Smith. spicil. t. 15. Gärtn. t. 33.

503. TRAPA L. (1737.)

Cal. quadripartitus.

Petala quatuor.

Stylus conicus, stigmatē capitato.

Nux bi-quadricornuta, tota vitello (*Gärtn.*) seu placenta radicali (*Tittmann.*) repleta. Albumen nullum. Radicula sursum progerminans.

Hydrocaryas Link. Schk. t. 25. *Gärtn.* t. 26. *Roxb. corom.* 3. t. 234. *Tittmann* Keimung der Pflanzen. t. 5.

94. *ANMANNIA* Houst. L. (1737.)

Cal. campanulatus, octodentatus, dentibus alternis minoribus.

Petala quatuor, summo calycis tubo inserta, aut nulla.

Stam. petalis alterna, rarius octo aut quinque.

Caps. quadrilocularis, polysperma. Placenta ad angulos loculorum centrales.

Salicariae. Willd. hort. berol. t. 6. 7. *Gärtn.* t. 112.

95. *LUDWIGIA* L. (1737.)

Cal. quadripartitus, persistens.

Petala quatuor, cum calycis lacinii alternantia, aut nulla.

Stamina lobis calycis opposita.

Stylus simplex, stigmatē capitato.

Caps. quadrilocularis, quadrivalvis, polysperma.

Lam. ill. t. 77. *Gärtn.* t. 31. [*Isuardia* L. Schk. t. 25. *Dauria Thuars.*]

†† *Flores imperfecti.*

I. FRUCTUS SUPERIOR.

PROTEACEAE.

A. *Fructus clausi.*

a. *Antherae distinctae.*

α. *Flores dioeci bracteati.*

96. *AULAX* Berg. (1767.)

Flores ♂ racemosi, ♀ subcapitati.

Cal. corollinus tetrasepalus medio staminifer.

Stigma clavatum obliquum.

Nux ventricosa, barbata.

Burm. afr. t. 70. f. 3. [*Protea Andr.* reposit. 76. 248.]

97. *LEUCADENDRON* Boerh. (1720.)

Flores ♂ et ♀ capitati.

Nux squamis strobili dilatatis inclusa.

Bot. reg. 402. 826. 979.

β. Flores hermaphroditi.

αα. *Antherae apicibus concavis calycis immersae*

αα. *Ovarium monospermum*

508. *PETROPHILA* R. Br. (1810.)

Calyx corollinus totus deciduus.

Squamae hypogynae nullae.

Stylus basi persistens. Stigma fusiforme apice attenu

Strobilus ovatus. Nux lenticularis, hinc comosa.

[*Protea Cav. ic. 6. t. 550. 551. Schrad. sert. hannov. 2*

509. *IsoPOGON* R. Br. (1810.)

Calyx corollinus persistens.

Squamae hypogynae nullae.

Stylus totus deciduus. Stigma fusiforme.

Nux sessilis ventricosa, undique comosa.

Bot. reg. 900. [*Protea Cav. anal. de cienc. nat. 1.*
Botan. Mag. 697.]

510. *PROTEA* L. (1737.)

Involucrum imbricatum, persistens. Receptaculum mune paleaceum.

Cal. corollinus bilabiatus, labii latioris laminis stans cohaerentibus.

Squamae quatuor hypogynae.

Stylus subulatus persistens.

Nux undique barbata, apice caudata.

Bot. reg. 20. 47. 206. 354. 569. 1023. *Gärtn. t. 51.*
dendron et *Pleuranthe Salisb.*]

511. *LEUCOSPERMUM* R. Br. (1810.)

Involucrum et receptaculum similia.

Cal. tubulosus, subbilabiatus, laminis staminiferis dist

Stylus filiformis, deciduus, stigmathe incrassato glal

Nux sessilis, ventricosa, laevis.

[*Protea Andr. repos. 17. 294. Leucadendron Berg. I*
atella Salisb.]

512. *MIMETES* R. Br. (1810.)

Involucrum et receptaculum Proteae, sed hoc pl paleis deciduis.

Cal. aequalis quadripartitus, laciniis distinctis.

Stylus filiformis deciduus, stigmathe cylindrico, gra

Nux ventricosa, sessilis, laevis.

Reichenb. hort. 1. t. 92. 93. [Lepidocarpodendron Boerl.
lugd. bat. 2. p. 194. Hypophyllocarpodendron ib.]

513. *SERRURIA* R. Br. (1810.)

Receptaculum convexum, paleis persistentibus.

Cal. quadrifidus, subaequalis, unguibus distinctis, b suffultus.

Squamae quatuor hypogynae.

- Stigma verticale glabrum.*
Nux brevissime pedicellata, ventricosa.
 [Serraria *Burm. afr.* t. 99.]
54. *NIVENIA* R. Br. (1810.)
Invulcrum tetraphyllum, quadriflorum, fructiferum nudum. Receptaculum planum nudum.
Cal. quadrifidus aequalis deciduus.
Squamae hypogynae liberae.
Stigma clavatum verticale.
Nux ventricosa, sessilis, basi integra.
 [Protea *Andr. repos.* 243. *Peranomus Salisb.*]
55. *SOROCEPHALUS* R. Br. (1810.)
Invulcrum tri-hexaphyllum, pauciflorum, fructiferum non mutatum. Receptaculum nudum.
Cal. quadrifidus aequalis, deciduus.
Stigma verticale clavatum.
Nux ventricosa, brevissime pedicellata, vel basi emarginata.
 [Protea *Andr. repos.* 517. *Sorauthe Salisb.*]
56. *SPATALLA* R. Br. (1810.)
Invulcrum di-tetraphyllum, uni-quadriflorum. Receptaculum nudum.
Cal. quadrifidus, lacinia interiore plerumque maiore.
Stigma obliquum, dilatatum.
Nux ventricosa, brevissime pedicellata.
 [Protea *Thunb. diss. de Protea,* t. 3. 4.]
57. *ADENANTHOS* Labill. (1804.)
Invulcrum uniflorum, subimbricatum, oligophyllum, calyciforme.
Cal. quadrifidus, infra circumscissus.
Squamae hypogynae, basi calycis persistentis adnatae.
Stigma verticale.
Nux ventricosa.
Labill. nov. holland. t. 36—38.
- ββ. *Ovarium dispernum.*
58. *GUEVINA* Molin. (1782.)
Cal. tetrasepalus, irregularis; sepalis tribus revolutis, quarto erecto.
Glandulae duae hypogynae anticae.
Stigma obliquum.
Drupa, putamine osseo, monosperma.
 [Nebu *Feuill.* 3. t. 34. (non 33.) *Quadria Ruiz. et Pav. fl. peruv.* t. 33. *Gärtn.* t. 220.]
- b. *Antherae exsertae, medio vel basi calycis insertae.*
 αα. *Glandulae hypogynae. Stamina sterilia quatuor.*
19. *BRABBIUM* L. (1737.)
Flores polygami.

Cal. tetrasepalus, regularis.
Stam. basi calycis inserta.
Vaginula perigyna glandulosa.
Stigma verticale.
Drupa exsucca monosperma, tomentosa.

Gärtn. t. 217. [*Amygdalus Breyn.* cent. t. 1.]

520. *PERSOONIA* Sm. (1798.)

Flores hermaphroditi.

Cal. tetrasepalus, regularis, sepalis apice recurvis, cidiis.

Stam. mediis sepalis inserta, erecta.

Glandulae hypogynae carnosae.

Stigma obtusum.

Drupa baccata, nuce 1-2 loculari, stylo persistente, ronata.

Andr. repos. 77. *Labill.* nov. holl. t. 45. *Rudge* in *transact.* 10. t. 16. *Gärtn.* t. 220. [*Liukia Cav.* ic. t. 3]

521. *CENARRHENES* Labill. (1804.)

Flores hermaphroditi.

Cal. tetrasepalus, regularis, patens.

Stam. basi calycis inserta, sepalis multo minora, ^{quasi} fertilia, totidemque cassia minora cum iis alternantia.

Drupa baccata.

Labill. nov. holl. t. 50.

ββ. *Glandulae* nullae hypogynae.

522. *AGASTACHYS* R. Br. (1810.)

Cal. regularis, quadripartitus, medio staminifer.

Filamenta distincta.

Stigma unilaterale.

Germen sessile, trigonum, monospermum.

Bot. reg. 1015.

523. *SYMPHYONEMA* R. Br. (1810.)

Cal. tetrasepalus, regularis.

Stam. basi sepalorum inserta: filamenta apice coalita.

Stigma truncatum.

Germen dispermum.

Reichenb. hort. 2. t. 107.

cc. *Antherae* tubo calycis adnatae.

524. *FRANKLANDIA* R. Br. (1810.)

Cal. hypocrateriformis, limbo quadripartito regulari.

Antherae inclusae.

Squamae hypogynae in vaginam connatae.

Nux fusiformis pedicellata, apice dilatato papposo.

R. Brown ad *Flinders.* it. t. 6.

dd. *Stamina hypogyna.*

88. *BILLENDENA* R. Br. (1810.)

Cal. tetrasepalus, regularis, patens.

Glandulae nullae hypogynae.

Sigma simplex.

Nux compressa, submonosperma.

Antherae cohaerentes, vicinarum lobis proximis loculum constituentibus, tandem distinctis.

89. *SINSIA* R. Br. (1810.)

Cal. tetrasepalus, regularis, laminis reflexis.

Stam. exserta.

Stammas hypogynae nullae.

Sigma dilatatum, concavum.

Nux obconica.

90. *COSPERMUM* Sm. (1798.)

Cal. tubulosus, ringens; lacinia suprema basi fornicata.

Antherae tres inclusae, laterales dimidiatae: superior

loba: primo cohaerentes, lobis proximis vicinarum loculum constituentibus.

Sigma liberum.

Nux obconica, pilis rigidis coronata.

Smith exot. bot. 2. t. 82. *Rudg.* in Linn. transact. 10. t. 17.

Hook. in Bot. Mag. 2850. *Gärtn.* t. 215.

91. *SYNAPHEA* R. Br. (1810.)

Cal. tubulosus, ringens; lacinia suprema latiore.

Antherae tres inclusae, laterales dimidiatae: inferior biloba: primo cohaerentes, lobis proximis vicinarum loculum constituentibus.

Sigma filamento superiori sterili connatum.

Nux obovata pilis rigidis coronata.

R. Brown ad Flinders. it. t. 7.

B. *Fructus dehiscentes.*

a. *Uniloculares.*

α. *Ovarium dispermum.*

αα. *Antherae* apicibus calycis concavis immersae.

92. *ANADENIA* R. Br. (1810.)

Cal. tetrasepalus.

Glandulae hypogynae nullae.

Sigma conicum.

Folliculus monospermus. Semen apterum.

93. *GREVILLEA* R. Br. (1810.)

Cal. irregularis, tetrasepalus, suboblique insertus, sepalis secundis.

Glandula hypogyna solitaria, dimidiata.

Stigma obliquum, depressum.

Folliculus unilocularis, loculo centrali, dispermus.

mina apice brevissime vel undique alata.

Bauer ill. nov. holl. t. 9. Bot. reg. t. 443. *Hooker*. ex
3. t. 216. [*Lysanthe* et *Stylurus Salisb.* *Embothria*
nov. holl. t. 9. *Cav.* ic. t. 386.]

531. *HAKKA* Schrad. (1795.)

Cal. irregularis, tetrasepalus, sepalis secundis.

Glandula hypogyna solitaria, dimidiata.

Stigma subobliquum, e basi dilatata mucronatum.

Folliculus unilocularis, loculo excentrico, dispermus.

mina apice longe alata.

Schrad. sert. hannov. t. 17. *Cav.* ic. t. 533—535.
nov. holl. t. 39—41. Bot. reg. 475. [*Conchocarpus*
Banksia Gärtn. t. 47. f. 2.]

bb. *Antherae* exsertae, laciniis calycis revolutis

532. *LAMBERTIA* Sm. (1798.)

Involucrum coloratum, uni-septemflorum. Receptaculum
planum, nudum.

Cal. tubulosus quadrifidus.

Stam. laciniis inserta.

Squamae hypogynae quatuor, distinctae vel connatae

Stigma subulatum.

Folliculus unilocularis. Sem. marginata.

Cav. anal. de cienc. 1. t. 15. ic. t. 547. *Andr.* repos. t.

533. *XYLOMELUM* Sm. (1798.)

Flores polygami spicati. *Involucrum* nullum.

Cal. tetrasepalus, regularis.

Stam. supra medium sepalorum inserta.

Glandulae hypogynae quatuor.

Stigma clavatum.

Folliculus unilocularis, loculo excentrico. Sem.
alata.

Reichenb. hort. 1. t. 90. [*Banksia pyriformis Gärtn.*
Hakea Cav. ic. t. 536.]

534. *ORITES* R. Br. (1810.)

Flores spicati. *Involucrum* nullum.

Cal. tetrasepalus, regularis.

Stam. supra medium sepalorum inserta.

Glandulae hypogynae quatuor.

Stylus strictus, stigmate obtuso verticali.

Folliculus unilocularis, loculo subcentrali. Sem.
alata.

535. *ROPALA* (Roupala) Aubl. (1775.)

Flores racemosi. *Involucrum* nullum.

Cal. tetrasepalus, regularis.

Stam. sepalis inserta.

Glandulae hypogynae quatuor, distinctae vel connatae.

Stigma clavatum.

Folliculus unilocularis. Sem. utrinque alata, nucleo centrali.

Abbl. guian. t. 32. Gärtn. t. 217. Rudg. guian. t. 39. Humb. nov. gen. 2. t. 118—121. Pohl plant. bras. t. 86—90. [Helicia Lou. Euplassa et Panopsis Salisb. Andriapetalum Pohl plant. bras. t. 91. 92. paullo magis soluta a sepalis habet stamina.]

β. Ovarium tetraspermum.

KNIGHTIA R. Br. (1810.)

Cal. tetrasepalus, regularis.

Stam. extra medium sepalis inserta.

Glandulae hypogynae quatuor.

Stigma verticale subclavatum.

Folliculus stylo persistente coronatus, unilocularis. Sem. apice alata.

Lin. transact. 10. t. 2.

γ. Ovarium polyspermum.

EMBOTHRIUM Forst. (1775.)

Cal. irregularis, hinc longitudinaliter fissus, inde quadridentatus.

Stam. apicibus concavis calycis immersa.

Glandula hypogyna solitaria semi- seu subannularia.

Stylus persistens, stigmatibus clavato.

Folliculus unilocularis. Semina apice alata.

Gärtn. t. 218. Cav. ic. t. 388. Labill. nov. holland. t. 44. [Telopea R. Br. bracteas habet plerumque, deciduas. Hylogyne Salisb.]

ORBOCALLIS R. Br. (1810.)

Cal. irregularis, hinc longitudinaliter fissus, inde quadridentatus.

Stam. apicibus concavis calycis immersa.

Glandula hypogyna nulla.

Stigma obliquum concaviusculum.

Folliculus cylindricus. Sem. apice alata.

[Embothrium Ruiz. et Pav. fl. per. 1. t. 95.]

LOMATIA R. Br. (1810.)

Cal. irregularis, sepalis distinctis secundis.

Stam. apicibus calycis concavis immersa.

Glandulae hypogynae tres secundae.

Germen longe stipitatum.

Stylus persistens, stigmatibus obliquo, subrotundo, dilatato.

Folliculus oblongus. Sem. apice alata, ala marginata.

Bot. reg. 442. [Embothrium Labill. nov. holl. t. 42. 43. Cav. ic. 4. t. 388. Tricoudylus Salisb.]

540. *STENOCARPUS* R. Br. (1810.)*Cal.* irregularis, sepalis distinctis secunda.*Stam.* apicibus calycis concavis immersa.*Glandula* hypogyna solitaria semiannularis.*Germen* stipitatum.*Stylus* deciduus, stigmate obliquo, subrotundo,*Folliculus* linearis. Sem. basi alata.Bot. reg. 441. [*Embothrium* Forst. char. gen. 1
Cybele Salisb.]b. *Fructus biloculares, dissepimento libero bij*541. *BANKSIA* L. fil. (1781.)Flores spicati. Bracteae ternae florum singulorum
sistentes.*Cal.* quadripartitus.*Stam.* apicibus concavis calycis immersa.*Squamae* hypogynae quatuor.*Folliculus* ligneus, dispermus. Sem. apice cuneatGärtn. t. 48. Cav. ic. 6. t. 537. 538. 540—546. 1
688. 697. 787.542. *DRYANDRA* R. Br. (1810.)Involucrum commune imbricatum. Receptaculum
ceum.*Cal.* quadripartitus.*Stam.* apicibus concavis calycis immersa.*Squamae* hypogynae quatuor.*Folliculus* ligneus dispermus.Linn. transact. 10. t. 3. [*Banksia* Labill. it. 1. t. 24
... phia Salisb.]543. *PHYLA* Lour. (1790.)Involucrum imbricatum polyphyllum. Receptaculum
leaceum, paleis binis singulos flores fulcientibus.*Cal.* corollinus quadrifidus, lacinia suprema emarg*Stam.* Filamenta brevia, bino ordine tubo affixae
therae bilobae subrotundae.*Stylus* brevis, stigmate crassiusculo,*Sem.* nuda receptaculo affixa.

T H Y M E L A E A E.

544. *CANSIERA* Juss. (1789.)*Cal.* globoso-urceolatus, quadridentatus.*Stamina* imo calyci inserta. Antherae subrotundae
exsertae.

Squamae perigynae quatuor.

Stylus pisiformis monosperma.

Roxb. corom. 2. t. 103.

DIAPETES Lam. (1792.)

Sam. aggregato-fasciculati. Receptaculum barbatum.

Cal. corollinus infundibuliformis, limbo quadrifido.

Antherae exsertae.

Gynopsis calyce tecta.

Journ. d'hist. nat. t. 10. f. 1.

PIRUTHIOLA L. (1767.)

Cal. corollinus bracteatus, tubo gracili, limbo quadrifido.

Glandulae squamaceae octo in fauce.

Gemmae in tubo recondita.

Sam. monosperma.

Smith exot. bot. 1. t. 46. Andr. repos. 43. 49. Gärtn.

t. 215.

CONOCEPHALUS Blum. (1828.)

Sam. saecii, capitati,

Cal. tubulosus, tri-quadrifidus.

Sam. Filamenta brevia, clavata. Antherae biloculares, basi discretis.

Stylus simplex, apice in stigma tubulosum abiens.

Sam. calyce baccato vestita, monosperma.

Urticeae Bot. reg. 1203. A. et B.

C H E N O P O D E A E.

TRIVINA Plum. (1703.)

Cal. quadripartitus, persistens.

Sam. quatuor, interdum octo.

Stylus brevis, stigmate obtuso.

Sam. monosperma.

Lam. ill. t. 81. Gärtn. t. 77.

CRYPTOCARPUS Kunth. (1817.)

Cal. campanulato-urceolatus, quadri-quinquedentatus.

Sam. exserta: filamenta basi subconnata: antherae didymae.

Stylus simplex.

Achenium calyce inclusum.

Humb. nov. gen. 2. t. 123.

PTERANTHUS Forsk. (1775.)

Cal. quadripartitus, persistens: laciniis concavis, duabus maioribus cristatis.

Filamenta basi connata.

Utriculus monospermus, calyce tectus.

Gärtn. t. 213. Lam. ill. t. 764. [Louichea Herit. stirp. 1. t. 66.]

551. CAMPHOSMA L. (1747.)

Cal. urceolatus, quadrifidus; laciniis oppositis ~~ma~~
recurvatis.

Stylus bi-trifidus. Stigmata acuta.

Garyopsis nuda, calyce tecta.

Kitaib. pl. hung. 1. t. 63. *Gärtn.* t. 213.

552. ANISACANTHA R. Br. (1810.)

Cal. campanulatus, quadrifidus, post anthesin ~~auc~~

Stam. tria, quatuor.

Stylus bipartitus.

Utriculus monospermus, calyce nucamentaceo,
spinis quaternis inaequalibus instructo, inclusus.

553. NYSSANTHES R. Br. (1810.)

Cal. quadripartitus, irregularis, laciniis duabus
ribus inaequalibus: bracteis spinescentibus.

Stamina bina, quaterna, basi connata, squamulis
positis. Antherae biloculares.

Stylus simplex, stigmate capitato.

Utriculus monospermus.

554. EMPLEURUM Soland. Ait. (1789.)

Flores subinde polygami.

Cal. quadrifidus, glandulosus.

Stam. hypogyna, exserta. Antherae apice glanduliferae

Germen apice connatum. *Stylus* lateralis.

Folliculus monospermus.

Diosmeae. *Smith.* exot. bot. 2. t. 63. *Lam.* ill.

Gärtn. t. 211. *Adr. Juss.* in *Mém. du mus.* 12. t. 20

555. OBOLARIA L. (1737.)

Cor. calycina, bracteis binis foliaceis suffulta, ~~can~~
lata, quadrifida, laciniis subincisis.

Stam. aequalia ex angulis corollae.

Stigma emarginatum.

Caps. unilocularis, bivalvis; polysperma.

Gentianeae. [*Orobanche Plukn.* t. 209. f. 6.]

556. MAIANTHEMUM Wiggers. (G. W. Weber. 1780.)

Cor. calycina quadripartita, rotata, subrevoluta.

Stamina basi corollae inserta, laciniis opposita.
rae ovales erectae.

Stylus filiformis, stigmate obtuso.

Bacca globosa, bi-trilocularis, di-trisperma.

Sarmentaceae. [*Convallaria* L. *Sturm.* fl. germ.

lacina *Desf.* *Sciophila Wibel.*]

557. ALCHEMILLA Trag. (1539.)

Cal. patens octofidus, laciniis alternis minoribus.

Discus glandulosus perigynus.

Stam. quaterna (bina in quatuor, unicum in una specie)
 laci calycis inserta. Antherae transverse dehiscentes.
Stylus lateralis, unicus in europaeis (2, 3, 4, 5 in ame-
 riana.)

Gynopsis solitaria in fundo calycis.

Rosaceae. Schk. t. 26. Sturm. fl. germ. Humb. nov. gen.
 6. t. 560. 561.

CONETES N. L. Burm. (1768.)

Indicrum tetraphyllum, triflorum.

Cor. calycina tetrapetala.

Stam. filamenta capillaria, corollam aequantia. Antherae
 subrotundae.

Stylus filiformis, stigmate trifido.

Cap. tricoeca.

— — Burm. fl. ind. t. 15. f. 5.

DAPHNITIS * (1825.)

Flores andioeci.

Cal. tetrasepalus, bracteatus.

Stylus simplex.

Fructus?

— — [Laurophyllus Thunb. ? Dilobeia Thuers.]

HEDYCARPUS Jack. (1825.)

Cal. quadri- (subinde quinque-) partitus.

Semina quatuor (quandoque quinque), cum lacinii ca-
 lycis alterna, Antherae bilobae.

Stylus brevissimus. Stigmata tria (quatuor) villosa.

Cap. baccata, trilocularis: loculis subdispermis. Semina
 integumento succulento cincta.

LAGENULA Lour. (1790.)

Cal. tetrasepalus, persistens.

Gyn. perigynus carnosus, quadrilobus.

Stam. filamentis calyci aequalibus, antheris incumbentibus.

Stylus crassus, stigmate simplici.

Bacca lagenaeformis bilocularis disperma.

II. FRUCTUS INFERI.

VISCUM Virgil. Plin.

Flores dioeci.

Cal. subnullus.

Cor. tetrapetala.

Antherae subsessiles, petalis adnatae.

Stigma sessile orbiculatum.

Bacca monosperma. Sem. pluribus embryonibus fetum.

Loranthaceae. Sturm. fl. germ. Rich. in ann. du mus. 12.
 t. 27. Mirbel. in ann. du mus. 16. t. 21.

563. *ELAEAGNUS* Tournef. (1719.)*Cal.* corollinus campanulatus, quadrifidus.*Stam.* calyci inserta.*Drupa* monosperma.*Elaeagnaceae.* Aeb. Rich. in act. soc. nat. paris.
Gärtn. t. 216.† 564. *MOLDENHAUERIA* * (1825.)

Flores dioeci.

Cal. tetrasepalus.*Stylus* radiatus.*Nux.*— — [*Cavanilla Thunb.*]565. *GYROCARPUS* Jacqu. (1763.)

Flores polygami.

Cal. quadri-octofidus.*Stam.* perigyna, laciniis calycis opposita. Anther
culares, loculis a basi dehiscentibus.*Stylus* brevissimus, stigmate capitato.*Drupa* apice bialata. Embryo inversus, cotyle
spiratim involutis.*Laurinae?* Roxb. corom. 1. t. 1. *Gärtn.* t. 97.566. *SANTALUM* Val. Cord. (1561.)*Cor.* calycina urceolata quadrifida.*Glandulae* nectariferae fauci insertae, cum sta
alternantes.*Stylus* simplex, stigmate tri-quadrilobo.*Drupa* apice marginata.*Santaleae.* [Sirium L. mant. *Lam.* ill. t. 74. *R*
rom. 1. t. 2.]567. *FUSANUS* L. Murr. (1774.)*Cal.* corollinus turbinatus, semiquadrifidus.*Stam.* filamentis linearibus, antheris quadrilobis.*Discus* perigynus glandulosus quadrilobus.*Stylus* brevissimus, stigmatibus quatuor.*Drupa* baccata globosa.*Santaleae.* *Lam.* ill. t. 73. [Colpoon *Berg.* cap.568. *OCTABILLUM* Lour. (1790.)*Cor.* hypocrateriformis, tubo tetragono brevi, limbo
dripartito carnoso.*Stam.* filamentis brevissimis fauci insertis, anthe
longis, bilocularibus, incumbenibus.*Stylus* turbinatus, stigmate crassiusculo.*Bacca* monosperma. Semen integumento octog
volutum.

— —

† 569. *CATONIA* P. Brown. (1756.)*Cal.* tetrasepalus, persistens.

Stam. filamentis calycem aequantibus, antheris subrotundis.

Stylus simplex.

Bacca disperma, calyce coronata.

370. GYMNOTACHYS R. Br. (1810.)

Flores in spadice cylindrico e spatha minuta.

Cal. quadripartitus.

Stam. basi calycis inserta.

Sigma, sessile.

Bacca monosperma.

D I G Y N I A .

I FLORES PERFECTI.

371. BEFFONIA Sauvag. L. (1747.)

Cal. tetrasepalus.

Petala quatuor integra.

Caps. compressa unilocularis, bivalvis, disperma.

Coryophylleae. Lam. ill. t. 87. Gärtn. t. 129.

372. HYPECOUM Dioscor.

Cal. disepalus.

Petala quatuor inaequalia, interiora saepe triloba.

Stamina filamentis capillaribus.

Caps. siliquaeformis, articulata: articuli monospermi, seminibus alternatim affixis.

Papavereae. Schk. t. 27. Gärtn. t. 115.

373. HAMAMELIS L. (1751.)

Cal. quadrididus, basi externa squamis binis suffultus.

Petala quatuor lingulata, basi squama lineari incumbente.

Stam. petalis alterna. Antherae bivalves subrotundae, valvis operculiformibus fere liberis.

Styli breves.

Caps. seminifera, bilocularis, apice bivalvis, valvis bifidis. Cocci monospermi, coriacei, bivalves.

Berberideis affinis Juss. *Fothergilleae* Nuttall. Schk. t. 27.

1. DICORYPHE Thuars. (1806.)

Cal. tubulosus, quadrilobus, basi persistente.

Petala quatuor.

Stam. petalis alterna, quaterna fertilia, cum totidem sterilibus alternantia. Antherae oblongae apice affixae.

Caps. suboctogona bicornis, inter stylorum vestigia dehiscens.

Cocci duo, in fundo haerentes, monospermi, apice stice dissilientes.

Hamamelidi affinis. Thuars. hist. ins. Afric. austr. t. 1.

575. PAGAMEA Aubl. (1775.)

Cal. turbinatus, quadridentatus.

Cor. urceolata, quadrifida, intus villosa.

Stam. Antherae subsessiles in tubo corollae.

Styli capillares.

Bacca calyci immersa, bilocularis: nuculis dispermis tero saepe abortiente.

Rubiaceae? Aubl. guian. t. 44.

576. SCHÄFFERA Jacqu. (1763.)

Flores dioeci.

Cal. quadripartitus, persistens.

Petala quatuor.

Stam. petalis opposita, filamentis erectis capillaribus theris erectis oblongis.

Stigmata subsessilia, bifida.

Bacca bilocularis, loculo altero abortiente.

Rhamnaceae? Swartz fl. ind. occ. t. 7. A.

577. NERTEA Banks. Gärtn. (1788.)

Cal. minimus, superus, quadridentatus.

Cor. infundibuliformis quadrifida.

Stam. filamentis basi corollae insertis.

Drupa baccata, di-tetrapyrena.

Rubiaceae. Gärtn. t. 26. Smith. ic. ined. 2. t. 28. danellae facie Feuill. 3. t. 44. Gomozia Mutis. L. Erythrodanum Thuars.]

578. MICROCHLAENA R. Br. (1810.)

Gramen paniculatum.

Involucrum bivalve minimum. Pedicellus barbatus.

Cal. bivalvis; uniflorus, valva utraque aristata suba-

Cor. bivalvis mutica, inaequalis.

Squamae duae hypogynae.

Caryopsis nuda.

[Ehrharta stipoides Labill. nov. holl. t. 118. Palis. agrost. t. 12. f. 6.]

II. FLORES IMPERFECTI.

579. ANTIARIS Leschen. (1812.)

Flores monoeci.

Involucrum cyathiforme, apice multifidum, multifloribus dense confertis, sessilibus.

Cal. tetrasepalus.

Antherae biloculares.

Styli filiformes.

Samina in drupam ex involucri aucto formatam confluentia.

Urticeae. *Leschen.* in ann. mus. 16. t. 22. *R. Brown* ad *Flinders* it. t. 5.

80. MORUS Virgil. Plin.

Flores monoeci amentacei.

Cal. quadripartitus.

Stam. filamentis erectis, antheris bilocularibus.

Stigmata duo sessilia.

Baccae cum calyce concreatæ, aggregatæ. *Caps.* sub-
disperma, embryo conduplicato.

Amentaceae — *Urticeae*. *Schk.* t. 290. *Gärtn.* t. 126.

81. PLANERA J. F. Gmel. (1791.)

Flores polygami.

Cal. campanulatus, membranaceus, quadri-quinquefidus.

Stam. perumque quaterna, exserta.

Stigmata duo glandulosa.

Nux coriacea, monosperma, squamulosa.

Amentaceae. [*Rhamnus* *Pall.* ross. 2. t. 40.]

82. CRUCITA Löffl. (1758.)

Cal. tetrasepalus, extus bracteis ternis munitus. Sepala
exteriora integerrima, interiora margine tenuissimo,
lacero.

Stam. filamentis capillaribus, calyce brevioribus.

Stigmata simplicia.

Caryopsis calyce tecta.

Chenopodeae.

83. MYRICA L. (1737.)

Flores amentacei dioeci.

Squamæ unifloræ.

Antheræ subsessiles, bipartitæ.

Styli filiformes.

Drupa baccata. Sem. exalbuminosum.

Amentaceae. *Schk.* t. 322. *Gärtn.* t. 39. [*Nagi Kämpf.*
amoen. exot. ic. p. 774. *Nageia* *Gärtn.* t. 39. sola albu-
minis præsentia differt.]

T R I G Y N I A.

† 584. BOSCHIA Thunb. (1794.)

Cal. quadridentatus.

Cor. tetrapetala.

Caps. quadrilocularis.

Terebinthaceae. [*Asaphea* *Caud.*]

85. PACHYSANDRA Mx. (1803.)

Flores spicati monoeci.

Cal. quadripartitus, bracteatus.

Stam. filamentis complanatis, antheris dorso affixis, cularibus.

Styli recurvi, persistentes.

Caps. trilocularis, loculis dispermis.

Triticaceae. Michaux fl. bor. am. t. 45. *Adr. Juss. E.* t. 1. f. 2. Bot. reg. 33.

586. *HYMENELLA* Sess. Cand. (1824.)

Cal. quadripartitus patens.

Petala quatuor, oblonga, integra, calycis longitudine.

Stam. petalis alterna, basi coronula petaloidea iuncta.

Caps. trilocularis.

Caryophylleae.

T E T R A G Y N I A.

587. *SAGINA* L. (1737.)

Cal. quadripartitus.

Petala quatuor, calyce minora, aut nulla.

Caps. unilocularis, quadrivalvis, polysperma.

Caryophylleae. Schk. t. 27. *Sturm.* fl. germ. 880. 881.

588. *MÖNCHIA* Ehrh. (1788.)

Cal. tetrasepalus, persistens.

Petala quatuor, lanceolata, calyce breviora.

Stam. filamenta subulata. Antherae bilobae.

Caps. unilocularis, apice dehiscens, octodentata, polysperma.

Caryophylleae. [*Sagina erecta* L. Engl. bot. 600.]

589. *RADIOIA* Dillen. (1719.)

Cal. quadripartitus, laciniis inciso-dentatis.

Petala quatuor.

Caps. octococca, coccis monospermis.

Lineae. Engl. bot. 893.

590. *BULLIARDA* Cand. (1799.)

Cal. quadripartitus.

Petala quatuor.

Squamae lineares ad basin petalorum.

Folliculi quatuor polyspermi, intus dehiscentes.

Crassuleae. Cand. succul. t. 74. [*Sedum minimum* V. bot. par. t. 10. f. 2. *Tillaea* fl. dan. 1510.]

591. *BRUCEA* J. S. Miller. (1780.)

Flores polygami.

Cal. quadripartitus.

Petala quatuor.

Discus perigynus, quadrilobus.

Stigmata quatuor papillosa.

Drupae quatuor monospermae.

Terebinthaceae. Herit. stirp. 1. t. 10. [Gonus et tradium *Lour.*]

6. *HYGINDA* Jacqu. (1763.)*Cal. minimus, quadrifidus.**Cor. quatuor patentia.**Stam. cum petalis alternantia.**Stigmata quatuor.**Superiora monosperma.**Celastrinae. Jacqu. ic. 2. t. 311. Humb. nov. gen. 7. t. 620. [Crossopetalum P. Braun jam. t. 17. f. 1. Rhaecosa L. 1759.]*7. *ILLEX* Lonicer. (1564.)*Cal. quadridentatus persistens.**Cor. quadrifida vel tetrapetala.**Stam. petalis alterna.**Stigmata vel distincta vel in unum confusa.**Bacca tetrapyrena, nucleis monospermis.**Celastrinae. Schk. t. 28. Gärtn. t. 92. [Paltaria R. et P. Macoucota Aubl.]*8. *ALVERENIA* Ruiz. et Pav. (1798.)*Cal. quadripartitus, bracteis duplicibus bifidis.**Petalis quatuor concava.**Filamenta compressa.**Cor. quadrilocularis, quadrivalvis, stylis coronata, polysperma.*9. *CICCA* L. (1767.)*Flores dichini.**Cal. quadripartitus, intus glandulis quatuor alternis, discove glanduloso instructis.**Cor. nulla.**Stam. filamentis liberis, antheris bilocularibus.**Styli quaterni, bifidi.**Fructus baccatus, quadricoccus, coccis submonospermis.**Tricoccae. Adr. Juss. Euphorb. t. 4. f. 13. [Averrhoa acida L.]*10. *POTAMOGETON* Dioscor.*Cor. calycina tetrapetala.**Stam. cum petalis alternantia. Antherae biloculares.**Nuculae quatuor.**Alismaceae. Schk. t. 28. Tittmanns Keimung, t. 6. [Hydrogeton Lour.]*11. *RUPPIA* L. (1737.)*Flores spicati, alternatim ad utrumque rhacheos latus dispositi.**Cal. nullus.**Cor. nulla.**Antherae quaternae biloculares, squamis ephippioides suffultae. Pollen cylindricum geniculatum.*

Germina quaterna obconica, *Nuculae* gibbae, dein pedunculatae. Semen totum albuminosum rostellum hoc cotyledonem Richardus et Trevis albumen radicem habet Richardus, vitellum Gärtz

Najades. Hook. in fl. loudin. 4. t. 53. *Reichenb.* t. 174. Fl. dan. 364. *Richard* in ann. mus. 17. t. 5. 43. *Treviran.* Entwickel. des Eyes, t. 1. f. 1-11.

598. OTTONIA* (1819.)

Flores amentacei, racemosi, squamis distincti.

Cal. nullus,

Cor. nulla,

Antherae ovatae, biloculares,

Stigmata quatuor,

Achenium quadrangulare. Albumen corneum.

axilis rectus, radícula supera.

Pipereae. Grundzüge, t. 3. f. 4. 5.

CLASSIS V. PENTANDRIA

CONSPECTUS

CLASSIS QUINTAE.

MONOGYNAE.

- Fructus perfectis.
 1. Genia monopetala.
 A. Fructibus superis,
 a. monospermis,
 1. Caryopseis, achenia vel utriculi 599 — 608.
 2. Nucis 609 — 634.
 3. Drupae aut baccae 635 — 645.
 4. Fructus follicularis 646.
 b. Fructibus dispersis.
 1. Membranaceis, 647 — 649.
 2. Drupis vel baccis 650 — 656.
 4. Fructibus trispermis.
 1. Membranaceis 656 — 658.
 2. Drupaceis, baccatis 659, 660.
 4. Fructibus tetraspermis.
 1. Membranaceis, 661 — 666.
 2. Drupaceis, baccatis 667 — 678.
 4. Fructibus penta-decapermis.
 1. Membranaceis, 679 — 681.
 2. Drupaceis, baccatis, 682 — 698.
 4. Fructibus polyspermis.
 1. Membranaceis,
 aa. unilocularibus.
 aa. Capsulis, 699 — 721.
 bb. Fructibus follicularibus geminis 722 — 742.
 bb. Capsulis aut folliculis bilocularibus, 743 — 762.
 yy. Capsulis trilocularibus, 763 — 769.
 zz. Capsula quadriloculari 770.
 aa. Capsulis quinquelocularibus, 771 — 782.
 2. Fructibus baccatis,
 aa. unilocularibus, 783 — 790.
 bb. bilocularibus, 791 — 803.
 yy. quadri-quinque-multilocularibus 804 — 808.
 3. Fructibus ignotis, 809 — 814.
 B. Fructibus inferis,
 a. capsularibus.
 a. Capsulis unilocularibus, 815 — 819.
 b. Capsulis bilocularibus 820 — 845.
 y. Capsulis tri-quinquelocularibus, 846 — 849.
 b. Fructibus baccatis, drupaceis,
 a. mono-dispermis, 850 — 866.
 b. tri-penta-hexaspermis, 867 — 879.
 y. polyspermis, 880 — 896.
 I. Corolla pleiopetala.
 A. Fructibus superis.
 a. Corolla tetrapetala 897.
 b. Corolla pentapetala.
 4. Fructibus mono-dispermis 898 — 913.
 b. Fructibus tri-pentaspermis 914 — 947.
 y. Fructibus deca-polyspermis 948 — 982.

B. Fructibus inferis.

a. Corolla tripetala 983.

b. Corolla pentapetala 984 — 997.

** Floribus imperfectis.

A. Fructibus superis, 998 — 1042.

B. Fructibus inferis, 1043 — 1051.

D I G Y N A E.

* Floribus perfectis.

I. Corolla monopetala: fructibus superis 1052 — 1113.

II. Corolla pentapetala.

A. Fructibus superis 1114 — 1117.

B. Fructibus inferis 1118 — 1191.

** Floribus imperfectis 1192 — 1202.

T R I G Y N A E.

* Floribus perfectis.

I. Fructibus superis.

A. Corolla monopetala 1203.

B. Corolla pentapetala 1204 — 1224.

II. Fructibus inferis 1225 — 1230.

** Floribus imperfectis 1231 — 1238.

T E T R A G Y N A E.

1239.

P E N T A G Y N A E.

* Floribus perfectis.

I. Fructibus superis 1240 — 1252.

II. Fructibus inferis 1253. 1254.

** Floribus imperfectis 1255. 1256.

P O L Y G Y N A E.

1257 — 1259.

M O N O G Y N I A.

* *Flores perfecti.*

I. COROLLA MONOPETALA.

A. *Fructus superi.*a. *monospermi.*1. *Caryopses, achenia vel utriculi.*

599. MIRABILIS Parkins. (1640.)

Involucrum quinquefidum, calyciforme, uniflorum.

Cor. calycina, infundibuliformis, basi ventricosa persi
Urceolus nectarifer filamenta promit, post anthes
crassatus seminis integumentum constituit.*Stamina* limbum corollae subaequantia. *Antherae*
tundae.*Stylus* simplex, stigmatе capitato.

Nucula monosperma, embryo peripherico, albumine centrali.

Nyctagineae. C. K. Sprengels entdeckt. Geheiman. t. 8. f. 16. Smith. exot. bot. 1. t. 23. Gärtn. t. 127.

ABRONIA Juss. (1789.)

Endocrum pentaphyllum, multiflorum.

Or. calycina, infundibuliformis: basi ventricosa, persistente, angulata: limbo quinquelobo.

Urceolus perigynus staminifer, post anthesin cum fructu coalescens.

Stamina inclusa, antheris oblongis.

Stylus simplex, stigmatibus subclavato.

Nucula angulata, monosperma. Embryo convolutus: albumen centrale.

Nyctagineae. Lam. ill. t. 105. Hooker exot. fl. t. 193. 194. Bot. mag. t. 2879. Gärtn. t. 214. [Tricratus Herit.]

TRICICLA Cav. (1800.)

Endocrum triphyllum uniflorum, foliis subrotundis venosis cordatis superantibus.

Or. calycina, infundibuliformis, limbo quinquelobo.

Stam. filamenta hypogyna, basi libera: antherae oblongae.

Stylus simplex abbreviatus.

Semen oblongum in basi corollae persistente inclusum.

Embryo periphericus, cum albumine centrali.

Nyctagineae. Cav. ic. 6. t. 598. Anal. de cienc. nat. 5. t. 40.

PLUMBAGO Plin. Clus. (1576.)

Cal. tubulosus, pentagonus, glandulosus, quinqueidentatus.

Cor. hypocrateriformis, limbo quinquefido.

Stam. filamentis receptaculo insertis, basi dilatatis.

Stylus simplex, stigmatibus quinquefido.

Semen unicum, filamentorum basibus valvaeformibus inclusum, styli basi conica coronatum, ex apice funiculi umbilicali curvati pendulum. Albumen periphericum.

Plumbagineae. Schk. t. 36. Gärtn. t. 59.

VOGELIA Lam. (1792.)

Cal. pentasepalus, sepalis ovatis magnis transversim undulato-sulcatis.

Cor. tubulosa, gracilis, plicata, calycem excedens, limbo quinquelobo.

Stam. hypogyna, filamentis capillaribus.

Stylus simplex, stigmatibus quinquefido.

Fructus?

Plumbagineae? Lam. ill. t. 149.

WEIGELIA Thunb. (1780.)

Cal. pentasepalus, sepalis aequalibus subulatis.

Cor. campanulata, tubo intus villosa, limbo erecto, semi-quinquefido.

- Stam.* filamenta tubo inserta, corollam fere aequant.
Antherae erectae, lineares, basi bifidae.
Stylus e basi germinis, filiformis, corollam superius
 stigmate peltato.
Sem. unicum nudum?
 — — *Thunb.* in act. holm. 1780. t. 5. fl. japon.
Kämpfer ic. select. t. 45.
605. **QUINCHAMALA W. (1798.)**
Cal. brevissimus, globosus, persistens, inaequaliter
 quinqueidentatus.
Cor. tubo gracili cylindrico, limbo quinquefido.
Stam. *Antherae* lineares, subsessiles, fauci corollae insertae.
Stylus simplex, stigmate capitato.
Caryopsis globosa, calyce persistente tecta.
 — — [Quinchamali *Deuill.* obs. 3. t. 44. *Lam.* ill. t. 44
Arjuna Cav. ?]
606. **ULLUCUS Lozan. (1809.)**
Cal. disepalus, coloratus, deciduus.
Cor. profunde quinquepartita, tubo brevissimo.
Stam. filamentis brevissimis: *antherae* erectae, biloculares.
Stylus filiformis, stigmate simplici.
Utriculus monospermus.
Portulacaceae.
607. **BRUNONIA Sm. (1810.)**
Capitulum involucreatum.
Cal. quinquefidus, bracteis quaternis suffultus.
Cor. infundibuliformis, limbo quadripartito, laciniis, quibus
 superioribus altius divisus.
Stamina hypogyna: *antherae* connatae.
Stigmatis indusium bivalve.
Utriculus monospermus, inclusus tubo calycis aucto
 durato, laciniis plumosis. *Sem.* exalbuminosum.
 — — *Smith* in *Linn.* transact. 10. t. 28. 29. *Bauer* ill. t.
608. **CORYMBIUM L. (1737.)**
Cal. basi bracteatus, prismaticus, disepalus: sepalis co
 niventibus, apice truncatis, obsolete tridentatis.
Cor. tubo brevissimo, limbo quinquepartito.
Stam. filamenta tubo insidentia: *antherae* in cylindri
 coalitae.
Stylus longitudine corollae, stigmate bipartito.
Caryopsis oblonga, lana papposa cincta et coronata
 calyce inclusa.
Aggregatae. *Burm.* afr. t. 70. *Gärtn.* t. 86.

2. Nuces.

ASPERIFOLIAE.

Boragineae Juss. *Familia plantarum herbacearum, frutico
 rum, raro arborescentium, labiatis proxima, maxime fruc*

bus quaternis. Sed differt defectu gynobaseos carnosae et embryonis situ inverso.

α. Binae.

CERINTHE Theophr. Plin.

Cal. pentasepalus inaequalis.

Cor. tubuloso - campanulata, limbo quinqueidentato.

Stam. filamenta crassiuscula tubo inserta. Antherae hastatae, auriculis filiformibus tortis antheras secum necantibus.

Nuces duae biloculares.

Sturm. fl. germ. Gärtn. t. 67.

β. Nuces quaternae,

aa. uniloculares,

aaa. cohaerentes.

COLDENIA L. (1747.)

Cal. quinquepartitus.

Cor. infundibuliformis, fauce ampliata, limbo plano quinquefido.

Stam. filamenta fauci inserta; antherae ovales.

Sylus stigmatate bifido.

Nuces apice subulatae.

Lam. ill. t. 89. [Lithospermum dichotomum R. et P. flor. peruv. 2. t. 3. f. 6. Tiquilia Pers.]

HELIOTROPIMUM Theophr.

Cal. quinquefidus.

Cor. hypocrateriformis, fauce pervia (interdum barbata), limbo quinquelobo, plicato.

Stam. inclusa.

Stigma conicum.

Schk. t. 29. Gärtn. t. 68.

ββ. Nuces distinctae,

aaa. Fundo calycis affixae,

aaaa. Faux corollae pervia.

LITHOSPERMUM Dioscor.

Cal. quinquepartitus.

Cor. infundibuliformis, limbo quinquelobo, fauce quandoque barbata.

Stam. inclusa.

Nuces basi planae, ovatae, plerumque lapideae.

Sturm. fl. germ. Jacq. ecl. 2. t. 136. Gärtn. t. 67. [Batschia J. F. Gmel., Arnebia Forsk., Mertensia Roth.]

613. *PULMONARIA* Plin.*Cal.* prismaticus, quinquefidus.*Cor.* campanulata limbo quinquefido, fauce subbarba*Stigma* emarginatum.*Nuces* turbinatae, basi planae.*Reichenb.* ic. 6. t. 501 — 506.614. *PURSHIA* * (1817.)*Cal.* quinquepartitus.*Cor.* tubulosa, limbo ventricosso connivente.*Stam.* filamentis inclusis: antheris sagittatis distinctis.*Stylus* corolla duplo longior.*Nuces* turbinatae, basi planae.[*Onosmodium Michaux* fl. bor. amer. t. 15.]615. *MOLTKIA* Lehm. (1817.)*Cal.* quinquepartitus.*Cor.* infundibuliformis.*Stam.* exserta, antheris oblongis incumbentibus.*Nuces* magnae, difformes, basi planae.*Lehmann* ic. t. 43. 44. [*Nonea phanerantha Vivian.*
lib. t. 1. f. 3.]616. *ONOSMA* Dioscor.*Cal.* quinquepartitus.*Cor.* tubuloso-campanulata.*Stam.* inclusa, vel corollam subaequantia: antherae
lateraliter nexae.*Nuces* ovatae, lapideae, basi planae.*Kitab.* plant. hung. 3. t. 279. *Desfont.* atlant. 1. t. 43617. *ECHIMUM* Numen. Heracleensis.*Cal.* quinquepartitus.*Cor.* campanulata, limbi laciniis inaequalibus.*Stam.* longitudine varia.*Nuces* gigartodes, angulatae, basi applanatae.*Sturm.* fl. germ. *Gärtn.* t. 67.618. *ECHIOCHILON* Desf. (1800.)*Cal.* quadripartitus.*Cor.* tubo gracili incurvo: limbo bilabiato, labio su-
bi-infer. trilobo.*Nuces* glabrae, tuberculatae.*Desfont.* fl. atl. t. 47.619. *LYCOPSIS* Dioscor.*Cal.* quinquefidus, in fructu inflatus, nutans.*Cor.* infundibuliformis, fauce barbata.*Antherae* in tubo subsessiles.*Nuces* turbinatae, basi foratae.*Reichenb.* ic. 4. t. 330. *Gärtn.* t. 67. [*Nonea Cand.*]620. *COLSMANNIA* Lehm. (1818.)*Cal.* quinquepartitus, maximus, persistens, basi pent-
nus, laciniis oblongis erectis, coloratis.

Cor. campanulata, calyce brevior, limbo quinquedentato.
Stam. filamenta abbreviata: antherae sagittatae, conni-
 ventae.

Stylus corollae longior, stigmate obtuso.
Noces triangulares, osseae, basi foratae.
 Berl. Bot. mag. 8. t. 1.

FRANIOSPERMUM Lehm. (1818.)

Cal. oblongus, quinquepartitus, laciniis aequalibus.

Cor. cylindrica, limbo erecto quinquedentato.

Stam. filamenta capillaria, corolla duplo longiora: antherae
 incumbentes.

Noces hemisphaericae, basi foveolatae.

Lehmans ic. t. 50.

DIOCLEA * (1824.)

Cal. quinquepartitus.

Cor. indubuliformis, villosa, tubo gracili; limbo patulo,
 quinquelobo.

Stam. tubum aequantia, antheris oblongis.

Stylus bifidus, stigmatibus capitatis.

Noces pyramidatae, basi foveolatae.

[*Anchusa asperima Delil.*]

βββ. *Faux corollae fornicibus clausa*.

MYRSOTIS Diosc.

Cal. quadrifidus vel quadripartitus, post anthesin clausus
 vel apertus.

Cor. hypocrateriformis, fauce squamis instructa.

Stam. inclusa.

Noces basi foveolatae, glabrae vel scabrae, rugulosae.

Sturm. fl. germ. Gärtn. t. 68.

EXARRHENA R. Br. (1810.)

Cal. quinquefidus.

Cor. hypocrateriformis: fauce fornicibus coarctata: limbo
 quinquelobo.

Stam. exserta, antheris peltatis.

Stylus longitudine staminum: stigma simplex.

Noces basi umbilicatae.

BORRAGO Braunschweig. (1500.)

Cal. quinquepartitus.

Cor. rotata, limbo quinquelobo. Faux fornicibus emar-
 ginatis et filamentis lanceolatis alternis obsessa.

Antherae incumbentes, oblongae vel lanceolatae, conni-
 ventae.

Noces turbinatae, basi planae.

Sturm. fl. germ. Bot. reg. 288.

26. **ANCHUSA** Theophr.

Cal. quinquepartitus, fructiferus erectus vel nutans inflatus.

Cor. infundibuliformis, fauce fornicibus erectis clausa.

Stam. tubo adnata, inclusa.

Stylus, stigmatē bilobo.

Nuces turbinatae, basi foveolatae.

Engl. bot. 662.

627. *Stomotechium* Lehm. (1818.)

Cal. quinqueangularis, quinquefidus.

Cor. tubuloso-subcylindrica: fauce fornicibus subre carnosus, extus muricatis clausa.

Stam. inclusa: antherae oblongae acuminatae.

Stylus filiformis, stigmatē obtuso.

Nuces subrotundae, basi foveolatae.

628. *Symphytum* Dioscor.

Cal. quinquepartitus.

Cor. cylindrico-campanulata, fauce fornicibus sub in conum conniventibus, clausa: limbo quinque-

Stam. filamentis tubo adnatis: antherae lanceolatae minatae cum fornicibus alternantes.

Stylus simplex, stigmatē obtuso.

Nuces ovatae, basi foveolatae.

Schk. t. 30. Gärtn. t. 67. Reichenb. ic. t. 220.

629. *Lobostema* Lehm. (1828.)

Cal. quinquepartitus.

Cor. infundibuliformis.

Fornices erectae, dorso staminiferae.

Nuculae quatuor, basi non perforatae.

bbb. *Nuces columnae centrali affixae, basi plan-*

630. *Trichodesma* R. Br. (1810.)

Cal. quinquepartitus.

Cor. subrotata: fauce nuda; limbo quinquepartito, lobatis mucronato-cuspidatis.

Stam. filamenta brevissima. Antherae exsertae, actae, in conum apice spiralem collectae, villis dorsifariis cohaerentes.

Nuces columnae centrali quadrialatae foveolis semmersae.

Jacqu. ic. rar. 2. t. 314. [Pollichia Medic.]

631. *Asperugo* Plin.

Cal. in fructu compressus, lamellis parallelis sinuatis.

Cor. subinfundibuliformis: fauce fornicibus clausa.

Stam. filamenta brevissima, tubo adnata: antherae subrotundae cum fornicibus alternantes.

Nuces compressae, columnae centrali affixae.

Schk. t. 31.

62. **CYNOGLOSSUM** Dioscor.*Cal.* quinquepartitus.*Cor.* infundibuliformis vel subrotata: fauce fornicibus erectis instructa.*Stam.* filamenta brevissima, tubo adnata: Antherae ovals cum fornicibus alternantes, vel exsertae.*Nuces* vel echinatae solidae: vel laeves marginatae, seu membrana plana aut urceolata cinctae.

Schk. t. 30. Gärtn. t. 67. [Omphalodes Tourn. Rindera Pall.]

63. **ECHINOSPERMUM** Sw. Lehm. (1818.)*Cal.* quinquefidus.*Cor.* hypocrateriformis: fauce fornicibus brevibus instructa: limbo quinquelobo.*Nuces* echinatae, columnae centrali affixae.*Stam.* fl. germ. [Myosotis L. Gärtn. t. 68.]bb. *Nuces* biloculares.64. **TIARIDIMUM** Lehm. (1818.)*Cal.* quinquefidus.*Cor.* hypocrateriformis: tubo angulato: fauce coarctata, quinquerradiata: limbo quinquelobo undulato.*Stylus* brevissimus, stigmate capitato.*Nuces* quatuor biloculares, mitraeformes, cohaerentes, basi clausae.

Lehm. ic. t. 6. [Heliotropium indicum L. Bot. mag. 1837.]

3. *Drupae, aut baccae.*65. **MONOTOCA** R. Br. (1810.)*Cal.* quinquepartitus.*Cor.* infundibuliformis, limbo quinquefido fauceque imberbibus,*Discus* perigynus cyathiformis lobatus.*Stam.* filamenta tubo inserta: antherae incumbentes, bivalves, subexsertae.*Stylus* simplex, stigmate capitato.*Drupa* baccata monosperma.*Epacridae.* [Styphelia glauca Labill. nov. holl. t. 61.
? Needhamia R. Br.]66. **VALLERIA** Ruiz et Pav. (1798.)*Cal.* quinquefidus.*Cor.* infundibuliformis, fauce inflata; limbo plano, aequaliter quinquefido, laciniis lanceolatis acutis, subobliquis.*Stam.* filamenta brevissima, tubo adnata: antherae sagittatae, faucem aequantes.*Stylus* filiformis, stigmate oblongo.

Drupae duae, monospermae; putamine ligneo, fibre reticulato.

Contortae. R. et P. fl. peruv. 2. t. 151. Humb.
gen. 3. t. 241. [*Rauwolfia glabra* Cav. ic. 3. t. 297.]

637. *CORDIA* Plum. (1703.)

Cal. tubulosus, quinque- (interdum quadri-) dentatus.

Cor. infundibuliformis vel campanulata, limbo plano
quinque- (raro septem-) fido: fauce pilosa aut nuda.

Stam. filamenta breviter fauci imposita: antherae biloculares.

Stylus exsertus, bifidus, stigmatibus quaternis, variis.

Germen subquadriloculare.

Drupa monoperyna, 1-3 locularis: loculis duobus
sistentibus.

Asperifoliae. Delil. fl. aeg. t. 19. 20. Humb. nov.
t. 207. Roxb. corom. 1. t. 58. [Varronia L.
journ. de bot. 1. t. 9. Patagonula L. Cerdana R.]

638. *ERYCIBE* Roxb. (1798.)

Cal. quinqueidentatus.

Cor. infundibuliformis, limbo quinquelobo, lobis bilobis.

Stam. filamenta brevissima: antherae sagittatae.

Stylus brevissimus; aut nullus, stigmate quinquesulcato.

Bacca cerasiformis, monosperma.

Convolvuleae. Roxb. corom. 2. t. 159.

639. *ARDISIA* Sw. (1788.)

Cal. quinquefidus, quinquepartitus.

Cor. hypocrateriformis, tubo brevissimo: laciniis limbi
sistentibus subreflexis.

Stam. filamentis subulatis tubo corollae insertis: Antherae
lanceolatae, conniventes.

Stylus subulatus, stigmate simplici.

Bacca monosperma: seminis cortice fragili. Embryo
peritremis transversus in albumine corneo.

Myrsineae. Swartz. fl. ind. occid. 1. t. 10. Roxb. corom.
t. 27. Bot. reg. 533. 638. 827. [Anguillaria Gärtn. t. 7.
Icacorea Aubl. guian. t. 368. Pyrgus Lour.]

640. *MYRSINE* L. (1737.)

Flores polygami, dioeci.

Cal. quinquefidus (subinde quadrifidus).

Cor. quinque- (nonnunquam quadri-) partita.

Stam. laciniis corollae opposita. Antherae erectae, bi-
locares.

Stigma saepe lobatum aut laciniatum.

Bacca drupacea, pisiformis, putamine crustaceo. Embryo
transversus, flexuosus in medio albumine corneo.

Myrsineae. Venten. Cels. t. 86. Humb. nov. gen. 1.
t. 245. [Rapanea Aubl. guian. t. 46. Athruphyllum Lou.
Römeria Thunb. Scleroxylon W. Manglilla et Badu
Juss. Caballeria R. et P.]

EMBELIA N. L. Burm. (1768.)*Cal.* quinquefidus.*Cr.* profunde quinquepartita, pentapetaloidea.*Stam.* laciniis corollae opposita. Filamenta basi corollae inserta, elongata, patula. Antherae peltatae, biloculares.*Stigma* truncatum.*Bacca* Myrsines.*Myrsineae.* N. L. Burm. fl. ind. t. 23.**GLADHIA** Thunb. (1781.)*Cal.* quinquepartitus, patens, laciniis inciso-serratis.*Cor.* rotata quinquepartita.*Stam.* filamenta brevissima, fauci inserta. Antherae cordatae, acutae in conum conniventes.*Stylus* filiformis, corolla longior: stigmatе acuto.*Bacca* monosperma, stylo persistente coronata. Sem. in tegumento laxo involutum.*Myrsineae.* Thunb. fl. jap. t. 18. Gärtn. t. 211.**BURHELIA** Theophr. Sw. (1788.)*Cal.* quinquepartitus.*Cor.* quinquepartita, tubo brevissimo. Squamae duae ad basin cuiusque lacinae.*Stam.* tubo inserta, laciniis opposita. Antherae subcordatae biloculares. Squamae petaloideae accessoriae, perumque denticulatae, cum staminibus alternantes et ante ea positae.*Styl.* persistens, stigmatе simplici. Germen pentaspermum.*Drupa* monosperma, stylo coronata.*Sapotaeae.* Swartz fl. ind. occ. t. 8. Gärtn. t. 202.*Humb.* nov. gen. 7. t. 617.**SIDEROXYLON** Dill. (1732.)*Cal.* parvus, quinquefidus, persistens.*Cor.* rotata, quinqueloba. Appendices aut squamae petaloideae serratae ad basin loborum corollae, cumque iis alternantes.*Stam.* lobis corollae breviora. Filamenta subulata. Antherae incumbentes.*Styl.* simplex, inferiore parte persistens.*Bacca* monosperma. Sem. testa ossea vestitum, cavitate una aut altera exsculptum. Albumen crassum, subcartilagineum.*Sapotaeae.* Lam. ill. t. 120. Venten. sel. t. 22. Gärtn. t. 202. [Sersalisia R. Br.]**THELA** Lour. (1790.)*Cal.* tribracteatus, tubulosus, subquinquefidus, coloratus, papillosus.*Cor.* hypocrateriformis, quinquefida.*Stam.* receptaculo inserta. Filamenta capillaria. Antherae oblongae erectae.

Styl. filiformis. Stigma quinquepartitum.
Bacca pentagona, monosperma.

— — 4. *Fructus follicularis.*

646. *AEGICERAS* Gärtn. (1788.)

Cal. pentasepalus, coriaceus, persistens.
Cor. infundibuliformis, quinquepartita.
Stam. ex annulo imi tubi corollae, exserta. *Antherae* longae biloculares, superne dehiscentes.
Styl. persistens. Stigma simplex.
Folliculus arcuatus, cylindricus, acuminatus, monospermus. Funiculus umbilicalis in arillum calyptraeformis abiens. Cotyledones minimae. Radicula maxima.
 — — Gärtn. t. 46. König in Ann. of bot. t. 1.
 [Rhizophora corniculata L.]

b. *Fructus dispermi,*

1. *membranacei.*

647. *POLYMERIA* R. Br. (1810.)

Cal. quinquepartitus.
Cor. infundibuliformis, plicata.
Styl. stigmatibus quaternis, senis, acutis.
Caps. disperma.

Convolvuleae.

648. *BONAMIA* Thuars. (1806.)

Cal. quinquepartitus, persistens.
Cor. campanulata, quinquefida.
Stamina tubo inserta, basi pilosa. *Antherae* dorso oblongae, biloculares.
Styl. bifidus, exsertus. Stigmata capitato-lobata.
Caps. bilocularis, disperma.

Thuars. ins. Afr. austr. t. 5.

649. *MURUOIA* Aubl. (1775.)

Cal. turbinatus, quinquelobus, persistens.
Cor. infundibuliformis, limbo patente, amplo, quinquefido.
Stam. fauci imposita. *Antherae* incumbentes.
Styl. elongatus. Stigma bilamellatum.
Caps. coriacea, bilocularis. Sem. solitaria, limbo vexa, inde plana.

Convolvuleae. Aubl. guian. t. 54. [Convolvulus spermus Willd.]

2. *Drupae vel baccae.*

650. *CERBERA* Mill. (1759.)

Cal. quinquepartitus.
Cor. infundibuliformis, fauce quinqueidentata, limbo quinquefido.
Stam. *Antherae* in summo tubo subsessiles, stigmatibus haerentes.
Styl. stigmatibus peltato, lobato.

Drupae duae exsuccae, putamine semibivalvi.

Contortae. Jacq. amer. t. 34. Gärtn. t. 123. 124.
[Ochrosia Jacq. ic. rar. 2. t. 321. Labill. nov. Caled.
t. 30.]

ARDUINA Mill. (1759.)

Cal. quinquepartitus, persistens.

Cor. subinfundibuliformis, limbo quinquesido.

Stam. inclusa.

Stylus bifidus.

Bacca disperma.

Contortae. Mill. ic. 2. t. 300.

CORTESIA Cav. (1797.)

Cal. campanulatus, truncatus, decemdentatus.

Cor. infundibuliformis, limbo quinquelobo.

Stam. exserta, filamentis basi dilatatis, tubo insertis;
antheris incumbentibus.

Stylus staminibus longior, bipartitus: stigmatibus glo-
bosis.

Bacca disperma: semina hinc plana inde convexa.

Convolvulaceae. Cav. ic. 4. t. 377.

HUNTERIA Roxb. (1821.)

Cal. quinquepartitus.

Cor. infundibuliformis.

Stigma bilobum.

Baccae duae, mono-dispermae.

Sapoteae Juss.

GÄRTNERA Lam. (1791.)

Cal. campanulatus, quinquedentatus, basi bracteis binis
suffultus.

Cor. infundibuliformis, limbo quinquepartio.

Stam. inclusa, tubo inserta; filamentis brevissimis; anthe-
ris oblongis, erectis.

Stylus exsertus, apice bifidus.

Bacca drupacea, dipyrena.

Loganiaceae R. Brown. Lam. ill. t. 167. Gärtn. t. 191.

DICARYUM W. herb. (1819.)

Cal. quinquedentatus, persistens.

Cor. quinquedentata, fauce pilosa.

Stam. fauci inserta.

Stylus simplex, stigmatibus obtuso.

Drupa disperma: nuce biloculari.

Strychnaceae Cand.

c. *Fructus trispermi.*1. *Membranacei.*656. *PHLOX* L. (1751.)*Cal.* prismaticus, quinquefidus.*Cor.* tubo elongato, limbo hypocrateriformi, quinque-*Stam.* inaequalia, tubo inserta.*Caps.* trilocularis, trivalvis, subtrisperma.*Polemoniae.* Schk. t. 38. Gärtn. t. 62.657. *BONPLANDIA* Cav. (1800.)*Cal.* tubulosus, quinqueidentatus.*Cor.* infundibuliformis, limbo quinquepartito, subbil-*Stam.* inaequalia, declinata, exserta.*Caps.* trilocularis, trivalvis, trisperma. Sem. mucosa.*Polemoniae.* Cav. ic. 6. t. 532. [*Caldasia Willd.* berol. t. 71.]658. *CRYPTANDRA* Sm. (1798.)*Cal.* pentasepalus.*Cor.* tubulosa, quinquefida: squamis quinque cucullatis
stamina in fauce corollae tegentibus.*Stigma* trifidum.*Caps.* trivalvis, valvis inflexis dissepimenta formantibus.
Sem. terna.*Rhamnaceae.* Rudg. in Linn. transact. 10. t. 18.2. *Fructus drupacei, baccati.*659. *CRYPTOSTOMUM* Schreb. (1789.)*Cal.* infundibuliformis, ventricosus, coloratus: limbo
quinquefido.*Cor.* infundibuliformis, tubo brevissimo calycis faucibus
serto: limbo quinquefido connivente.*Fornix* stipitatus, quinqueidentatus, orificium corollae
claudens.*Stam.* filamenta nulla. Antherae sessiles sub dentibus
fornicis.*Stylus* cylindricus, stigmatibus capitato.*Bacca* drupacea, trilocularis. Sem. in quovis loculo
litaria, mucilagine cincta.*Büttneriae* affine. [*Moutabea Aubl. guian. t. 274.*]660. *LANDOLPHIA* Pal. Beauv. (1806.)*Cal.* imbricatus, coriaceus.

Cor. tubulosa, fauce villosa, limbo quinquefido, obliquo.
Stam. tubo inserta.

Stylus apice incrassatus, stigmate bifido.

Caps. unilocularis, trisperma.

Contortae. Pal. Beauv. fl. owar. t. 34.

d. *Fructus tetraspermi.*

1. *Membranacei.*

HYDROPHYLLUM Joncquet. (1659.)

Cal. quinquepartitus.

Cor. tubulosa, limbo erecto quinquelobo.

Stam. filamenta tubo imo inserta, medio pilosa, basi laciniis corollae binis prominulis nectariferis munita. Antherae exsertae incumbentes.

Stylus simplex, stigmate bifido.

Caps. unilocularis, bivalvis, tetrasperma. Sem. albuminosa.

Hydrophyllaeae R. Br. Bot. reg. 244. 331. Gärtn. t. 110.

PHACELIA Juss. (1789.)

Cal. quinquepartitus.

Cor. subcampanulata, limbo erecto.

Stam. exserta: filamenta e plicis tubi corollae in sulcos absentibus surgentia: antherae incumbentes.

Stylus bifidus, stigmatibus acutis.

Caps. bivalvis, valvis dissepimentum formantibus, tetrasperma.

Hydrophyllaeae. Michaux fl. bor. amer. t. 16. Jacq. fil. ecl. t. 91. [Aldecoa R. et P. fl. 2. t. 14.]

NEMOPHILA Barton. (1818.)

Cal. patens, decempartitus, laciniis alternis reflexis.

Cor. campanulata, limbo quinquefido.

Stam. corolla breviora: filamenta tubo inserta: scrobiculi gemini cuiusvis filamenti basi adstantes. Antherae incumbentes, versatiles.

Stylus bifidus.

Caps. unilocularis, tetrasperma.

Hydrophyllaeae. Bot. reg. 740.

NAVARRETIA R. et P. (1799.)

Flores capitati, involucrati.

Cal. campanulatus, quinquefidus, laciniis tridentatis.

Cor. infundibuliformis, tubo pentagono, limbo quinquelobo.

Stam. tubo inserta.

Stylus stigmate bifido.

Caps. bilocularis, tetrasperma.

Hydrophyllaeae.

665. *PORANA* N. L. *Burm.* (1768.)*Cal.* quinquepartitus.*Cor.* campanulata; limbo quinquepartito aequali,*Stam.* longitudine corollae: filamenta tubo adnata: thecae subrotundae.*Stylus* bifidus, stigmatibus capitatis.*Caps.* bilocularis, tetrasperma, calyce tumido cincta.*Convolvuleae.* *Jacqu. ecl.* 2. t. 165.666. *BREWERIA* R. Br. (1810.)*Cal.* quinquepartitus, non mutatus.*Cor.* infundibuliformis, plicata.*Stylus* bipartitus, stigmatibus capitatis.*Caps.* bilocularis, tetrasperma, valvata.*Convolvuleae.*2. *Fructus drupacei, baccati.*667. *MARIPIA* Aubl. (1775.)*Cal.* quinquepartitus, imbricatus, lobis obtusis, 5-ventibus.*Cor.* campanulata, plicata, extus tomentosa.*Stam.* intra tubum: filamenta breviter: antherae sagittatae.*Stylus* filiformis, exsertus: stigmate peltato, decem-*Drupa* tetrasperma.*Convolvuleae.* *Aubl. guian.* t. 91.668. *ARGYREIA* Lour. (1790.)*Cal.* pentasepalus, connivens.*Cor.* cylindrico-infundibuliformis; limbo plicato, lobis quinquelobo.*Stam.* inclusa, inaequalia: filamenta tubo adnata, incrassata: antherae erectae sagittatae.*Stylus* filiformis, stigmate didymo-capitato.*Bacca* quadrilocularis, tetrasperma.*Convolvuleae.* *Bot. reg.* 661. [*Lettsomia Roxb.*]669. *TECTONIA* L. fil. (1781.)*Cal.* campanulatus, quinquefidus, persistens.*Cor.* infundibuliformis, tubo brevi, limbo quinquefidus*Stam.* filamenta brevissima, fauci inserta: antherae bogae exsertae.*Stylus* stigmate bilobo.*Drupa* hirsuta, spongiosa-suberosa; putamine quadriloculari, tetraspermo.*Vitaceae.* *Roxb. corom.* 1. t. 6. *Gärtn.* t. 57.670. *TOURNEFORTIA* L. (1737.)*Cal.* quinquepartitus.*Cor.* hypocrateri-infundibuliformis, tubo basi globoso, limbo quinquelobo.

Stam. filamenta breviora, subulata, tubo inserta: antherae acuminatae, conniventes, inclusae.

Stylus clavatus, stigmate integro.

Drupa quadrilocularis, di-tetrasperma.

Asperifoliae. Jacq. ic. 1. t. 31. Gärtn. t. 76. Bot. reg. 464. [Pittosia Plum. Messerschmidia L.]

BOURRERIA Jacq. (Bourreria P. Brown 1756.)

Cal. subbilabiatus, quinquedentatus.

Cor. infundibuliformis, limbo quinquepartito.

Stam. filamenta subulata, tubo inserta: antherae oblongo-lanceolatae, exsertae.

Stylus simplex, aut bifidus, stigmatibus capitatis.

Drupa tetrapyrena, ossiculis bilocularibus, mono-disper-

ma. Embryo erectus.

Asperifoliae. P. Brown jam. t. 15. f. 2. Jacq. amer. t. 173. Gärtn. t. 212. [Freylinia Coll. Morelosia Lessert.]

PABBLEA Mart. (1826.)

Cal. quinquepartitus.

Cor. infundibuliformis, limbo decedentato, dentibus alternis minoribus, introrsum conniventibus.

Flos fasciculis pilorum quinque.

Antherae conniventes, apice barbatae, latae, membrana saetae.

Drupa tetrapyrena.

Asperifoliae. Mart. nov. gen. 2. t. 164.

EBRETIA P. Brown. (1756.)

Cal. quinquedentatus, vel irregulariter incisus.

Cor. campanulato-infundibuliformis, limbo quinquepartito.

Stam. filamenta subulata, tubo inserta: antherae basi bilobae, biloculares, exsertae.

Stylus bifidus, stigmatibus acutis vel obtusis.

Drupa dipyrena, ossiculis dispermis, vel tetrapyrena, tetrasperma. Embryo inversus.

Asperifoliae. P. Brown jam. t. 16. f. 1. Humboldt nov. gen. t. 208. 209. Bot. reg. 1097.

RHABDIA Mart. (1826.)

Cal. quinquepartitus.

Cor. campanulata, tubo brevi, limbo quinquefido.

Stam. imo tubo inserta. Antherae erectae, oblongae, biloculares, inclusae.

Stylus filiformis. Stigma bilobum.

Drupa tetrapyrena, pyrenis monospermis.

Asperifoliae. Mart. nov. gen. 2. t. 195.

ELLISIA L. (1764.)

Cal. quinquepartitus.

Cor. subcampanulata, calyce minor, limbo quinquelobo, lobis emarginatis.

Stam. tubo breviora, antheris subrotundis.

Stylus brevis, stigmate bifido.

Receptaculum ante maturitatem succulentum, deinde succum, hivalve, tetraspermum.

Asperifoliae. Lam. ill. t. 97. Gärtn. t. 184.

676. *MENALS* Löff. (1757.)

Cal. trisepalus, persistens.

Cor. tubulosa, tubo cylindrico calycem superante, quinquelobo,

Stam. filamenta brevissima, tubo inserta: antherae sublatae, ad faucem corollae.

Stylus filiformis, stigmatibus duobus oblongis.

Bacca globosa, quadrilocularis, tetrasperma.

Asperifoliae?

677. *ROTULA* Lour. (1790.)

Cal. cyathiformis, quinquepartitus, persistens.

Cor. rotata, quinquepartita.

Stam. filamenta subulata, corollae basi inserta, ipsa breviora. Antherae sagittatae.

Stylus linearis planiusculus, stigmate emarginato.

Bacca unilocularis, tetrasperma.

678. *COUMA* Aubl. (1775.)

Cal. quinquefidus.

Cor. infundibuliformis, laciniis angustis primum, teris

Stam. brevissima. Antherae sagittatae.

Discus hypogynus.

Stigma bifidum.

Drupa tri-pentapyrena.

Contortae. Aubl. guian. t. 392.

e. *Fructus* penta-decasermi.

1. *Membranacei.*

679. *CORIS* Diosc. Matth. (1560.)

Cal. campanulatus, quinquedentatus, sub dentibus quoque setis patentibus instructus.

Cor. tubulosa, limbo erecto quinquefido, laciniis ob bifidis.

Stam. laciniis corollae opposita: filamenta basi corollae inserta: antherae erectae, oblongae, corollam aequantes.

Stylus simplex, stigmate capitato.

Caps. quinquevalvis. Placenta centralis quinque- pentasperma.

Primuleae. Bot. reg. 536. Gärtn. t. 183.

1. **ANDROSACE** Matth. (1560.)*Cal.* pentagonus, quinqueidentatus.*Cor.* hypocrateriformis, fauce glandulosa, limbo quinquelobo.*Stam.* intra tubum.*Caps.* quinquevalvis, penta- aut decasperma.*Primulaceae.* Schk. t. 33. *Reichenb. ic.* t. 248. *Gärtn.* t. 50. [*Aretia* L.]2. **DOUGLASIA** Lindl. (1827.)*Cal.* obconicus, angulatus, quinqueidentatus.*Cor.* infundibuliformis, tubo ventricoso, limbo plano quinquepartito, fauce callo lineari sub quovis sinu instructa.*Caps.* unilocularis quinquevalvis, pentasperma. Placenta centralis libera pedicellata quinqueangulata.*Primulaceae.*2. *Drupacei vel baccati.*3. **HUBBERTIA** Cominers. Juss. (1789.)*Cal.* coriaceus, quinquepartitus.*Cor.* tubulosa subcampanulata, limbo erecto quinquelobo.*Stam.* longe exserta, filamentis corollae hinc insertis: antherae subsagittatae, biloculares.*Stylus* longissimus, incurvus, stigmate obtuso.*Bacca* substipitata, bilocularis, loculis di-trispermis.*Convolvuleae.* Lam. ill. 103. [*Endræchium* Juss. Thouinia Sm. ic. ined. t. 7.]83. **STYPHELIA** Sm. (1793.)*Cal.* quinquefidus, bracteatus.*Cor.* tubulosa: ad basin tubi fasciculi quinque villorum: limbus revolutus, barbatus.*Stam.* exserta.*Squamae* hypogynae quinque distinctae, raro connatae.*Drupa* subexsucca, putamine osseo pentaspermo.*Epacrideae.* Andr. bot. repos. 72. Bot. reg. 24. [*Pentachondra* R. Br.]84. **ASTROLOMA** R. Br. (1810.)*Cal.* quinquefidus, bracteatus.*Cor.* tubuloso-ventricosa: ad basin tubi fasciculi quini villorum: limbus brevis, patens, barbatus.*Stam.* inclusa.*Discus* perigynus, cyathiformis, integer.*Drupa* subexsucca, putamine osseo, pentasperma.*Epacrideae.* Bot. mag. 1439. [*Ventenatia* Cav. ic. 4. t. 380.]35. **STENANTHERA** R. Br. (1810.)*Cal.* quinquepartitus, multibracteatus.*Cor.* tubuloso-ampullacea, limbo brevi, quinquefido, semibarbato. Fasciculi villorum nulli in basi tubi.

697. JACQUINIA L. (1764.)

Cal. quinquepartitus, persistens.*Cor.* subcampanulata, limbo decemlobo, duplici lobis interioribus minoribus.*Stam.* basi corollae inserta. Antherae posticae.*Stylus* brevissimus, stigmate obtuso.*Bacca* globosa subpentasperma. Albumen carnosum.*Myrsineae.* Humb. nov. gen. 3. t. 246. Gärtn. t. 20

698. LEEA Royen. L. (1740.)

Flores monoeci.

Cal. campanulatus, quinquefidus.*Cor.* tubo brevi, limbo quinquefido, laciniis saccatis.*Urceolus* perigynus cylindricus quinquefidus, bifidis.*Stam.* cum laciniis urceoli alternantia, brevia. Antherae incumbentes.*Stylus* simplex, stigmate incrassato, lacero.*Bacca* penta - hexasperma.*Sapoteae.* *Melieae.* Andr. repos. 355. [Aquil.]*Cav.* diss. 7. t. 218. Gärtn. t. 218.]f. *Fructus polyspermi.*l. *Membranacei.*aa. *Uniloculares.*aa. *Capsulae.*

699. EUTOCIA R. Br. (1823.)

Cal. quinquepartitus, persistens.*Cor.* subcampanulata. Plicae, seu membranulae geminae utrinque filamentis adstantes.*Stam.* exserta. Filamenta tubo inserta. Antherae incumbentes, biloculares.*Stylus* bifidus.4 *Caps.* unilocularis, bivalvis, polysperma; valvis indiviso medio placentiferis.*Hydrophyllaeae.* Franklin. it. arctic. t. 27. Bot. reg. 11

P R I M U L E A E.

700. CORTUSA Matth. (1560.)

Cal. quinquefidus, persistens, laciniis dentatis.*Cor.* subrotata: faux tuberculis basi annulatis clau limbus quinquepartitus.*Stam.* tubo innata, filamentis brevissimis, antheris bicularibus.*Stylus* filiformis, corolla longior, stigmate simplici.

Caps. unilocularis, apice quinquedentata. Placenta columnaris centralis, libera.

Jacqu. ic. rar. 1. t. 32. Andr. repos. t. 1. Gärtn. t. 50.

ANAGALLIS Dioscor.

Cal. quinquepartitus, persistens.

Cor. rotata, quinquepartita.

Stam. basi corollae inserta: filamenta inferne hirsuta.

Antherae incumbentes, biloculares.

Stylus simplex, stigmate capitato.

Caps. globosa, unilocularis, circumscissa. Placenta globosa, libera. *Sem. angulata.*

Sturm. fl. germ. Gärtn. t. 50.

LYSIMACHIA Diosc. Trag. (1551.)

Cal. quinquepartitus, persistens.

Cor. rotata, quinquepartita.

Stam. basi corollae inserta: filamenta subulata, basi subcomata.

Caps. globosa, quinque-decemvalvis. Placenta centralis globosa.

Schk. t. 36. Gärtn. t. 50.

LUBINIA Venten. (1800.)

Cal. quinquepartitus.

Cor. subhypocrateriformis vel campanulata, quinquepartita.

Stam. exserta, filamentis tubo adnatis.

Stylus simplex, persistens.

Caps. subealvis: placenta centralis libera.

Venten. hort. Cels. t. 96. Link et Otto. ic. t. 27.

CYCLAMEN Nicand.

Cal. campanulatus, semiquinquefidus.

Cor. basi globosa, fauce prominula, limbi laciniis reflexis.

Stam. filamenta minima, tubo inserta. *Antherae conniventes*, sagittatae.

Stylus filiformis exsertus.

Caps. baccata, demum quinquefariam dehiscens. *Sem. in pulpa nidulantia.*

Sturm. fl. germ. Gärtn. t. 183.

DODECATHEON L. (1751.)

Cal. quinquepartitus, laciniis reflexis.

Cor. quinquepartita, laciniis limbi reflexis.

Stam. filamenta brevissima, latiuscula, tubo insidentia.

Antherae sagittatae, in rostrum conniventes.

Stylus staminibus longior, simplex.

Caps. apice dehiscens dentibus quinque.

Schk. t. 34. Gärtn. t. 50.

706. SOLDANELLA C. Gesn. (1550.)

Cal. quinquefidus.

Cor. subcampanulata, limbo erecto fimbriato-lacero.

Stam. filamenta subulata: antherae sagittatae.

Stylus longitudine corollae, persistens.

Caps. cylindrica, apice muldentata. Placenta centralis angulata, libera.

Sturm. fl. germ. *Gärtn.* t. 183.

707. PRIMULA Matth. (1560.)

Flores subumbellati, involucrati.

Cal. tubulosus, quinquefidus vel quinquedentatus, persistens.

Cor. tubulosa, fauce vel nuda vel glandulosa, limbo quinquelobo.

Stam. inclusa: filamenta brevissima, tubo inserta: antherae oblongae, conniventes.

Stylus filiformis, stigmate capitato.

Caps. cylindrica, apice decedentato dehiscens. Placenta centralis libera.

Sturm. fl. germ. *Gärtn.* t. 50.

708. HOTTONIA Boerh. (1720.)

Cal. quinquepartitus.

Cor. hypocrateriformis, limbo quinquelobo.

Stam. intra tubum.

Stylus persistens.

Caps. globosa, evalvis. Placenta globosa centralis libera.

Engl. bot. 364. *Fl. dan.* 1867. *Gärtn.* t. 193.

709. METTERNICHIA Mikan. (1825.)

Cal. campanulatus, quinquefidus, persistens.

Cor. infundibuliformis, plicata, leviter quinquefida.

Stam. tubo inserta. Antherae subcordatae erectae, teribus dehiscentes.

Styl. filiformis. Stigma peltatum, bilobum, lobis reflexis.

Caps. bivalvis, valvis bifidis. Placenta septiformis, libera. Sem. linearia.

Convolvuleae. *Mikan.* del. fl. brasil.

710. CALIBRACHOA Lexarz. (1825.)

Cal. pentasepalus.

Cor. infundibuliformis, fauce angulata, limbo plicato quinquelobo: fovea in medio lobo inferiori.

Stam. filamenta tubi basi inserta: antherae subrotundae.

Stylus filiformis, stamina aequans.

Caps. conica, unilocularis, bivalvis. Placenta conica.

711. RAMONDIA Rich. (1805.)

Cal. quinquefidus.

Cor. rotata, quinquepartita.

Stam. approximata, antheris longitudinaliter dehiscentibus.

Caps. unilocularis, bivalvis, valvis placentiferis.

Solanaceae. [*Verbascum* *Trew* plant. sel. t. 57.]

SESSHA Ruiz. et Pav. (1798.)

Cal. tubulosus quinquedentatus.

Cor. infundibuliformis, limbo plicato.

Stam. filamenta curvata, villosa.

Stigma bilobum.

Caps. cylindrica, bivalvis, valvis bifidis. Sem. imbricata, compressa, margine membranacea.

Personatae? R. et P. fl. peruv. 2. t. 115. 116.

BACOPA Aubl. (1775.)

Cal. quinquepartitus, inaequalis, lacinia suprema maiore, duabus inferioribus deflexis.

Cor. subhypocrateriformis, limbo quinquelobo.

Stam. fauci inserta: antherae sagittatae, exsertae.

Stylus brevis.

Caps. membranacea.

Personatae. Aubl. guian. t. 49.

G E N T I A N E A E.

11. ERYTHRAEA Renealm. (1611.)

Cal. quinquepartitus, pentagonus.

Cor. infundibuliformis, limbo quinquefido.

Stam. filamenta tubo corollae adnata: antherae defloratae spirales.

Stylus brevis: stigmata duo subrotunda.

Caps. bivalvis, valvis introflexis loculos mentientibus, placentiferis.

[*Gentiana Schk.* t. 59. *Chironia Sturm.* fl. germ.]

12. SABBATIA Adans. (1763.)

Cal. quinque - (duodecim -) partitus.

Cor. urceolata, limbo quinque - (duodecim -) lobo.

Stam. filamenta tubo inserta: antherae demum spiratim tortae.

Stylus brevis: stigmata spiratim torta.

Caps. valvis inflexis placentiferis.

Salisb. parad. lond. t. 32. *Reichenb.* hort. 2. t. 118.

16. HIPPION. * (1824.)

Cal. quinquefidus cum capsula demum concreescens.

Cor. campanulata vel hypocrateriformis, limbo quinquelobo.

Antherae biloculares (non tortae), longitudinaliter dehiscentes.

Stigma capitatum.

Caps. bipartibilis, valvis inflexis placentiferis.

[*Exacum Smith* ic. ined. t. 18. *Gentiana Plum.* ic. t. 81. f. 2. *Curtia Schlechtend.* *Schubleria Mart.* nov. gen. 2. t. 186—188.]

17. CHIRONIA L. (1737.)

Cal. quinquefidus, post anthesin tumescens.

Cor. campanulata vel hypocrateriformis, limbo quinque
Antherae oblongae erectae, post anthesin spiratim tor
Stigma capitatum.

Caps. cum calyce concreta (subbaccata), valvis in
dissepimentum fingentibus.

Lam. ill. t. 108. *Gärtn.* t. 114.

718. MYRMECIA Schreb. (1789.)

Cal. tubulosus, quinqueangularis, quinquedentatus.

Cor. infundibuliformis, limbo quinquefido.

Stam. exserta. *Antherae* erectae, siccitate haud mut

Stylus filiformis, stigmate bilamellator

Caps. bivalvis, valvis in placentas intumescens.

[*Tachia Aubl. guian.* t. 29. *Mart. nov. gen.* 2. t. 189.

719. LITA Schreb. (1789.)

Cal. tubulosus, quinquefidus, coloratus, persistens.

Cor. tubo longissimo, limbo hypocrateriformi quinque

Antherae didymae in fauce corollae sessiles.

Stylus filiformis, stigmate capitato.

Caps. valvis inflexis placentiferis.

[*Voyria Aubl. guian.* t. 83.]

720. MENYANTHES Dalechamp. (1586.)

Cal. quinquepartitus, persistens.

Cor. infundibuliformis, quinquefida, intus dense villosa.

Stam. filamenta brevia: antherae basi bilobae.

Stylus filiformis, stigmate bilobo.

Caps. bivalvis, valvis axi placentiferis.

Sturm fl. germ.

721. VILLARSIA Venten. (1803.)

Cal. quinquepartitus.

Cor. subrotata, quinquepartita, basi hirsuta.

Stam. filamenta tubo corollae affixa: antherae corda

Stylus brevis, stigmate bilobo.

Glandulae circa ovarium quinque, cum staminibus
alternantes.

Caps. subcarnosa compressa (evalvis in aquatilibus, b
valvis in terrestribus). Sem. compressa, marginato-alat

Sturm. fl. germ. Venten. select. t. 9. [*Menyanthes*]

*Limnanthemum S. G. Gmel.; Waldschmidia Wigg
Schweikharta C. C. Gmel.]*

bb. *Fructus folliculares gemini.*

C O N T O R T A E.

A P O C Y N E A E.

α α α. *Semina plerumque ad umbilicum comosa.*

α α α. *Faux corollae nuda.*

722. ECHITES P. Brown. (1756.)

Cal. quinquepartitus.

Cor. infundibuli - vel hypocrateriformis.

Stam. inclusa aut exserta: antherae apice cassae in conum circa stigma conniventes, cui adhaerent.

Squamae quinque perigynae.

Stylus simplex, stigmate crassiusculo cum antherarum cono cohaerente.

Jacqu. ic. rar. 1. t. 53. amer. t. 21 — 27. Humb. nov. gen.

3. t. 243. [Parsonsia R. Br. Forsteronia G. F. W. Meyer.]

BRAUMONTIA Wallich. (1825.)

Cal. pentasepalus, patens.

Cor. campanulato-ventricosa maxima, limbo quinquefido.

Stam. apici tubi inserta: antherae sagittatae, circa stigma cohaerentes.

Stylus filiformis arcuatus. Stigma fusiforme.

Squamae perigynae nullae.

Ovarium singulare.

Indl. in Bot. reg. t. 911.

VALLARIS N. L. Burm. (1768.)

Cal. quinquepartitus.

Cor. hypocrateriformis, limbo quinquefido.

Stam. fauci inserta, exserta. Antherae sagittatae, tergo vesiculam (N. L. Burm.) seu gibberem (R. Br.) aut corpus peltatum (Roth.) gerentes, conum circa stigma, cum quo cohaerent formantes.

Squamae quinque hypogynae, ciliatae.

Stylus simplex, stigmate clavato.

[Pergularia L. Houttuyn syst. 3. t. 20. f. 1. Emericia Röm. et Schult., Peltanthera Roth.]

ICHTHOCARPUS R. Br. (1809.)

Cal. quinquefidus.

Cor. hypocrateriformis, laciniis falcatis.

Stam. inclusa. Antherae sagittatae, a stigmate liberae, postice cassae.

Filamenta quinque sterilia perigyna cum staminibus alternantia.

Stylus filiformis, stigmate ovato, acuminato.

[Apocynum Burm. zeyl. t. 12. f. 1.]

HOLARRHENA R. Br. (1809.)

Cal. quinquepartitus, laciniis subulatis.

Cor. hypocrateriformis, laciniis aequilateribus.

Stam. imo tubo inserta: filamenta vix ulla: antherae a stigmate liberae, lanceolatae, longitudinaliter polliniferae.

Filamenta nulla perigyna.

Stylus brevissimus, stigmate cylindrico.

[Carissa Vahl. symb. 3. t. 59.]

APOCYNUM Matth. (1560.)

Cal. quinquefidus.

Cor. campanulata quinquefida: denticulis tubi q
acutis inclusis.

Stam. receptaculo inserta: filamenta brevissima.
rae sagittatae, postice cassae, conum circa
formantes.

Glandulae quinque perigynae cum staminibus
nantes.

Stylus subnullus, stigmate capitato.

Schk. t. 53.

728. *CRYPTOLEPIS* R. Br. (1809.)

Cal. quinquefidus.

Cor. infundibuliformis: squamæ obtusae tubi cum
limbi alternantes.

Stam. inclusa, imo tubo inserta. *Antherae* sagitt

Squamæ quinque perigynae cum staminibus alteri

Stylus nullus. Stigma capitato - conicum.

729. *THENARDIA* Kunth. (1818.)

Cal. quinquepartitus.

Cor. hypocrateriformis, quinquefida.

Stam. exserta. Filamenta membranacea, plani
medio connata, apice incurva. *Antherae* sagittat
conum circa stigma cohaerentes.

Stylus filiformis, stigmate pentagono.

Squamæ perigynae quinque carnosae.

Humb. nov. gen. 3. t. 240. [Echites? *Labill.* nov.
t. 31.]

730. *ALSTONIA* R. Br. (1809.)

Cal. quinquefidus.

Cor. hypocrateriformis, limbo quinquepartito, l
obliquis.

Stam. tubo inclusa: filamenta brevissima: anthe
berae, longitudinaliter polliniferae.

Squamæ nullae perigynae.

Stylus simplex, stigmate dilatato, subbifido.

Folliculi longissimi. Semina utrinque ciliis elo
comosa.

Labill. nov. Caled. t. 32.

bbb. *Faux corollae coronata.*

731. *PRESTONIA* R. Br. (1809.)

Cal. quinquepartitus.

Cor. hypocrateriformis. Corona faucis duplex, e
annularis, indivisa: interior squamis quinque, c
ciliis limbi alternantibus.

Stam. filamenta brevissima, tubo inserta. *Anther*
gittatae, in conum conniventes.

Squamae quinque perigynae.

Stylus filiformis, stigmate incrassato.

Humb. nov. gen. 3. t. 242.

732. **BALFOURIA** R. Br. (1809.)

Cal. basi squamulis decem, corollae adstantibus.

Cor. infundibuliformis, limbi laciniis rectis aequilateribus.

Faux simpliciter coronata, tubo crenulato.

Stam. fauci inserta. Antherae sagittatae, mucronatae, medio stigmati cohaerentes.

Stylus filiformis, stigmate angulato.

733. **NERIUM** Diosc.

Cal. quinquepartitus.

Cor. infundibuli-hypocrateriformis. Faux coronata squamis laceris.

Stam. tubo adnata. Antherae sagittatae, cohaerentes, apice filamentosae.

Stylus subclavatus, stigmate truncato.

Schk. t. 52.

734. **STROPHANTHUS** Cand. (1802.)

Cal. quinquepartitus.

Cor. infundibuliformis, limbi laciniis filiformibus caudatis.

Faux coronata squamis bifidis.

Stam. intra tubum: antherae sagittatae, apice aristatae.

Squamae quinque perigynae.

Stylus filiformis, stigmate subcylindrico.

Desfont. anal. mus. 1. t. 27. *Bot. reg.* t. 469. *Reichenb. hort.* 2. t. 159.

735. **WRIGHTIA** R. Br. (1809.)

Cal. pentasepalus: basi squamis quinque cinctus.

Cor. hypocrateriformis. Faux coronata squamis multifidis.

Stam. exserta: filamenta fauci adnata. Antherae sagittatae cum medio stigmate cohaerentes.

Stylus filiformis, apice dilatato.

Bot. reg. 933.

$\beta\beta\beta$. *Semina nuda.*

736. **VINCA** Plin.

Cal. quinquepartitus.

Cor. hypocrateriformis, fauce quinqueplicata, limbi laciniis oblique truncatis.

Stam. filamenta apice dilatata. Antherae membranaceae biloculares.

Stylus apice dilatatus. Stigma peltatum.

Glandulae duae inter germina.

Schk. t. 51.

737. *TABERNAEMONTANA* Plum. (1703.)*Cal.* quinquefidus.*Cor.* hypocrateriformis. Faux fornicibus clausa.*Stam.* inclusa. Antherae sagittatae, conniventes.*Stylus* filiformis, stigmatе capitato, peltato.*Sem.* pulpaе immersa.*Vahl.* eclog. 3. t. 6. Bot. reg. 338. 716. [*Vahea* Lam.
t. 169. *Urceola* Roxb.]738. *CAMERARIA* Plum. (1703.)*Cal.* quinquedentatus.*Cor.* infundibuli - hypocrateriformis, laciniis limbi obli-*Stam.* tubo inserta. Antherae acuminatae, conniventes,
apice in filum productae.*Styli* vix ulli, stigmatibus obsoletis.*Folliculi* reflexi, basi utrinque lobati. *Sem.* membrana
peculiari inserta.*Lam.* ill. t. 173. *Aubl.* guian. t. 102.739. *ASPIDOSPERMA* Mart. (1826.)*Cal.* quinquepartitus.*Cor.* tubo ventricoso, laciniis limbi patentibus obliquis.*Stam.* tubo inclusa. Filamenta brevissima. Antherae
ovatae, biloculares, longitudinaliter dehiscentes.*Stylus* cylindricus. Stigma clavatum.*Folliculi* obovati, lignosi. *Sem.* compressa, alata,
niculo umbilicali centrali.*Mart.* nov. gen. 1. t. 34—36.740. *AMSONIA* Walt. (1788.)*Cal.* quinquefidus.*Cor.* tubo cylindrico, fauce barbata, limbi laciniis quinque-
que subobliquis.*Stam.* inclusa. Antherae ovatae.*Stylus* simplex, stigma peltatum.*Sem.* teretia, apice truncata.

Bot. reg. 151. \

741. *ALLAMANDA* L. (1771.)*Cal.* quinquepartitus.*Cor.* infundibuliformis ventricosa, limbo quinquefido.*Stam.* faucibus inserta, antheris subsessilibus sagittatis.*Annulus* perigynus.*Stylus* filiformis. Stigma orbiculo duplici peltatum.*Caps.* coriacea compressa, undique aculeis longis echinata,
bivalvis. *Sem.* margine alata.*Gärtn.* t. 61. [*Orelia* *Aubl.* guian. t. 106.]742. *ANABATA* Willd. Schlechtend. (1819.)*Cal.* campanulatus, brevissime quinquedentatus.*Cor.* campanulata, limbo quinquefido.

Antherae in tubo sessiles.

Stylus exsertus.

Caps. tetragona, stylo persistente coronata, valvis placentiferis.

Contortae? [Sulzeria Röm. et Schult., Rauwolfia oppositifolia N. Entdeck. 3. p. 33.]

$\beta\beta$. *Capsulae* aut *folliculi* biloculares.

MITREOLA Rich. (1823.)

Cal. quinquepartitus, persistens.

Cor. urceolata.

Stam. inclusa.

Stylus stigmatе indiviso.

Caps. bifurcata, bilocularis: placentis parietalibus internis.

— — Rich. in act. soc. nat. paris. 1. t. 3.

LYONSIA R. Br. (1809.)

Cal. quinquefidus.

Cor. infundibuliformis, limbo quinquepartito aequali.

Stam. exserta: filamenta medio tubo inserta. *Antherae* sagittatae, postice cassae, cum medio stigmatе coherentes.

Stylus filiformis, apice dilatato.

Quamae hypogynae connatae.

Caps. cylindrica, bilocularis, valvis folliculiformibus, dissepimento parallelo libero, utrinque seminifero.

Contortae.

ROMANZOVIA Chamiss. (1820.)

Cal. quinquepartitus.

Cor. hypocrateriformis, fauce nuda.

Stam. inclusa, filamentis tubo adnatis, antheris erectis.

Stylus simplex, persistens. Stigma capitatum.

Caps. compressa, bilocularis, dissepimento valvis contrario utrinque placentifero.

Nees hor. phys. berol. t. 14.

SIBTHORPIA L. (1764.)

Cal. quinquepartitus.

Cor. subrotata, quinqueloba (interdum minus aut plus divisa.)

Stam. tubo inserta, quatuor, quinque, septem, octo.

Stylus simplex, stigmatе capitato.

Caps. compressa, bilocularis: dissepimentum valvis contrarium, utrinque placentiferum.

Personatae. Engl. bot. 649. Gärtn. t. 55. Humb. nov. gen. 2. t. 176. 177. [Disandra L. Bot. mag. 268. Gärtn. t. 185.]

747. VERBASCUM Plin.

Cal. quinquefidus.

Cor. rotata, limbo inaequaliter lobato.

Stam. basi corollae inserta. Filamenta barbata. rae plerumque diffformes.

Stylus simplex, stigmatate incrassato.

Caps. valvis inflexis dissepimentum formantibus, incertas incrassatis.

Personatae. Schk. t. 42. Gärtn. t. 55.

748. HYOSCYAMUS Dioscor.

Cal. campanulatus, limbo quinquefido aut quidentato.

Cor. infundibuliformis, limbo patente oblique quinquefidus (interdum lateraliter fisso, in unica specie subtruncato.)

Stamina basi corollae inserta.

Stylus simplex, stigmatate capitato.

Caps. operculo circumscisso, bilocularis: dissepimento utrinque placentas incrassatas gerente.

Solaneae. Sturm fl. germ. Bot. reg. 180.

749. NICOTIANA Dalechamp. (1586.)

Cal. tubulosus, quinquefidus.

Cor. infundibuliformis, limbo plicato.

Caps. apice quadridentata, placentis discretis.

Solaneae. Schk. t. 44. Gärtn. t. 35.

750. PETUNIA Juss. (1803.)

Cal. quinquepartitus, lacinijs subspathulatis.

Cor. infundibuliformis.

Stam. inaequalia.

Stylus simplex, stigmatate bilobo.

Caps. apice bifida, bilocularis: dissepimento utrinque placentas gerente.

Solaneae. Juss. in ann. mus. 2. t. 47.

751. LEHMANNIA * (1817.)

Cal. disepalus.

Cor. ringens, tubo gracili incurvo.

Stam. exserta.

Solaneae. [Nicotiana tomentosa R. et P. fl. p. t. 429.]

752. LAMARCKIA Pers. (1804.)

Cal. tubulosus, pentagonus, quinquefidus.

Cor. hypocrateriformis, quinquepartita, limbo subobtus.

Caps. teres polysperma.

Solaneae. [Markea Rich.]

753. FABIANA Ruiz. et Pav. (1799.)

Cal. quinquefidus.

Cor. infundibuliformis, limbo plicato revoluto.

Filamenta duo longiora.

Stigma emarginatum.

Convolvulaceae. R. et P. fl. peruv. 2. t. 122. *Hilar.* in
mém. mus. 12. t. 9.

734. *NIEREMBERGIA* Ruiz. et Pav. (1799.)

Cal. basi tubulosus, apice quinquepartitus, patens.

Cor. tubo gracili elongato, limbo quinquelobo.

Stam. filamentis basi connatis.

Stylus filiformis, stigmate infundibuliformi.

Caps. bilocularis, bivalvis, valvis bifidis, dissepimento
valvis parallelo, demum libero: placentae dissepimento
arcte nexae.

Convolvulaceae? Solanaceae? Humb. nov. gen. 3. t. 198.

735. *CONVOLVULUS* Plin.

Cal. quinquepartitus, nudus vel bracteatus.

Cor. campanulata vel infundibuliformis, limbo plicato.

Stylus simplex, stigmate capitato vel fuso.

Caps. 1-2-3-4-locularis, oligo-polysperma.

Sturm fl. germ. *Jacq.* ic. rar. 1. t. 36. ecl. t. 44. *Gärtn.*

134. [*Ipomoea* L., quam iam in horto Cliffortiano p. 66.
hand rite distingui fatetur. *Calystegia* R. Br., *Morenoa*
Lexarz.]

736. *RETZIA* Thunb. (1776.)

Cal. quinquepartitus, laciniis inaequalibus lanceolatis.

Cor. tubulosa, villosa; limbo erecto quinqueudentato.

Stam. corolla breviora. *Antherae* sagittatae.

Stylus filiformis, stigmate bifido.

Caps. bilocularis, bivalvis, bisulca.

Convolvulaceae. Act. soc. lundin. 1. t. 1. *Lam.* ill.
t. 103.

737. *GENIOSTOMA* Forst. (1775.)

Cal. turbinatus quinquefidus.

Cor. urceolato-infundibuliformis, intus villosa; limbo
quinquepartito.

Stam. filamenta faucis corollae inserta: *Antherae* oblon-
gae, exsertae.

Stylus filiformis, stigmate sulcato, didymo.

Caps. bilocularis, placentis filiformibus ad dissepimentum
nexus. Sem. angulosa.

Contortis affinis. *Forst.* char. t. 12. [*Anasser* Juss.]

738. *LOGANIA* R. Br. (1810.)

Cal. quinquepartitus.

Cor. campanulata, fauce villosiuscula, limbo quinque-
partito.

Stam. intra tubum, filamentis brevissimis, antheris ob-
longis versatilibus, bilocularibus.

Stylus brevis. Stigma clavatum, bilobum.

Caps. bilocularis: placentae incrassatae utrinque ad sepimentum. Sem. peltata, reticulata.

Gentianeae, Rubiaceis et Contortis affinis.

Entdeck. 1. t. 2. [Euosma Andr. 420. Exacum nov. holl. t. 51.]

759. *SPIGELIA* L. (1747.)

Cal. quinquepartitus.

Cor. infundibuliformis, fauce imberbi, limbo quinque-

Stam. intra tubum, inclusa vel exserta.

Stylus basi persistens, cum stigmate articulatus.

Caps. didyma, bilocularis, loculis bivalvibus. Sem. angulata. Placentae columnares, demum liberae.

Gentianeis et Rubiaceis affinis. P. Brown jam.

f. 3. Humb. nov. gen. 3. t. 226. Mart. nov.

t. 192 — 194. Gärtn. t. 198.

760. *LISIANTHUS* P. Brown. (1756.)

Cal. subcampanulatus, quinquefidus, lacinüs diaphanis et incumbentibus.

Cor. infundibuliformis, fauce imberbi, limbo quinque-

Stam. basi tubi inserta. Antherae sagittatae, bilocales.

Stylus elongatus. Stigma bilamellatum vel bicrura.

Caps. bilocularis, dissepimento e valvis introflexis. Sem. angulata.

Gentianeae. P. Brown. jam. t. 9. Smith. ic. t. 29.

Bot. reg. 883. Gärtn. t. 183. [Eustoma Sm.]

parad. lond. t. 34. Helia Mart. nov. gen. 2. t. 491.

Irlbachia ib. t. 179.]

761. *SPHENOCLEA* Gärtn. (1788.)

Cal. urceolatus quinquefidus, basi bracteatus.

Cor. minor, calyci adnata; quinquefida.

Stam. tubo inserta.

Stylus nullus. Stigma capitatum.

Caps. circumscissa, polysperma.

Aizoideae. Gärtn. t. 24. [Pongatium Juss. Gärtn.

Retz. Rapinia Lour.]

762. *VELLEIA* Sm. (1798.)

Cal. tri-pentasepalus, inaequalis.

Cor. tubulosa, basi ovario accreta, limbo bilabiato: h. sup. bi-infer. tripartito, lacinüs alatis undulatis.

Stam. perigyna. Glandula inter filamenta duo anterior.

Antherae distinctae, oblongae, erectae.

Stylus simplex, stigmate obliquo, bilobo.

Caps. basi bilocularis, valvis bipartitis. Sem. imbricata compressa.

Goodeniaceae. Juss. in ann. mus. 18. t. 1. Bot. reg. 5.

971. Hook. exot. fl. 1. t. 24.

*γγ. Capsulae triloculares.*763. **DIAPENSIA** L. (1735.)*Cal.* quinquepartitus, basi tribracteatus.*Cor.* hypocrateriformis, limbo quinquelobo.*Stam.* corollae inserta. Antherae biloculares.*Stylus* cylindricus, stigmate obtuso.*Caps.* trilocularis: placentae demum liberae. Sem. membrana obvoluta.*Ericaceae.* Fl. dan. t. 47. *Wahlenb.* fl. lapp. t. 9. Svensk bot. t. 517. [*Pyxidantha Michaux.* fl. bor. amer. t. 17. antheris differt. transverse dehiscentibus appendiculatis. *Lepuropetalon Elliott.*]764. **POLEMONIUM** Diosc.*Cal.* urceolatus, quinquefidus.*Cor.* subcampanulata, limbo quinquelobo.*Stam.* filamentis basi dilatatis, antheris incumben-^{ti}b^{us}.*Stylus* filiformis, stigmate trifido.*Caps.* trilocularis, trivalvis, apice trifariam dehiscens.

Dissepimenta valvis contraria.

Polemoniacae. Schk. t. 38. *Gärtn.* t. 62.765. **CANTUA** Juss. (1789.)*Cal.* urceolatus, tri-quinquefidus.*Cor.* infundibuliformis, limbo quinquelobo.*Stam.* exserta.*Stigma* trifidum.*Caps.* trilocularis, trivalvis, calyce cincta, Sem. apice alata.*Polemoniacae.* Cav. ic. 4. t. 363. 364. [*Periphragmos Ruiz. et Pav.* fl. per. 2. t. 131—133. *Vestia W.*]766. **GILIA** Ruiz. et Pav. (1798.)*Cal.* campanulatus, quinquefidus.*Cor.* infundibuli-hypocrateriformis, limbo patente quinquefido.*Stam.* fauci inserta. Antherae biloculares.*Stylus* simplex, stigmate trifido.*Caps.* trilocularis, trivalvis. Sem. duplici serie, angulata, nuda.*Polemoniacae.* Bot. maj. 2883. Bot. reg. 1170. [*Ipomopsis Mx., Ipomeria Nuttall.*]767. **HOITZIA** Juss. (1789.)*Cal.* quinquepartitus, basi bracteis plerumque serratis suffultus.*Cor.* infundibuliformis, limbo quinquelobo.*Stam.* exserta.*Caps.* trilocularis, trivalvis, oligo-polysperma.*Polemoniacae.* Cav. ic. 4. t. 365. 366. [*Löselia L. Lam.* ill. t. 527. *Gärtn.* t. 62.]

768. *COLLOMIA* Nuttall. (1818.)

Flores aggregati, involucrati.

Cal. campanulatus, quinquefidus, glandulosus.

Cor. tubulosa, limbo subhypocrateriformi, quinquefidus.

Stam. tubo inserta, subinclusa. Antherae ovales, culares.

Styl. filiformis. Stigmata tria.

Caps. trilocularis, trivalvis, loculis mono- & dispermiis.

liber, triqueter. Sem. mucosae obvoluta, qui tracheaeformibus scatet.

Polemoniaceae. Bot. reg. 1166. 1174. Bot. mag. 2893.

769. *COBAEA* Cav. (1791.)

Cal. campanulatus, quinquefidus, quinqueangulatus, gulis alatis.

Cor. campanulata, quinqueloba, lobis trilobis.

Stam. declinata, filamentis demum spiratim tortis.

Stigma tri-quinquefidum.

Caps. obovata, tri-quinquelocularis, totidem valvis hiscens. Sem. imbricata, marginata.

Bigoniaceae. Cav. ic. 1. t. 16. 17. 500. Andr. repos.

♂♂. *Capsula quadrilocularis.*

770. *DATURA* Garc. ab Orto. (1563.)

Cal. tubulosus, quinquefidus, basi peltata persistens.

Cor. infundibuliformis, plicata, limbo dentato.

Stylus filiformis, stigmatibus bilamellato.

Caps. echinata vel inermis, semi-quadrilocularis, quadrivalvis, placentis discretis.

Solanaceae. Schk. t. 43. Gärtn. t. 132.

ss. *Capsulae quinqueloculares.*

771. *AZALEA* L. (1737.)

Cal. quinquedentatus.

Cor. infundibuliformis, limbo quinquefido.

Stam. receptaculo inserta, inaequalia.

Caps. tri-quinquelocularis, valvis dissepimentum formantibus. Sem. nuda.

Ericaceae. Andr. repos. 16. Engl. bot. 865. Gärtn. t.

772. *EPACRIS* Forst. (1775.)

Cal. coloratus, bracteis imbricatis suffultus.

Cor. tubulosa, limbo imberbi.

Stam. tubo inserta. Antherae horizontales, biloculares.

Squamae quinque perigynae.

Stylus simplex, stigmatibus capitato.

Caps. placentis columnae centrali adnatis, demum liberis.

Epacridaceae. Forst. char. 10. Labill. nov. holl. t. 5.

58. Smith exot. bot. 1. t. 39. 40.

DRACOPHYLLUM Labill. (1798.)*Cal.* quinquepartitus.*Cor.* tubulosa, limbo imberbi, quinquefido.*Stam.* hypogyna, vel basi corollae inserta: antherae dorso adnatae.*Styl.* simplex. Stigma trilobum.*Squamae* quinque hypogynae.*Caps.* placentis ab apice columnae centralis pendulis.*Epacrideae*, Labill. it. 2. t. 40. *Reichenb.* hort. 2. t. 108.**LYSINEMA** R. Br. (1810.)*Cal.* coloratus, multibracteatus.*Cor.* hypocrateriformis, limbi laciniis imberbibus, dextrorsum flexis.*Stam.* hypogyna. Antherae horizontales.*Squamae* quinque perigynae.*Caps.* placentis columnae centrali adnatis.*Epacrideae*. [Epacris Cav. ic. 4. t. 346.]**ANDERSONIA** R. Br. (1810.)*Cal.* coloratus, bracteatus.*Cor.* limbo basi barbato.*Stamina* hypogyna. Antherae infra medium affixae.*Squamae* quinque perigynae subconnatae.*Caps.* placentis columnae centrali annexis.*Epacrideae*.**SPRENGELIA** Smith. (1794.)*Cal.* coloratus.*Cor.* subrotata, quinquepartita, imberbis.*Stam.* hypogynae. Antherae connatae barbatae, vel liberae imberbes.*Squamae* nullae.*Caps.* placentis columnae centrali annexis.*Epacrideae*, Smith in act. holm. 1794. t. 8. *Andr.* repoa. t. 2.**77. PONCELETIA** R. Br. (1810.)*Cal.* foliaceus.*Cor.* breviter campanulata, quinquefida, imberbis.*Stam.* hypogyna. Antherae horizontales, septo marginato.*Squamae* nullae.*Caps.* placentis columnae centrali annexis.*Epacrideae*.**78. COSMELIA** R. Br. (1810.)*Cal.* foliaceus, imbricatus.*Cor.* tubulosa.*Stam.* corollae adnata. Antherae apicibus ciliatis filamentorum adnatae.

Squamae quinque hypogynae.

Caps. placentis columnae centrali annexis.

Epacrideae.

779. *CYSTANTHE* R. Br. (1810.)

Cal. foliaceus.

Cor. clausa calyptraeformis, transverse dehiscens, basi truncata persistente.

Stam. hypogyna, persistentia.

Squamae nullae.

Caps. placentis ab apice columnae centralis pendulis solutis.

Epacrideae.

780. *RICHEA* R. Br. (1810.)

Cal. membranaceus, ebracteatus.

Cor. clausa calyptraeformis, transverse dehiscens, basi truncata persistente.

Stam. hypogyna, persistentia.

Squamae quinque hypogynae.

Caps. placentis ab apice columnae centralis pendulis solutis.

Epacrideae.

781. *PRIONOTES* R. Br. (1810.)

Cal. ebracteatus.

Cor. tubulosa, limbo imberbi.

Stam. hypogyna, filamentis tubo semiadhaerentibus.

Squamae quinque hypogynae.

Epacrideae. [*Epacris* Labill. nov. holl. 1. t. 59.]

782. *CAMPYLUS* Lour. (1790.)

Cal. tubulosus, quinquepartitus, tuberculosus, laciniis subulatis elongatis.

Cor. tubulosa, bilabiata: labio sup. subulato, infer. ovato.

Stam. inaequalia, imo tubo inserta. Antherae ovatae, erectae.

Stylus simplex, stigmate quinquelobo.

Caps. subrotunda, quinquelocularis.

2. *Fructus baccati.*

aa. uniloculares.

783. *STRYCHNOS* Wachendorf. (1747.)

Cal. quinquepartitus.

Cor. tubulosa, limbo quinquepartito.

Stam. fauci inserta.

Stigma capitatum.

Bacca crustacea. Sem. peltata, placentae centralis affixa.

Contortae. Roxb. coron. 1. t. 4. Lam. ill. t. 119. Lesschenault in ann. mus. 16. t. 23. Gärtn. t. 179.

84. *THEOPHRASTA* Royen. L. (1740.)*Cal.* quinquedentatus.*Cor.* campanulata, limbo quinquelobo.*Stam.* corolla breviora, in urceolo perigyno.*Bacca* placentae centrali libera.*Contortae.* *Plum.* ic. t. 126. *Jacq.* hort. schönbrenn.1. t. 116. [*Clavija R.* et *P.* prodr. t. 30. *Oucinus Lour.*]85. *LEONIA* R. et P. (1798.)*Cal.* quinquelobus.*Cor.* rotato-campanulata.*Filamenta* connata in tubum pentagonum. *Antherae* didymae, transverse dehiscentes, quadrivalves.*Stylus* stigmati simplici.*Bacca* corticosa: placentatio parietalis.*Contortae.* *R.* et *P.* fl. peruv. 2. t. 222. *Mart.* nov.

gen. 2. t. 168. 200. f. 4.

86. *DORAENA* Thunb. (1784.)*Cal.* quinquepartitus, laciniis ovatis concavis.*Cor.* rotata, quinqueloba.*Stam.* *Antherae* subsessiles, tetragonae.*Squamulae* quinque perigynae persistentes.*Stylus* simplex.*Bacca* sicca, globosa, striata, squamis quinque coronata.*Placentae* centralis libera, hispida. *Sem.* angulata.— — *Gärtn.* t. 184.87. *HANCORNIA* Gomez. (1812.)*Cal.* minimus quinquepartitus.*Cor.* limbus quinquefidus patentissimus.*Stam.* intra tubum.*Stigma* bifidum.*Bacca* polysperma.

S O L A N E A E.

88. *CESTRUM* L. (1737.)*Cal.* urceolatus, quinquedentatus.*Cor.* infundibuliformis, limbo plicato quinquefido.*Stam.* tubo inserta, saepe denticulata.*Caps.* unilocularis, polysperma. *Placenta* centralis libera.*Herit.* stirp. 1. t. 35. 36. *Gärtn.* t. 77.89. *DARTUS* Lour. (1790.)*Cal.* quinquefidus.*Cor.* tubo subgloboso, limbo quinquefido patente.*Stam.* tubo inserta. *Antherae* incumbentes.*Stylus* brevissimus. *Stigma* quinquelobum.*Bacca* excoriata, diaphana, unilocularis, polysperma.90. *MANDRAGONA* Theophr.*Cal.* quinquefidus.

Cor. campanulata quinquefida.

Stam. filamenta basi dilatata, conniventia.

Bacca carnosio-spongiosa. Sem. reniformia periph.

Tournef. instit. t. 12. *Gärtn.* t. 131.

β β. biloculares.

791. ATROPA L. (1737.)

Cal. quinquepartitus.

Cor. infundibuliformis, campanulata, quandoque sutata, limbo quinque- (interdum decem-) lobo.

Stam. subexserta.

Bacca bilocularis, placentis liberis.

Schk. t. 45. *Gärtn.* t. 131. [*Saracha R.* et *P.* differt a subrotata, bacca abortu subuniloculari.]

792. ULLOA Pers. (1805.)

Cal. ovatus, inflatus, quinquepartitus, coloratus.

Cor. tubulosa, fauce gibbosa, limbo patente minimo.

Stam. tubo inserta.

Bacca bilocularis, calyce inflato tecta.

[*Juanulloa R.* et *P.* fl. per. 2. t. 185.]

793. PHYSALIS L. (1737.)

Cal. quinquefidus.

Cor. subrotata.

Stam. basi corollae inserta. Antherae longitudine dehiscentes.

Bacca bilocularis, calyce inflato colorato tecta.

Schk. t. 45. *Jacq. ecl.* 2. 137. *Gärtn.* t. 131.

794. CAPSICUM Fuchs. (1542.)

Cal. quinquefidus.

Cor. rotata, limbo plicato, quinquefido.

Stam. filamenta brevissima. Antherae conniventes, gitudinaliter dehiscentes.

Bacca polymorpha, bilocularis, placenta basilari. superne dissepimento adnata.

Schk. t. 47. *Gärtn.* t. 132.

795. SOLANUM Plin.

Cal. quinque- (decem-) partitus.

Cor. subrotata, plicata (quinque- interdum decem)

Antherae conniventes, apice poro gemino dehisce (interdum inaequales.)

Bacca bi- (tri- quadri-) locularis, placentis septo natis. Sem. glabra.

Schk. t. 46. *Dunal hist. solan.* t. 2. 4—26. *Gärtn.* t. [Bassovia *Aubl.*, *Nycterium Vent.*, *Androcera Nu Aquartia Jacqu.*]

796. LYCOPERSICON Anguill. (1560.)

Cal. quinque- (sex-) partitus.

- Cor.* rotata, quinque - (sex -) loba.
Antherae conicae, membrana apice elongata connatae, longitudinaliter dehiscentes.
Bacca bi - (tri -) locularis. Sem. villosa.
Dunal hist. solan. t. 3. 16. Bot. mag. 2814.
79. **WITHERINGIA** Herit. (1788.)
Cal. urceolatus, quinque - (quadri -) fidus.
Cor. campanulato - rotata, tubo gibboso.
Antherae longitudinaliter dehiscentes.
Bacca bilocularis, calyce persistente suffultae.
Herit. sert. angl. 1. Bot. mag. 2768.
80. **DIERBACHIA** * (1825.)
Cal. urceolatus, quinquedentatus.
Cor. infundibuliformis, limbo plicato.
Stam. inclusa, tubo inserta. Filamenta tripartita, cuspidem media antherifera. *Antherae* longitudinaliter dehiscentes.
Stylus filiformis, stigmate capitato.
Bacca bilocularis, placentis septo adnatis.
[Dunalia Kunth. Humb. nov. gen. 3. t. 194.]
799. **JABOROSA** Juss. (1789.)
Cal. quinquefidus.
Cor. tubuloso - campanulata, limbo quinquefido.
Stam. fauci inserta, filamentis subclavatis.
Stylus simplex, stigmate bilobo.
Bacca bilocularis, placentis crassis dissepimento adnatis.
Link et Otto ic. t. 48. Lam. ill. t. 114.
800. **ANISODUS** Link. (1824.)
Cal. campanulatus, inflatus, angulatus, quinquedentatus, persistens.
Cor. campanulata quinqueloba.
Stam. tecta, basi corollae inserta, basi cohaerentia, corolla minora.
Bacca exsucca, bilocularis, operculo quadrivalvi. Placenta scrobiculata.
[Nicandra Link. et Ott. ic. t. 35.]
801. **LYCIUM** Diosc.
Cal. campanulatus, quinquedentatus, saepe lateraliter fissus.
Cor. infundibuliformis.
Stam. tubo corollae inserta: filamentis basi villosis.
Bacca bilocularis, dissepimento in placentam incrassato.
Schk. t. 46. *Gärtn.* t. 132.
802. **CARISSA** L. (1767.)
Cal. quinquepartitus.
Cor. infundibuliformis.
Stam. inclusa.

Bacca bilocularis. Sem. compressa.

Contortae. Roxb. corom. 1. t. 77.

803. *WILLUGHBEIA* Scop. (1778.)

Cal. quinquepartitus.

Cor. hypocrateriformis, limbo quinquepartito, undula obliquo.

Stam. tubo inserta, filamentis brevissimis, antheris gittatis.

Stylus simplex. Stigma ovatum, orbiculo plano inside bicuspidatum.

Bacca bilocularis. Sem. in pulpa nidulántia vel dissemento affixa.

Contortae. [Ambelania et Pacouria *Aubl.* guian. t. 4
105. *Fagraea Thunb.*]

γγ. Fructus baccati quadri-quinque-multilocularis

804. *CAMAX* Schreb. (1789.)

Cal. quinquepartitus.

Cor. rotata, limbo quinquelobo.

Stam. corollae inter lacinias inserta.

Stylus capillaris. Stigmata quatuor.

Bacca ovalis, villosa, quadrilocularis. Sem. in pulpa nidulántia.

— — [Ropourea *Aubl.* guian. t. 78.

805. *NICANDRA* Adans. (1763.)

Cal. quinquefidus, angulatus.

Cor. campanulata, subintegra.

Stam. basi corollae inserta, inferne dilatata, connivent
Antherae ovatae.

Bacca quadri-quinquelocularis. Placentae inter dissepimenta.

Solaneae. Gärtn. t. 131. [Alkekengi *Feuill.* obs. 1. t. 1.
Calydermos *R. et P.*]

806. *SOLANDRA* Sw. (1787.)

Cal. campanulatus, quinquefidus.

Cor. maxima, infundibuliformis, ventricosa, limbo undula quinquefido.

Stam. ascendentia. Antherae versatiles.

Stylus inflexus. Stigma bilobum.

Bacca quadrilocularis, placentis dissepimento adnatis.

Solaneae. Swartz in act. holm. 1787. t. 11. fl. ind. o
t. 9. *Salisb.* in *Linn. transact.* 6. t. 6. *Corr.* in *al*
mus. 8. t. 4.

807. *DESFONTAINIA* Ruiz. et Pav. (1798.)

Cal. quinquepartitus.

Cor. campanulata, limbo quinquefido.

Stam. fauci inserta. Antherae biloculares, extus dehiscentes.

Stylus simplex, stigmate capitato.

Bacca quinquelocularis.

Solanaceae. R. et P. fl. peruv. 2. t. 186. Humboldt pl. sequin. 1. t. 45. [Linkia Pers.]

809. **CERIUM** Lour. (1790.)

Cal. quinquepartitus, laciniis subulatis.

Cor. campanulata, limbo quinquelobo.

Stam. tubo adnata, exserta.

Stylus subulatus, stigmate crassiusculo.

Bacca multilocularis, loculis monospermis.

g. *Fructus ignoti.*

810. **BARRERIA** Scop. (1780.)

Cal. quinquedentatus.

Cor. quinquepartita. Laciniae superne concavae scrobiculo duplici, superiori ovato bifido, inferiori trifido.

Stam. filamentis latis emarginatis. Antherae articulatae tetragonae, subquadrialatae, foveis corollae receptae.

Stylus simplex, stigmatibus tribus.

— — [Poraqueiba Aubl. guian. t. 47.]

811. **TORREYA** * (1821.)

Cal. quinquelobus, patens, coloratus.

Cor. tubulosa, limbo quinquefido, reflexo.

Stam. filamenta tubo adnata, corolla longiora, spiralia.

Antherae didymae.

Stylus simplex, stigmate acuto.

Germen nucem referens, valvis introrsum flexis.

Nyctagineae?

812. **TRATTINICKIA** Willd. (1805.)

Cal. campanulatus, tridentatus.

Cor. campanulata, tridentata.

Stam. receptaculo inserta.

Stylus subulatus.

Terebinthaceae.

813. **MINA** Lexarz. (1824.)

Cal. carnosus, quinquefidus, laciniis sub apice mucronatis.

Cor. ventricosa, quinqueangularis, limbo angustato quinquedentato.

Stam. summo tubo inserta, ascendencia. Antherae biloculares.

Glandula carnosae perigyna.

Stylus filiformis. Stigma globosum.

13. **DIPLOCALYMMA** * (1822.)

Cal. duplex, ext. bivalvis, inter. decedentatus.

Cor. infundibuliformis subplicata.

Antherae sagittatae intra tubum.

Stigma urceolatum, subbilobum.

Convolvuleae.

814. *NECTOUXIA* Kunth. (1818.)

Cal. quinquepartitus.

Cor. hypocrateriformis: faux coronata annulo promianulo integerrimo: limbus quinquefidus.

Stam. intra tubum. Antherae erectae, biloculares.

Stylus filiformis. Stigma bifidum.

Solaneae. Humb. nov. gen. 3. t. 193.

B. *Fructus inferi.*

a. *Capsulares.*

a. *Capsulae uniloculares.*

815. *SPEERMADICTION* Roxb. (1816.)

Cal. quinquefidus.

Cor. infundibuliformis, tubo gracili elongato, limbi quinquefidi laciniis ovatis.

Stam. tubo inserta.

Stylus filiformis, stigmatе quinquefido.

Caps. apice quinquefariam dehiscens, pentasperma. Sem. membrana reticulata obvoluta.

Rubiaceae. Roxb. corom. 3. t. 236. Bot. reg. 348-4235. [Hamiltonia Don.]

816. *BELONIA* Plum. (1703.)

Cal. semiinferus, quinquepartitus, persistens.

Cor. rotata, limbo quinquelobo.

Stam. basi corollae inserta. Antherae conniventes.

Stylus subulatus.

Caps. laciniis calycis coronata, polysperma.

Rubiaceae. Lam. ill. t. 149.

817. *SAMOLUS* Plin.

Cal. semiinferus, quinquefidus.

Cor. campanulata, hypocrateriformis, limbo quinquelobo, denticulis inter lobos.

Stam. tubo inserta.

Caps. apice quinquedentata, placenta centrali libera.

Primulaeae. Fl. dan. 198. Gärtn. t. 30. Sheffieldia Forst. Labill. nov. holl. t. 54.

818. *LYSIPOMA* Kunth. (1818.)

Cal. quinquefidus, laciniis inaequalibus.

Cor. tubulosa, subbilabiata: lab. sup. bi-infer. trilobo.

Antherae connatae, barbatae.

Stigma bilobum ciliatum.

Caps. unilocularis seminafera, placentatione peripherica.

Lobeliaceae. Humb. nov. gen. 3. t. 266. 267.

819. **ROELLA** L. (1737.)

Cal. turbinatus, quinquepartitus, lacinii dentatis.

Cor. infundibuliformis, limbo quinquepartito.

Squamae quinque perigynae, concavae, staminiferae, persistentes.

Stam. filamenta subulata. Antherae conniventes.

Stylus filiformis. Stigmata duo oblonga.

Caps. unilocularis. Sem. angulata, scabra, funiculis apici capsulae affixis.

Campanulaceae. Linn. hort. Cliff. t. 35. Bot. mag. 378. Gärtn. t. 31.

β. Capsulae biloculares.

820. **MACHAONIA** Bonpl. (1808.)

Cal. quinquepartitus.

Cor. infundibuliformis, fauce villosa staminifera.

Stigma bifidum.

Caps. bilocularis, disperma, basi dehiscent.

Rubiaceae. Humb. plant. aequin. 1. t. 29.

821. **CANEPHORA** Juss. (1789.)

Involucrum tri-sexflorum, campanulatum, subquinquefidum. Flores singuli squamis distincti.

Cal. proprius quinque-sexfidus.

Cor. campanulata, quinque-sexloba.

Antherae oblongae sessiles.

Stigma bifidum.

Caps. globosa, bilocularis, disperma.

Rubiaceae. Lam. ill. t. 151.

822. **RONDELETIA** Plum. (1703.)

Cal. quinquepartitus, subpentasepalus.

Cor. infundibuliformis, fauce inflata.

Stam. inclusa.

Stigma bifidum.

Caps. subglobosa, bilocularis, loculis dispermis. Sem. solida.

Rubiaceae. Plum. ic. t. 242. f. 1. Gärtn. t. 181. Humb. nov. gen. 3. t. 290. 291.

823. **CHIMARRHIS** Jacq. (1762.)

Cal. integer, persistens.

Cor. infundibuliformis, limbo quinquefido: lacinii subtus medio hirsutis.

Stam. inter lacinias corollae inserta. Antherae ovales, erectae.

Stylus filiformis, stigmate bifido.

Caps. bilocularis, bivalvis; valvis apice bifidis; locale mono - dispermis.

Rubiaceae.

824. *SIPANEA* Aubl. (1775.)

Cal. quinquepartitus, patens. (Dentes glandulosi minimi laciniis calycis interiecti in una specie).

Cor. infundibuli - hypocrateriformis, tubo gracili, limbo quinquefido.

Stamina tubo inserta. *Antherae* conniventes.

Stylus filiformis, stigmate bipartito.

Caps. globosa, angulata, bilocularis, bivalvis. *Placentae* crassae utrinque ad dissepimentum. *Sem.* numerosa, aptera.

Rubiaceae. Aubl. *Guian.* t. 56. [Virecta L. fil. *Gärtn.* t. 184.]

825. *MACROCNEUM* P. Brown. (1756.)

Cal. turbinatus, obsolete dentatus. *Bractae* saepe magnae petiolatae adstantes.

Cor. infundibuli - campaniformis, limbo quinquepartito patulo.

Stam. fauci inserta, exstantia.

Stylus simplex, stigmate bilobo.

Caps. bilocularis, polysperma. *Sem.* obsolete marginata.

Rubiaceae. Vahl. *symb.* 2. t. 30. Ruiz et Pav. *fl. per.* 2. t. 189. 190.

826. *CINCHONA* L. (1749.)

Cal. quinquedentatus.

Cor. infundibuliformis (ut plurimum extus pubescens), limbo patente quinquefido.

Stam. tubo adnata, vel inclusa, vel tubum corollae superantia. *Antherae* lineares, biloculares, longitudinaliter dehiscentes.

Stylus simplex, stigmate bifido.

Caps. bilocularis, polysperma, medio intus dehiscens.

Sem. alata, margine lacero.

Rubiaceae. Humb. *plant. aequin.* 1. t. 10. 19. 39. 47. Vahl. *in act. havn.* 1. t. 1. 2. Ruiz et Pav. *fl. per.* 2. t. 191. 195. 196. *Gärtn.* t. 33. [Hymenodictyon Wallisch.] *Cosmibuena* R. et P. *fl. peruv.* 3. t. 198. 226.

827. *PINKNEYA* Michaux. (1803.)

Cal. quinquepartitus, laciniis inaequalibus (una aut altera longiore latioreque).

Cor. tubulosa, pubescens, laciniis limbi quinquefidi revolutis.

Stam. tubo inserta, exstantia. *Antherae* versatiles.

Stylus simplex, stigmate bilobo.

Caps. bilocularis, polysperma, valvis medio semiseptiferis.
Sem. alata.

Rubiaceae. Michaux fl. bor. am. t. 13.

88. EXOSTEMA Bonpl. (1808.)

Cal. quinquedentatus, regularis.

Cor. tubulosa, tubo longissimo, limbi laciniis quinque patentibus.

Stam. summo tubo inserta, exstantia.

Stylus exsertus, stigmatе incrassato simplici.

Caps. bilocularis, polysperma. *Sem.* alata, margine integerrimo.

Rubiaceae. Humb. plant. aequin. 1. t. 38.

89. MUSSAENDA Raj. (1704.)

Cal. irregulariter quinquefidus, laciniis una alterave bracteiformibus.

Cor. infundibuliformis, limbo patente quinquepartito.

Antherae subsessiles, tubo corollae inclusae.

Stylus simplex, stigmatе bifido.

Caps. bilocularis, placentis ad dissepimenta transversis.

Sem. solida, non alata.

Rubiaceae. Bot. reg. 517. Lam. ill. t. 157. Cailloud it. Meroë, 2. t. 62. Gärtn. t. 28.

890. HYMENOPOGON Wallich. (1825.)

Cal. quinquepartitus.

Cor. longe tubulosa, hypocrateriformis, limbo quinquefido.

Stam. intra tubum.

Caps. bilocularis, quadrivalvis. *Sem.* utrinque membrana appendiculata.

Rubiaceae.

81. LEGANANTHUS Jack. (1824.)

Cal. carnosus, amplus, coloratus, inaequaliter bi-quadrilobus.

Cor. calyce minor, tubo abbreviato, limbo quinquefido.

Stylus bifidus.

Caps. bilocularis, polysperma.

Rubiaceae.

82. ALSKIS Henr. Schott. (1827.)

Cal. quinquepartitus.

Cor. cyathiformis, fauce barbata, limbo quinquelobo.

Stam. ad basin corollae usque libera, exserta.

Stigma bipartitum.

Germen biloculare, multi-ovulatum.

Rubiaceae.

83. DANAIS Commers. Juss. (1789.)

Flores subdioeci.

Cal. quinquedentatus.

Cor. infundibuliformis, intus hirta.

Stam. tubo inserta. Antherae incumbentes.

Stigma bifidum.

Caps. globosa, bilocularis, apice dehiscens, septis in centas incrassatis. Sem. alata.

Rubiaceae. Gärtn. t. 195. [*Paederia* Juss. Lam. t. 166. f. 2.]

834. *DENTELLA* Forst. (1775.)

Cal. quinquepartitus, laciniis subulatis.

Cor. infundibuliformis, laciniis limbi tridentatis.

Stam. inclusa.

Stigma bifidum.

Caps. globosa, bilocularis, polysperma.

Rubiaceae. Forst. char. gen. t. 13.

835. *PORTLANDIA* P. Brown. (1756.)

Cal. pentasepalus, persistens.

Cor. infundibuliformis, ventricosa, amplissima, quinquefido.

Stam. declinata, fundo tubi inserta, corollam aequa Antherae lineares erectae.

Stylus simplex. *Stigma* oblongum, obtusum.

Caps. quinqueangularis, bilocularis, bivalvis, apice hiscens: dissepimento valvis contrario. Sem. elliptica, scabra, horizontalia, strophiole pyramidalia.

Rubiaceae. P. Brown jam. t. 11. Gärtn. t. 34.

836. *OPHIORRHIZA* L. (1747.)

Cal. turbinatus, quinquedentatus.

Cor. tubulosa, ventricosa, intus hirsuta, limbo quinque-

Stam. tubo inserta. Antherae subsagittatae biloculares

Stylus filiformis, apice bifidus. *Stigmata* globosa.

Discus epigynus bilobus, persistens.

Caps. transverse latior, biloba, lobis divergentibus, bilocularis, polysperma. Placentae dissepimento affixae ceterum liberae. Sem. minuta, angulata.

Rubiaceae. Linn. mat. med. t. 1. Gärtn. t. 55. Rk in act. soc. nat. paris. 1. t. 2.

837. *SICKINGIA* Willd. (1801.)

Cal. quinquedentatus.

Cor. campanulata, quinquedentata.

Stam. tubo inserta, corollam aequantia. Antherae neares erectae.

Stylus subulatus, staminibus brevior.

Caps. subrotunda, lignosa, bilocularis, bivalvis, apice alata.

Rubiaceae. [? *Platycarpum* Bonpl. in Humb. plant. noct. 2. t. 104.]

838. *LOBELIA* L. (1737.)

Cal. quinquedentatus.

Cor. tubulosa, irregularis, hinc longitudinaliter fissa: limbo bilabiato, labio sup. bi - infer. trifido.

Antherae connatae, barbatae.

Stylus simplex. Indusium ciliatum sub stigmatē.

Caps. bilocularis, dissepimento utrinque placentifero.

Lobeliaeae. Schk. t. 269. Willd. hort. berol. t. 83. 86.

Ann. mus. 18. t. 1. f. 1.

ISOTOMA Lindl. (1826.)

Cal. quinquepartitus.

Cor. tubulosa, limbo hypocrateriformi, laciniis aequalibus.

Antherae connatae, imberbes, duabus inferioribus aristatis.

Stylus simplex. Stigma capitatum.

Caps. bilocularis, dissepimento utrinque placentifero.

Lobeliaeae. Lindl. in Bot. reg. 964.

LECHENAULTIA R. Br. (1810.)

Cal. quinquepartitus.

Cor. tubulosa, hinc longitudinaliter fissa, bilabiata: lab. sup. bi - inf. trilobo.

Antherae sub anthesi cohaerentes.

Stigma obsoletum, in fundo indusii bilabiati.

Caps. prismatica, bilocularis, quadrivalvis, valvis oppositis, medio septiferis.

Goodeniaeae. Bot. reg. 916.

GOODENIA Sm. (1794.)

Cal. quinquepartitus.

Cor. tubulosa, uni - bilabiata: limbi quinquepartiti laciniis margine diaphanis subalatis.

Antherae distinctae.

Stigmatis indusium simplex subciliatum.

Caps. bilocularis, dissepimento parallelo. Sem. compressa.

Goodeniaeae. Andr. repos. 68. Cav. ic. 6. t. 306. 307.

508. Labill. nov. holl. t. 75. [Calogyne R. Br. style trifido differt.]

MOQUINIA * (1828.)

Cal. semisuperus, subinteger, basi involucratus.

Cor. tubulosa, hinc longitudinaliter fissa: Lacinae quinque lineares longissimae, spiratim revolutae, altera alteri saepe adhaerente.

Antherae lineares elongatae.

Stylus longe exsertus. Stigma capitatum.

Goodeniaeae?

FORGESIA Commers. Juss. (1789.)

Cal. turbinatus, quinquefidus.

Cor. quinquepartita, intus furfuracea.

Stam. *Antherae* oblongae filamentis adnatae.

Stigma bilobum.

Caps. semiinfera, bilocularis, polysperma, apice bivalvi
et stylo persistente coronata.

— — [Defforgia Lam. ill. t. 125.]

844. *PSILOBIUM* Jack. (1826.)

Cal. amplus, quinquepartitus.

Cor. tubus brevis, limbo quinquefido.

Stam. basi corollae inserta.

Stigma clavatum, decemalatum, exsertum.

Fructus siliquaeformis, bilocularis, polyspermus.

— —

845. *ARGOSTEMMA* Wallich. (1825.)

Cal. quadri-quinquedentatus.

Cor. rotata, quadri-quinquepartita.

Antherae conniventes, exsertae.

Fructus bilocularis, polyspermus.

— —

γ. Capsulae tri-quinqueloculares.

846. *PHYTEUMA* Matth. (1560.)

Cal. quinquefidus.

Cor. quinquepartita, laciniis elongatis apice cohaerentibus.

Filamenta basi dilatata.

Stylus simplex. *Stigma* bi-trifidum.

Caps. trilocularis, placenta centrali.

Campanuleae. Schk. t. 39. Gärtn. t. 30.

847. *TRACHELIUM* Pon. (1617.)

Cal. quinquedentatus.

Cor. tubo gracili, longissimo, limbo quinquefido.

Filamenta basi aequalia.

Stylus exsertus. *Stigma* subtrilobum.

Caps. trilocularis, basi poris dehiscens.

Campanuleae. Schk. t. 40. Bot. reg. 72. Gärtn. t. 31.

848. *CAMPANULA* Dodon. (1557.)

Cal. quinquefidus.

Cor. campanulata vel subrotata, limbo quinquefido.

Filamenta basi dilatata.

Stigma tri-quinquefidum.

Caps. tri-quinquelocularis, poris lateralibus vel apice
valvato dehiscens.

Campanuleae. Schk. t. 39. Gärtn. t. 31.

1. *Medium.* Calycis sinus reflexi, basi producti. *Defforgia*
sel. e cor. *Tournef.* t. 30. 31.

2. *Adenophora* Fisch., *Codonopsis* Wallich. *Glossocodon*
Don. Nectarium cylindricum pistillum inferne adnatum
iens. *Stylus* elongatus. *Reichenb. hort.* 1. t. 2. 3.
23. 32. 45.

3. *Campanopsis* R. Br., *Wahlenbergia* Schrad. *Cervicula*
Delil. (fl. aeg. t. 5. abortu triandra). *Caps.* valvis apice
dehiscens. *Reichenb. ic.* 5. t. 480. *hort.* 2. t. 165.

4. *Erinus*. Cor. inaequalis. Caps. apice dehiscens. *Allion.*
fl. pedem. t. 7. f. 1.

5. *Primatocarpus Herit.* Cor. subrotata. Caps. prismatica. Ann. mus. 11. t. 18. Bot. reg. 56. Engl. bot. 375.

DIERVILLA Tournef. (1706.)

Cal. quinquefidus, bracteatus.

Cor. infundibuliformis, limbo quinquefido subirregulari.

Stam. exserta.

Stigma capitatum.

Caps. quadrilocularis, polysperma.

Caprifolia. Linn. hort. Cliff. t. 7.

b. *Fructus* baccati, drupacei.

a. *mono-dispermi.*

RHYTIDEA Cand. (1807.)

Cal. quinquepartitus.

Cor. infundibuliformis, limbo quinquepartito.

Stam. fauci inserta.

Stigma bisulcum.

Bacca monosperma. Semen globosum.

Rubiaceae.

51. ZUCCARINIA * (1827.)

Cal. unilateralis trifidus.

Cor. infundibuliformis, limbo quinquefido.

Antherae in tubo sessiles.

Stigma bilobum.

Bacca monosperma?

— — [*Jackia Wallich.* in *Roxb. fl. ind.* 2. p. 323.]

52. AIDIA Lour. (1790.)

Cal. tubulosus, quinquedentatus.

Cor. hypocrateriformis: fauce lanosa: limbo quinquepartito.

Antherae lineares sessiles, incisuris corollae insertae.

Stylus stigmate oblongo.

Bacca monosperma.

— —

53. DASUS Lour. (1790.)

Cal. campanulatus, quinquelobus.

Cor. campanulata, pilosissima, limbo quinquefido.

Stam. basi corollae inserta. *Antherae* biloculares.

Stylus stigmate quinquefido.

Bacca monosperma.

— —

54. HELICANTHERA Lour. (1790.)

Cal. truncatus, coloratus, squama carnosae suffultus.

Cor. hypocrateriformis, quinquefida.

Stam. filamenta filiformia, fauci corollae insidentia.

therae lineares, spiratim revolutae.

Urceolus perigynus quinquefidus.

Stylus stigmatē crassiusculo.

Bacca monosperma.

855. *DIASPASIS* R. Br. (1810.)

Cal. quinquefidus.

Cor. hypocrateriformis, subregularis; tubo fisso, pube cohaerente.

Stam. intra tubum. Antherae liberae, imberbes.

Stigma indusio semibivalvi.

Drupa exsucca monosperma.

Goodeniaceae.

856. *DAMPIERA* R. Br. (1810.)

Cal. abbreviatus vel obsoletus.

Cor. bilabiata, extus barbata, pilosa: tubo hinc fissi, laciniis labii superioris margine interiori auriculatis.

Stam. persistentia. Antherae imberbes connatae.

*Stigmati*s indusium ore nudo.

Drupa monosperma.

Ann. mus. 18. t. 2. f. 1. 2.

857. *SCHÖPFIA* Schreb. (1789.)

Cal. duplex, exterior inferus subbifidus, interior superius integer.

Cor. campanulata, basi quinqueangularis, decemlimbo, limbo quinquefido.

Antherae didymae fauci insertae.

Stylus basi glandula semiglobosa cinctus. *Stigma* subtrilobum.

Drupa monosperma.

Caprifolia. [*Codonium Vahl. act. soc. hist. nat. havn.* 1. t. 6.]

858. *PSYCHOTRIA* L. (1759.)

Cal. urceolatus, quinquefidus (interdum integer.)

Cor. infundibuliformis, fauce barbata: limbo quinquefido, revoluto.

Stam. fauci inserta, exstantia.

Stylus stigmatē bifido.

Drupa dipyrrena, sulcata, calyce coronata.

Rubiaceae. *Humb. nov. gen.* 3. t. 282. 283. *Labill. nov. Caled.* t. 47. *Lam. ill.* t. 161. *Gärtn.* t. 26. [*Psychotrophum P. Brown jam.* t. 17. *Ronobea Aubl. guian.* t. 59. *Simira ib.* t. 65. ? *Antherura Lour.* *Geophila Don.*]

859. *PALICOURBA* Aubl. (1775.)

Cal. urceolatus, quinquefidus.

Cor. tubulosa, basi ventricosa, hinc gibba: infra medium interne barbata, limbo quinquefido reflexo.

Stam. exserta.

Stylus stigmatē bifido.

Drupa dipyrena, sulcata, calyce coronata.

Rubiaceae. Aubl. guian. t. 66. Humb. nov. gen. 3. t. 285. [*Stephanium* Schreb.]

CEPHAËLIS Sw. (1788.)

Flores capitato - congesti, involucrati.

Cal. minimus quinquedentatus, bracteis suffultus.

Cor. subinfundibuliformis, fauce barbata, limbo quinquefido.

Stam. inclusa.

Stylus stigmatē bipartito.

Drupa dipyrena sulcata.

Rubiaceae. Sw. fl. ind. occ. 1. t. 10. Lam. ill. t. 152.

[*Callicocca* Schreb. Brot. in Linn. transact. 6. t. 11. *Tapogomea* Aubl. guian. t. 60. *Carapichea* ib. t. 64.]

RUDGEA Salisb. (1807.)

Cal. quinquefidus.

Cor. tubus longissimus, gracilis, laciniis limbi refractis.

Stam. fauci inserta.

Stigma bilamellatum.

Bacca disperma.

Rubiaceae. Salisb. in Linn. transact. 8. t. 18. 19.

COFFEA L. (1737.)

Cal. minimus, quinquedentatus.

Cor. infundibuliformis, fauce nuda.

Stam. exserta.

Stigma bifidum.

Bacca disperma. Sem. testa chartaceo - membranacea involuta.

Rubiaceae. Ellis historical account of Coffee. Lond. 1774. 4. Lam. ill. t. 160. Gärtn. t. 25.

CHIOCCOA P. Brown. (1756.)

Cal. urceolatus, quinquedentatus.

Cor. infundibuliformis, fauce barbata.

Stam. inclusa.

Stigma indivisum.

Bacca drupacea didyma disperma. Embryo inversus.

Rubiaceae. Andr. repos. 284. Gärtn. t. 26. [*Phallaria* Schumacher.]

PAEDERIA L. (1767.)

Cal. quinquefidus.

Cor. infundibuliformis, intus hirsuta.

Stam. intra tubum.

Stylus bipartitus.

Bacca disperma. Sem. globosa. Embryo erectus, ledonibus magnis foliaceis.

Rubiaceae. Lam. ill. t. 166. f. 1. Kämpfer ic posth. t. 9. Gärtn. t. 195. [Lygodysodea Ruiz & fl. peruv. 2. t. 188. ? Landolphia Pal. Beauv. fl. 1. t. 34.]

865. *SERISSA* Commers. Juss. (1789.)

Cal. quinquefidus.

Cor. infundibuliformis, laciniis limbi trifidis.

Antherae sessiles.

Stylus bifidus.

Bacca disperma.

Rubiaceae. [Buchozia Herit. monogr. Lycium Japo Thunb. fl. jap. t. 17. Bot. mag. 361. Dysoda Low]

866. *CANTHIUM* Lam. (1783.)

Cal. quinque - quadrifidus.

Cor. tubo abbreviato, limbo quinque - quadrifido.

Stam. fauci corollae inserta, (4, 5.)

Stigma capitatum.

Bacca dipyrena, pyrenis osseis. Embryo inversus curvus.

Rubiaceae. Cav. ic. 5. t. 436. Roxb. corom. 1. Labill. nov. holl. t. 94. [Benzonia Schumacher. p. 113.] (Canthium dubium Lindl. Bot. reg. 1026. num ob stigma bifidum.)

β. *Fructus baccati, drupacei, tri - penta - hexasp*

867. *PLOCAMA* Ait. (1789.)

Cal. urceolatus, quinquedentatus.

Cor. campanulata, limbo quinquefido, laciniis linearit

Antherae subsessiles, fauci insertae.

Stylus simplex, stigmatibus obtusis.

Bacca trilocularis, trisperma. Sem. oblonga, hinc cava. Embryo erectus.

Rubiaceae. Gärtn. t. 196. [Bartlingia Reichenb. 1. t. 11.]

868. *COLLADONIA* * (1825.)

Cal. quinquefidus.

Cor. hypocrateriformis, calyce maior.

Stam. intra tubum.

Bacca trilocularis, trisperma.

Rubiaceae.

869. *TRIOSTEUM* Mill. (1737.)

Cal. quinquepartitus.

Cor. tubulosa calycem aequans.

Stam. inclusa.

Stylus filiformis. Stigma capitatum.

Bacca tripyrena.

Caprifolia. Schk. t. 41. Gärtn. t. 26. [Triosteospermum Dill. eltham. t. 293.]

LEONICERA L. (1737.)

Cal. quinquedentatus.

Cor. tubulosa, basi gibba, limbo subirregulari, bilabiato, quinquefido.

Stam. exserta.

Bacca uni - trilocularis, subtrisperma.

Caprifolia. Schk. t. 40. [Caprifolium Tourn. Juss. Gärtn. t. 27. floribus capitato - verticillatis. *Xylosteum* Tourn. pedunculis bifloris, baccis vel coadunatis vel distinctis.]

SYMPHORIA Pers. Pursh. (1814.)

Cal. quadri - quinquedentatus.

Cor. subcampanulata, abbreviata, quinquefida.

Stam. brevia.

Stigma globosum.

Bacca quadrilocularis, loculis uno aut altero abortientibus.

Caprifolia. Bot. mag. 2211. [Symphoricarpos Dill. eltham. t. 278.]

MORINDA Vaill. (1722.)

Flores arcte capitati.

Cal. quinquedentatus.

Cor. infundibuliformis.

Stam. inclusa.

Stigma bifidum.

Drupa angulata, tetrapyrena.

Rubiaceae. Lam. ill. t. 153. Gärtn. t. 29. Labill. nov. Caled. t. 49.

S. SCAEVOLA L. (1767.)

Cal. quinquefidus, persistens.

Cor. tubulosa, hinc fissa. Limbus secundus quinquepartitus, laciniis alatis conformibus.

Stam. receptaculo inserta. Antherae exsertae liberae.

Stigmatis indusium ciliatum.

Drupa bi - quadrilocularis, subtetrasperma.

Goodenias Ann. mus. 18. t. 1. f. 4. Labill. nov. holl. t. 78 - 80. Cav. ic. 6. t. 509. 510. [Goodenia Andr. repos. 22. Smith nov. holl. t. 5.]

GUETTARDA Osbek. (1757.)

Flores submonoeci.

Cal. campanulatus subintegerrimus.

Cor. tubo longissimo, limbo hypocrateriformi.

Stam. numero incerto inclusa.

Stigma capitatum.

Drupa putamine quadri-sexloculari, loculis spermis.

Rubiaceae. Lam. ill. t. 154. Vent. sel. t. 1. eclog. 1. t. 6. Humb. nov. gen. 3. t. 292. Gärtn. [Halesia P. Brown jam. t. 20. f. 1. Laugieria Jacq. symb. 3. t. 57. Matthiola L.]

875. *NONATELIA* Aubl. (1775.)

Cal. quinquedentatus, persistens.

Cor. infundibuliformis, limbo quinquefido.

Stam. tubo inserta, exstantia.

Stylus bifidus. Stigmata obtusa.

Bacca sulcata, quinquelocularis, pentasperma.

Rubiaceae. Aubl. guian. t. 70—74. [Oribasia Retiniphyllum Bonpl. in Humb. plant. aequin. 1. differt involucro calycis triphylo, stigmate si clavato.]

876. *CUVIERIA* Cand. (1807.)

Cal. quinquepartitus.

Cor. quinquefida, persistens, laciniis acutissimis spinescentibus.

Stigma umbraculiforme, subpentagonum.

Bacca pentasperma.

Rubiaceae. Ann. mus. 9. t. 15.

877. *ANCYLANTHUS* Desf. (1818.)

Cal. quinquefidus, persistens.

Cor. tubulosa, incurva, limbo subbilabiato quinque laciniis duabus superioribus longioribus.

Antherae in tubo sessiles.

Stylus stigmate quinquelobo.

Ovarium quinqueloculare.

Rubiaceae. Mém. mus. 4. t. 2.

878. *VANGUERIA* Juss. (1789.)

Cal. minimus quinquedentatus.

Cor. parva, campanulato-globosa, intus hirta, quinquefido.

Stam. antheris oblongis.

Bacca pomiformis, umbilicata, quinquelocularis, pentasperma.

Rubiaceae. Lam. ill. t. 159. [Vavanga Rohr i soc. nat. havn. 2. t. 7.]

† 879. *LICHTENSTEINIA* Wendl. (1866.)

Cal. duplex semisuperus monophyllus tri-quinqueden

Cor. quinquefida.

Filamenta extra corollam connata.

Nectarium epigynum.

Bacca pentasperma.

— Wendl. collect. 1. t. 39.

7. *Fructus baccati, drupacei, polyspermi.*

NAUCLEA L. (1764.)

Flores arcte capitati, receptaculo globoso.

Cal. quinquedentatus, persistens.

Cor. infundibuliformis, elongata, limbo quinquefido.

Antherae subsessiles, infra limbi sinus insertae.

Stylus longissimus exsertus. *Stigma* capitatum.

Fructus drupacei congesti, dicocci, coccis polyspermis, seminibus membranaceo-marginatis.

Rubiaceae. Roxb. corom. 1. t. 53. 54. Lam. ill. t. 153.
Hunter in Linn. transact. 9. t. 22. Bot. reg. 895. [*Adina*
Baliab. parad. lond. t. 115. *Uncaria* Schreb. Willd. in
Usteri Mag. 2. t. 3. *Cephalanthus* L. fl. seyl. Lam.
Oucouparia Aubl. guian. t. 68. *Cephalina* Thouning apud
Schumacher. guin. p. 105.]

WEBBERA Schreb. (1789.)

Cal. quinquefidus.

Cor. infundibuliformis, tubo elongato, ad faucem ampliato, laciniis recurvis.

Antherae subsessiles, fauci insertae, exstantes.

Stylus in stigma clavatum decangulare transiens.

Bacca bilocularis, loculis tetra-pentaspermis.

Rubiaceae. Bot. reg. 119. Gärtn. t. 192.

82. BRITHALIS Patr. Brown. (1756.)

Cal. urceolatus, quinquedentatus.

Cor. subinfundibuliformis, limbo patente quinquepartito.

Stam. exstantia.

Stylus stigmate acuto.

Bacca decemlocularis, loculis monospermis.

Rubiaceae. P. Brown jam. t. 17. f. 3. Gärtn. t. 26.

83. HAMELIA Jacqu. (1762.)

Cal. minimus quinquedentatus.

Cor. tubo elongato quinqueangulari, limbo quinquepartito.

Stam. tubo corollae inserta.

Stylus filiformis. *Stigma* obtusum.

Bacca quinquelocularis, polysperma. Placentae membranaceae.

Rubiaceae. Smith. exot. bot. 1. t. 24. Gärtn. t. 191. 196.

84. SCHWENKFELDIA Schreb. (1789.)

Cal. quinquepartitus, subpentasepalus.

Cor. tubulosa, pilosa, limbo quinquefido.

Antherae tubo inclusae.

Stigma quinquefidum.

Bacca quinquelocularis polysperma.

Rubiaceae. Swartz. fl. ind. occ. t. 10. [Sabicea Aubl.
 guian. t. 75. 76. Urophyllum Wallich.]

885. BURCHELLIA R. Br. (1820.)

Flores capitati, involucriati.

Cal. quinquepartitus.

Cor. tubuloso-clavata.

Antherae subsessiles, inclusae.

Stigma clavatum.

Bacca bilocularis polysperma.

Rubiaceae. Bot. reg. 466. 891. [*Lonicera bubalina*]

886. STYLOCORYNA Cav. (1798.)

Cal. urceolatus, quinquedentatus.

Cor. infundibuliformis, limbo patente, quinquefido.

Antherae subsessiles, lineares, inter lacinias limbi.

Stylus clavatus, stigmate simplici.

Bacca bilocularis, polysperma. Sem. in pulpa nidula

Rubiaceae. Cav. ic. 4. t. 368. Labill. nov. Caled.
(baccis quadrilocularibus.)

887. GENIPA Plum. (1703.)

Cal. truncatus, integer.

Cor. hypocrateriformis.

Antherae subexsertae.

Stigma clavatum.

Bacca subquadrilocularis. Sem. in pulpa nidulantia.

Rubiaceae. Plum. ic. t. 136. Gärtn. t. 190. [Duroi]

888. GARDENIA Ellis. (1759.)

Cal. quinquepartitus aut quinquedentatus.

Cor. hypocrateriformis, tubo cylindrico, laciniiis obliquis.

Antherae sessiles.

Stigma bilobum.

Bacca subbilocularis, seminibus bi-quadriseialibus.

Rubiaceae. Roxb. corom. 2. t. 134 — 136. Bot.
735. 975. Gärtn. t. 193. [*Randia L.* *Oxyceras L.*
Foscarenia Vand.? *Thunbergia Montin.* *Rothme*
Thunb.]

889. HIPPODIS Ruiz et Pav. (1799.)

Cal. spathaceus, hinc fissus, integerrimus, coloratus.

Cor. infundibuliformis, tubo curvato, limbo quinquel-

Stam. filamentis medio tubo insertis, recurvis, basi

sis. *Antherae* ovaes, biloculares.

Urceolus perigynus, quinquecrenatus.

Stylus filiformis. *Stigma* bilobum.

Bacca bilocularis, polysperma.

Rubiaceae. R. et P. fl. peruv. 2. t. 201.

890. UCRIANA Willd. (1798.)

Cal. campanulatus, quinquedentatus.

Cor. tubo longissimo, limbi laciniiis lanceolatis patenti

Stam. fauci inserta.

Stigma clavatum.

Bacca seminibus compressis quadriseriatis.

Rubiaceae. [Tocoyena Aubl. guian. t. 50. Gärtn. t. 90.]

SOLENA Willd. (1798.)

Cal. turbinatus quinquefidus.

Cor. tubo longissimo recurvo, fauce pilis cincta, limbi laciniis reflexis.

Stam. fauci inserta, inaequalia.

Stylus setaceus, stigmate trifido.

Bacca cortice duro, bilocularis. Sem. in pulpa nidulantia. Embryo dorsalis, cotyledonibus cordatis, in albumine cartilagineo.

Rubiaceae. Rudg. guian. 1. t. 40, 41. [Posoquenja Aubl. guian. t. 51. Gärtn. t. 193, 195. Cyrtanthus Schreb.]

OXYANTHUS Cand. (1807.)

Cal. apice constrictus, acutissime quinquefidus.

Cor. infundibuliformis, tubo longissimo.

Stam. fauci inserta. Antherae exsertae, acutissimae.

Fructus bilocularis.

Rubiaceae. Bot. mag. 1992. [Gardenia tubiflora Andr. repos. 183.]

BERTIERA Aubl. (1775.)

Cal. turbinatus quinquedentatus.

Cor. breviter tubulosa, fauce villosa, limbo quinquefido.

Antherae subsessiles, tubo insertae.

Stylus filiformis, basi glandula cinctus. Stigma bilobatum.

Bacca bilocularis: placentae fungosae utrinque ad dissepimentum.

Rubiaceae. Aubl. guian. t. 69. Lam. ill. t. 165. Gärtn. t. 192.

LEYCESTERIA Wallich. (1824.)

Cal. inaequaliter quinquepartitus.

Cor. infundibuliformis, supra basin gibba, limbo quinquelobo.

Stam. exserta.

Stigma capitatum.

Bacca quinquelocularis, polysperma.

Caprifoliae.

SELLIERA Cav. (1799.)

Cal. quinquepartitus.

Cor. tubulosa, hinc fissa, limbi secundi quinquepartiti laciniis lanceolatis.

Stam. tubo adnata.

Stylus simplex. Stigma truncatum.

Bacca unilocularis, polysperma. Sem. alata.

Goodeniaceae. Anal. de cienc. 1. t. 5. Cav. ic. 5. t. 474.

896. *BAEBOBOTRYS* Forst. (1775.)*Cal.* quinqueidentatus, bracteis duabus suffultus.*Cor.* campanulata quinquefida.*Stylus* brevissimus.*Bacca* unilocularis, polysperma. Placenta centrali*Ericaceae*. Forst. char. t. 11. [Maesa Forsk. Juss.]

II. COROLLA PLEIOPETALA.

A. *Fructus superi.*a. *Corolla tetrapetala.*897. *STRÖMIA* Vahl. (1790.)*Cal.* patens tetrasepalus.*Petala* quatuor unguiculata (raro nulla).*Stam.* receptaculo appendiculato inserta, basi cont*Siliqua* unilocularis, bivalvis. Sem. triseriaria in nidulantia.*Capparideae*. [Cleome Burm. ind. t. 46.]b. *Corolla pentapetala.*a. *Fructus mono-dispermi.*898. *HIRTELLA* L. (1737.)*Cal.* quinquepartitus vel pentasepalus, reflexus.*Petala* subrotundo-elliptica.*Stam.* quinque (tria aut numero indefinita), calyci inserta. Filamenta longissima, ante anthesin sp involuta.*Stylus* simplex lateralis.*Bacca* corticosa, monosperma.*Chrysobalanaceae*. Aubl. guian. t. 98. Vahl, sy t. 31. Grundzüge, t. 7. f. 1—4. Humb. nov. t. 565. vol. 7. t. 658. [Moquilea Aubl. guian.*Mart.* nov. gen. 2. t. 166. Cosmibuena R. et P. fl. peruv. t. 2.899. *OPILIA* Roxb. (1799.)*Cal.* quinqueidentatus, parvus.*Pet.* oblonga.*Glandulae* perigynae, clavatae, cum staminibus nantes.*Stylus* nullus. Stigma capitatum.*Bacca* globosa monosperma.*Rhamnaceae*. Roxb. corom. 2. t. 158.900. *VILLAREZIA* R. et P. (1802.)*Cal.* minimus, foliolis imbricatis.*Petala* oblonga, patentia, subcarinata.*Antherae* subcordatae.

Stylus brevissimus, inclinatus, stigmate truncato.
caps. corticosa, bivalvis. Nux tetragona, monosperma,
 pericarpio carnoso.

Rutaceae. R. et P. fl. peruv. 3. t. 231.

PUERTEA R. et P. (1802.)

Cal. campanulatus, quinquedentatus.

Petala calyci appressa, ovata.

Stam. cum petalis alternantia. Antherae ovatae.

Stylus brevis. Stigma bitidum, acutum.

Drupa, nuce monosperma.

Rhamneae. R. et P. fl. peruv. 3. t. 227.

ICACINA Adr. Juss. (1823.)

Cal. quinquefidus, persistens.

Petala quinque, e disco hypogyno orta.

Stam. cum petalis alterna. Antherae cordatae, biloculares.

Stylus simplex incurvus, stigmate truncato.

caps. apice dehiscens monosperma.

— — Act. soc. hist. nat. paris. 1. t. 9.

TRUJANO Lexarz. (1825.)

Cal. duplex, exter. trifidus, inter. quinquefidus.

Petala ovata.

Antherae subsessiles, cum glandulis quinque alternantes.

Stylus brevissimus, stigmate trifido.

Utriculus monospermus. Sem. lenticulare, visco indutum.

SABIA Colebr. (1817.)

Cal. quinquepartitus.

Cor. pentapetala.

Stam. basi cohaerentia, petalis opposita.

Stigma simplex.

Drupa monosperma.

Colebrook in Linn. transact. 12. t. 14.

SYNDESMIS Wallich. (1826.)

Cal. spathaceus, coloratus, hinc fissus.

Petala quinque vel quatuor, basi cum filamentis juncta.

Antherae sagittatae.

Stylus lateralis, stigmate simplici.

Germen globosum monospermum.

CORYNOCARPUS Forst (1775.)

Cal. pentasepalus, subcoloratus.

Petala erecta, unguiculata, subrotunda.

Glandulae quinque, appendiculis petaliformibus, cum
 staminibus alternantes.

Stam. basi petalorum inserta. Antherae erectae oblongae.

Stylus brevis, filiformis. Stigma obtusum.

Nux clavata, monosperma.

— — Forst. char. t. 16.

ANDRIA. MONOGYNIA.

opposita iisque subiecta.

tricocca.

Rhamneae. Lam. ill. t. 129. Vahl, symb. 3. t. 58.
n. 3. t. 270. Humb. nov. gen. 7. t. 614. Gärtn. t.

919. CELASTRUS L. (1737.)

Cal. quinquelobus.

Petala unguiculata, plana.

Discus glandulosus.

Stam. antheris erectis subcordato-ovatis.

Stylus crassiusculus abbreviatus. Stigmata tria.

Caps. trilocularis, loculis monospermis. Sem. arillo
noso cincta.

Rhamneae. Schk. t. 47. Herit. stirp. t. 25. ser.
t. 10. Gärtn. t. 95.

920. MAITENUS Feuill. (1725.)

Flores polygami.

Cal. quinquefidus, persistens.

Petala cum sepalis alternantia, patentissima.

Stamina cum petalis alterna.

Discus carnosus perigynus.

Stigma sessile, bi-trilobum.

Caps. subtrivalvis. Sem. tria, quatuor, quinque,

Rhamneae. [Celastrus Ruiz. et Pav. fl. per. 3.
f. B. t. 230. f. A. Senecia Lam.]

921. POMADERIS Labill. (1804.)

Cal. quinquefidus.

Petala squamaeformia, concava.

Stam. petalis opposita iisque abscondita.

Stylus brevissimus triqueter. Stigmata tria.

Caps. tricocca, trisperma.

Rhamneae. Labill. nov. holl. t. 86. 87.

922. HOVENIA Thunb. (1781.)

Cal. quinquepartitus, patens.

Petala cum laciniiis calycis alternantia, convoluta.

Stam. petalis opposita, iisque abscondita.

Stylus simplex. Stigmata tria.

Pedunculi demum incrassati, succulenti.

Caps. tricocca.

Rhamneae. Bot. reg. 501.

923. ELAEODENDRON Jacqu. (1781.)

Cal. minimus quinquelobus.

Petala patentia, basi latiora.

Stam. cum petalis alternantia, disco perigyno inse

Stylus brevissimus.

Drupa exsucca 2-5 locularis, loculis monospermis
aut altero subinde abortiente.

Rhamneae. Jacqu. ic. rar. 1. t. 48. [Schreber
obs. 6. t. 3. Cassine Vent. sel. t. 23. Hartogia
Rubentia Juss., Portenschlagia Trattun.]

POLYCARDIA Juss. (1789.)

al. minimus quinquelobus.

etala rotundata.

am. filamentis abbreviatis, cum petalis alternantia. Antherae subrotundae.

stylus brevissimus. Stigma lobatum.

sp. tri - quinquelvalvis, valvis medio septiferis. Sem. pauca, arillo calyciformi laciniato.

Rhamneae. Lam. ill. t. 132.

EVONYMUS Theophr.

al. quadri - quinquepartitus.

etala patentia.

stylus nectarifer peltatus in fundo calycis.

am. cum petalis alternantia, glandulis e disco promissis inserta. Antherae biloculares.

sp. tri - quadri - quinquelocularis. Sem. solitaria, arillo longoso.

Rhamneae. Sturm. fl. germ. Gärtn. t. 113.

ATAVIA Dahl. Thunb. (1787.)

res capitati. Rec. lanatum, bracteis radialibus marginatis coloratis.

al. quinquepartitus.

loblonga.

m. calyci inserta.

stylus bifidus.

am. mono - tri - pentasperma.

Brunieae. Wendl. coll. t. 22. Ann. des sc. nat. 8. t. 36

INTERLIA * (1825.)

al. urceolatus, quinquelfidus, decemcostatus.

al. cum laciniis calycis alterna, ipsiusque costis imposita.

m. calyci inserta, cum petalis alterna.

sp. trivalvis trisperma.

Caryophylleae. [Sellowia Roth.]

LAYTONIA Gronov. (1743.) L. (1751.)

al. bivalvis.

etala unguiculata staminifera.

stylus trifidum.

sp. trisperma.

Portulacaeae. Schk. t. 50. Ann. mus. 7. t. 6. Persh. Amer. t. 3. C. A. Mayer in act. mosqu. 7. t. 3. [Limnia L. act. holm. 1746. t. 5.]

ORIDULA L. Murr. (1774.)

al. pentasepalus, persistens.

al. oblonga, aequalia.

m. filamentis hypogynis, apice curvatis. Antherae anaeolatae, semibilibidae, apice poro gemino dehiscentes, basi glandula didyma instructae.

Stam. petalis opposita iisque subiecta.

Stigmata tria.

Caps. sicca, tricoeca.

Rhamneae. Lam. ill. t. 129. *Vahl.* symb. 3. t. 58.
ic. 3. t. 270. *Humb.* nov. gen. 7. t. 614. *Gärtn.* t. 1

919. *CELASTRUS* L. (1737.)

Cal. quinquelobus.

Petala unguiculata, plana.

Discus glandulosus.

Stam. antheris erectis subcordato-ovatis.

Stylus crassiusculus abbreviatus. *Stigmata* tria.

Caps. trilocularis, loculis monospermis. Sem. arillo
noso cincta.

Rhamneae. Schk. t. 47. *Herit.* stirp. t. 25. ser.
t. 10. *Gärtn.* t. 95.

920. *MAITENUS* Feuill. (1725.)

Flores polygami.

Cal. quinquefidus, persistens.

Petala cum sepalis alternantia, patentissima.

Stamina cum petalis alterna.

Discus carnosus perigynus.

Stigma sessile, bi-trilobum.

Caps. subtrivalvis. Sem. tria, quatuor, quinque, ari

Rhamneae. [*Celastrus* Ruiz. et Pav. fl. per. 3. t.
f. B. t. 230. f. A. *Senecia* Lam.]

921. *POMADERIS* Labill. (1804.)

Cal. quinquefidus.

Petala squamaeformia, concava.

Stam. petalis opposita iisque abscondita.

Stylus brevissimus triqueter. *Stigmata* tria.

Caps. tricoeca, trisperma.

Rhamneae. Labill. nov. holl. t. 86. 87.

922. *HOVENIA* Thunb. (1781.)

Cal. quinquepartitus, patens.

Petala cum lacinis calycis alternantia, convoluta.

Stam. petalis opposita, iisque abscondita.

Stylus simplex. *Stigmata* tria.

Pedunculi demum incrassati, succulenti.

Caps. tricoeca.

Rhamneae. Bot. reg. 501.

923. *ELAEODENDRON* Jacqu. (1781.)

Cal. minimus quinquelobus.

Petala patentia, basi latiora.

Stam. cum petalis alternantia, disco perigyno inserti.

Stylus brevissimus.

Drupa exsucca 2-5 locularis, loculis monospermis,
aut altero subinde abortiente.

Rhamneae. Jacqu. ic. rar. 1. t. 48. [*Schrebera*
obs. 6. t. 3. *Cassine* Vent. sel. t. 23. *Hartogia* f.
Rubentia Juss., *Portenschlagia* Trattin.]

924. POLYCARDIA Juss. (1789.)*Cal.* minimus quinquelobus.*Petala* rotundata.*Stam.* filamentis abbreviatis, cum petalis alternantia. Antherae subrotundae.*Stylus* brevissimus. Stigma lobatum.*Caps.* tri-quinquevalvis, valvis medio septiferis. Sem. pauca, arillo calyciformi laciniato.*Rhamneae.* Lam. ill. t. 132.**925. EVONYMUS** Theophr.*Cal.* quadri-quinquepartitus.*Petala* patentia.*Discus* nectarifer peltatus in fundo calycis.*Stam.* cum petalis alternantia, glandulis e disco prominis inserta. Antherae biloculares.*Caps.* tri-quadri-quinquelocularis. Sem. solitaria, arillo fungoso.*Rhamneae.* Sturm. fl. germ. Gärtn. t. 113.**926. STAAVIA** Dahl. Thunb. (1787.)*Flores* capitati. Rec. lanatum, bracteis radialibus marginatis coloratis.*Cal.* quinquepartitus.*Pet.* oblonga.*Stam.* calyci inserta.*Stylus* bifidus.*Caps.* mono-tri-pentasperma.*Brunieae.* Wendl. coll. t. 22. Ann. des sc. nat. 8. t. 36**927. WINTERLIA** * (1825.)*Cal.* urceolatus, quinquefidus, decemcostatus.*Pet.* cum laciniis calycis alterna, ipsiusque costis imposita.*Stam.* calyci inserta, cum petalis alterna.*Caps.* trivalvis trisperma.*Caryophylleae.* [Sellowia Roth.]**928. CLAYTONIA** Gronov. (1743.) L. (1751.)*Cal.* bivalvis.*Petala* unguiculata staminifera.*Stigma* trifidum.*Caps.* trisperma.*Portulacaeae.* Schk. t. 50. Ann. mus. 7. t. 6. *Parsh.*
amer. t. 3. C. A. Mayer in act. mosqu. 7. t. 3. [Limnia
L. act. holm. 1746. t. 5.]**929. RORIDULA** L. Murr. (1774.)*Cal.* pentasepalus, persistens.*Pet.* oblonga, aequalia.*Stam.* filamentis hypogynis, apice curvatis. Antherae lanceolatae, semibifidae, apice poro gemino dehiscentes, basi glandula didyma instructae.

Stylus simplex, stigmatē trilobo.

Caps. trilocularis, trivalvis, loculis monospermis. Sem. scabro - scrobiculata.

Droseraceae. Lam. ill. t. 141. Gärtn. t. 62. [Ireos N. L. Burm. prodr. fl. cap. p. 6. ad calc. fl. ind.]

930. AYENIA L. (1756.)

Cal. quinquefidus.

Pet. unguiculatā, apice laminis stellam glanduliferam formantia.

Urceolus quinquedentatus staminifer. Antherae sub stella petalorum.

Stigma quinquelobum.

Caps. quinque cohaerentes, pentacoccae. Cocci monospermi.

Büttneraceae. Cavan. diss. t. 147. Gärtn. t. 79.

931. BÜTTNERIA Löfl. L. (1757.)

Cal. quinquepartitus.

Pet. unguiculata, apice cucullata aut auriculata, dorso in ligulam producta, auriculis cum tubo stamineo coalitis.

Tubus staminifer urceolatus, quinque-decudentatus. Antherae geminae in apice tubi sessiles, auriculis petalorum tectae.

Stylus brevis. Stigma quinquefidum.

Caps. pentacocca, subechinata. Cocci monospermi. Cotyledones convolutae.

Büttneraceae. Cav. diss. t. 148—150. Humb. nov. gen. 5. t. 481. a. et b. Hilar. flor. brasil. t. 27—29.

932. ALMEIDEA Hilar. (1823.)

Cal. quinquepartitus, deciduus.

Pet. unguiculata, spathulata, erecta.

Stam. hypogyna. Filamenta medio barbata. Antherae biloculares.

Stylus simplex. Stigma quinquelobum.

Cocci quinque monospermi.

Rutaceae. Hilar. flor. brasil. t. 18. Adr. Juss. in mém. mus. 12. t. 23. f. 33. [Aruba Nees in act. acad. nat. cur. 11. t. 27. 28.]

933. SPIRANTHERA Hilar. (1823.)

Cal. cupulaeformis, pentagonus, quinquefidus.

Pet. linearia, subfalcata, in unguem latiusculum continuata.

Stam. cum petalis alternantia. Filamenta glanduloso-tuberculata. Antherae lineares basi bifidae, post anthesin spiratim revolutae.

Urceolus campanulatus perigynus.

Stylus simplex. Stigma quinquelobum.

Cocci quinque, bivalves, monospermi.

Rutaceae. *Hilar.* plant. us. brasil. t. 17. *Adr. Juss.* in mém. mus. 12. t. 23. f. 32. [*Terpanthus Mart.* in act. acad. nat. cur. 11. t. 31.]

934. DIOSMA L. (1737.)

Cal. quinquepartitus, aut pentasepalus.

Pet. patentia.

Discus perigynus lobatus, vel in stamina sterilia filiformia, vel in laminas petaloideas excrecens.

Stam. filamenta filiformia. Antherae apice glanduliferae.

Cocci quinque bivalves, apice mucronati, submonospermi.

Embryo inversus.

Rutaceae. *Adr. Juss.* in mém. mus. 12. t. 18. f. 13.

1. *Adenandra* Willd., staminibus sterilibus glandula terminatis. *Adr. Juss.* l. c. t. 19. f. 16.

2. *Coleonema* Bartl., unguibus petalorum basi canaliculatis, stamina sterilia basi recipientibus. *Adr. Juss.* l. c. f. 17.

3. *Acmadenia* Bartl., staminibus sterilibus brevissimis, glandula terminatis. *Adr. Juss.* l. c. t. 18. f. 14.

4. *Barosma* Willd., staminibus sterilibus latiusculis subpetaloideis, exunguiculatis, apice obscure glanduliferis. *Adr. Juss.* l. c. t. 29. f. 18.

5. *Agathosma* Willd., staminibus sterilibus petaloideis, spathulatis, apice obscure glanduliferis. *Adr. Juss.* l. c. t. 20. f. 19. [*Hartogia* Berg. Bucco Wendl.]

6. *Macrostylis* Bartl., filamentis sterilibus nullis, petalis intus hirsutis, stylo elongato. *Adr. Juss.* l. c. t. 20. f. 20. Limitea huius divisionis neutiquam discernendi: hinc nullo modo genera esse possunt.

935. LASIOPE TALUM Smith. (1798.)

Cal. corollinus campanulatus, persistens, quinquefidus, bractea tripartita suffultus.

Petala squamaeformia.

Stam. petalis opposita, filamentis abbreviatis. Antherae biloculares, apice poro gemino dehiscentes.

Stylus brevissimus.

Caps. trivalvis, trisperma. Sem. strophiole laciniato.

Lasiopetaleae. *Vent.* Malmais. t. 59. *Gay.* in mém. mus. 7. t. 18. 19. *Reichenb.* Mag. t. 39.

936. GUICHENOTIA Gay. (1821.)

Cal. corollinus persistens, quinquefidus, bractea tripartita suffultus.

Petala squamaeformia.

Stam. petalis opposita. Antherae rima longitudinali dehiscentes.

Stylus brevissimus.

Germen quinqueloculare.

Lasiopetaleae. *Gay* in mém. mus. 7. t. 20.

937. THOMASIA Gay. (1821.)

Cal. corollinus campanulatus, persistens, quinquefidus, bractea tripartita suffultus.

Petala squamaeformia.

Stam. filamentis basi connatis, vel appendiculatis, aut sterilibus alternantibus. *Antherae* lineares, rima laterali dehiscentes.

Stylus persistens, elongatus.

Caps. trivalvis, tri-hexasperma. Sem. strophiole crenato.

Lasiopetaleae. Gay in mém. mus. 7. t. 21. [*Lasiopetalum* Bot. mag. 1485. 1486. *Labill.* nov. holl. t. 86.]

938. *WALKERIA* Schreb. (1789.)

Cal. quinquepartitus, persistens.

Petala patentia.

Stam. filamenta capillaria. *Antherae* subrotundae.

Stylus setaceus, stigmate simplici.

Drupae quinque monospermae.

Ochnaeae. [Meesia Gärtn. t. 70.]

939. *EUTHEMIS* Jack. (1822.)

Cal. pentasepalus.

Petala patentia.

Stam. hypogyna. *Antherae* apice poro gemino dehiscentes.

Stylus filiformis.

Bacca pentasperma. Sem. arillata.

940. *MELICYTUS* Forst. (1775.)

Flores dioeci.

Cal. quinquedentatus.

Petala patentissima.

Stamina clavata cyathiformia, antheras quadrisulcas intus gerentia.

Stylus subnullus. Stigma quinquefidum.

Bacca quinquelocularis, pentasperma.

Terebinthaceae. Forst. char. 62. Gärtn. t. 44.

941. *OLINIA* Thunb. (1799.)

Cal. campanulatus, quinquedentatus.

Petala lineari-lanceolata, persistentia.

Squamae basi petalorum insertae, concavae.

Antherae subsessiles, didymae, squamis tectae.

Stylus brevissimus. Stigma subbifidum.

Drupa nuce pentasperma.

942. *TODDALIA* Juss. (1789.)

Flores diclini.

Cal. quinquedentatus.

Petala patentissima.

Stam. hypogyna, filamentis elongatis. *Antherae* subrotundae, biloculares.

Stylus quinquesulcus, stigmate quinquelobo.

Cor. tubulosa, irregularis, hinc longitudinaliter fissa:
limbo bilabiato, labio sup. bi-infer. trifido.

Antherae connatae, barbatae

Stylus simplex. Indusium ciliatum sub stigmatate.

Caps. bilocularis, dissepimento utrinque placentifero.

Lobeliaeae. Schk. t. 269. Willd. hort. berol. t. 85. 86.

Ann. mus. 18. t. 1. f. 1.

83. *Isotoma* Lindl. (1826.)

Cal. quinquepartitus.

Cor. tubulosa, limbo hypocrateriformi, laciniis aequalibus.

Antherae connatae, imberbes, duabus inferioribus
aristatis.

Stylus simplex. Stigma capitatum.

Caps. bilocularis, dissepimento utrinque placentifero.

Lobeliaeae. Lindl. in Bot. reg. 964.

84. *Lechenaultia* R. Br. (1810.)

Cal. quinquepartitus.

Cor. tubulosa, hinc longitudinaliter fissa, bilabiata: lab.
sup. bi-inf. trilobo.

Antherae sub anthesi cohaerentes.

Stigma obsoletum, in fundo indusii bilabiati.

Caps. prismatica, bilocularis, quadrialvis; valvis opposi-
tis, medio septiferis.

Goodeniaceae. Bot. reg. 916.

84. *Goodenia* Sm. (1794.)

Cal. quinquepartitus.

Cor. tubulosa, uni-bilabiata: limbi quinquepartiti laciniis
margine diaphanis subulatis.

Antherae distinctae.

Stigmatis indusium simplex subciliatum.

Caps. bilocularis, dissepimento parallelo. Sem. compressa.

Goodeniaceae. Andr. repoa. 68. Cav. ic. 6. t. 506. 507.

508. Labill. nov. holl. t. 75. [Calogyne R. Br. stylo
trifido differt.]

42. *Wendlandia* Bartl. Cand. (1830.)

Cal. tubo globoso, limbo quinquefido.

Cor. tubulosa, limbo quinquefido.

Stam. summo tubo inserta, antheris exsertis oblongis.

Styl. stigmatate bilobo, raro integro clavato.

Caps. bilocularis, bivalvis. Sem. minima numerosa.

Rubiaceae. [Rondeletia paniculata, tinctoria, exserta Roxb.,
thyrsiflora Roth., proxima Don., cinerea, coriacea, pendula
et longifolia Wallich. densiflora Blum.]

43. *Forgesia* Commers. Juss. (1789.)

Cal. turbinatus, quinquefidus.

Cor. quinquepartita, intus furfuracea.

Stam. *Antherae* oblongae, filamentis adnatae.

Stigma bilobum.

*Caps. semiinfera, bilocularis, polysperma, apice biva
et stylo persistente coronata.*

Saxifrageae. [Defforgia Lam. ill. t. 125.]

844. *PSILOBIUM* Jack. (1826.)

Cal. amplus, quinquepartitus.

Cor. tubus brevis, limbo quinquefido.

Stam. basi corollae inserta.

Stigma clavatum, decemalatum, exsertum.

Fructus siliquaeformis, bilocularis, polyspermus.

845. *ARGOSTEMMA* Wallich. (1825.)

Cal. quadri-quinquedentatus.

Cor. rotata, quadri-quinquepartia.

Antherae conniventes, exsertae.

Styl. stigmatе globoso.

*Caps. disco operculari coronata, bilocularis, superne
diatim dehiscens. Sem. numerosa, angulata.*

γ. Capsulae tri-quinqueloculares.

846. *PHYTEUMA* Matth. (1560.)

Cal. quinquefidus.

Cor. quinquepartita, lacinis elongatis apice cohaerentib

Filamenta basi dilatata.

Stylus simplex. Stigma bi-trifidum.

Caps. trilocularis, placenta centrali.

Campanuleae. Schk. t. 39. Gärtn. t. 30.

847. *TRACHELIUM* Pon. (1617.)

Cal. quinquedentatus.

Cor. tubo gracili, longissimo, limbo quinquefido.

Filamenta basi aequalia.

Stylus exsertus. Stigma subtrilobum.

Caps. trilocularis, basi poris dehiscens.

Campanuleae. Schk. t. 40. Bot. reg. 72. Gärtn. t.

848. *CAMPANULA* Dodon. (1557.)

Cal. quinquefidus.

Cor. campanulata vel subrotata, limbo quinquefido.

Filamenta basi dilatata.

Stigma tri-quinquefidum.

*Caps. tri-quinquelocularis, poris lateralibus vel ap
valvato dehiscens.*

Campanuleae Schk. t. 39. Gärtn. t. 31.

1. *Medium. Calycis sinus reflexi, basi producti. Desj
sel. e cor. Tournef. t. 30. 31.*

2. *Adenophora Fisch., Codonopsis Wallich. Glossoc
Don. Nectarium cylindricum pistillum inferne a
iens. Stylus elongatus. Reichenb. hort. 1. t. 2.
23. 32. 45.*

3. *Campanopsis R. Br., Wahlenbergia Schrad. Cervi
Delil. (fl. aeg. t. 5. abortu triandra). Caps. valvis a
dehiscens. Reichb. ic. 5. t. 480. hort. 2. t. 165.*

Stylus brevis.

Caps. compressa, bipartibilis, dicocca: coccis bi-trivalvibus. Sem. plurima resinosa.

Pittosporae. Car. ic. 4. t. 350. Bot. mag. 1767. [Itea Andr. repos. 314.]

54. **RUYSCHIA** Jacq. (1763.)

Cal. quinquepartitus, imbricatus, involucratus bractea tripartita, cuius lacinia media clavata.

Petala planiuscula.

Stylus subnullus. Stigma cruciatum.

Bacca quinquelocularis polysperma.

Guttiferae. Jacq. amer. t. 51. Sw. fl. ind. occ. 1. t. 11. [Souroubea Aubl. guian. t. 97.]

55. **HYDNOCARPUS** Gärtn. (1788.)

Flores diclini.

Cal. pentasepalus.

Petala cum sepalis alternantia, basi squamis nectariferis.

Bacca globosa: placentatio parietalis. Sem. rugosa.

Flacourtiace. Gärtn. t. 60.

56. **CALISPERMUM** Lour. (1790.)

Cal. quinquefidus, persistens.

Petala concava staminifera.

Stigma crassum.

Bacca unilocularis: placentatio parietalis. Sem. nidulantia, plurima.

57. **BREXIA** Noronh. Thuars. (1810.)

Cal. quinquefidus, persistens.

Petala rotundata, caduca, subcoriacea.

Urceolus fimbriatus staminifer.

Filamenta incurva. Antherae erectae, oblongae, biloculares.

Stylus vix ullus. Stigma quinquelobum.

Drupa quinqueangularis, quinquelocularis: loculi polyspermi.

— — Bot. reg. 730. 872.

58. **ESCALLONIA** Mutis. L. fil. (1781.)

Cal. semiglobosus, quinquedentatus.

Petala calyci inserta.

Stamina calyci imposita. Antherae biloculares.

Stylus erectus. Stigma bilobum.

Caps. baccata, bilocularis, basi irregulariter rumpens.

Placentae in quovis loculo liberae.

Escalloniae. Smith. ic. ined. t. 30. 36. Bot. mag. 2890. [Stereoxylon R. et P. fl. per. 3. t. 234 — 238.]

59. **FLINDERSIA** R. Br. (1814.)

Cal. quinquefidus, persistens.

Petala obtusa, plana.

Discus perigynus, cyathiformis, decomplicatus, subcrenatus.
Stam. disco imposita, sterilia filiformia cum fertilibus alternantia.

Stylus simplex pentagonus. Stigma peltatum, quinquelobum.

Caps. lignea, echinata, quinquelocularis: dissepimenta e placenta centrali prodeuntia. Sem. apice alata.

Melieis affinis. *R. Br.* ad *Flinders.* it. t. 1.

960. *CEDRELA* P. Brown. (1756.)

Cal. quinqueidentatus.

Pet. erecta, in tubi speciem conniventia, cum urceolo staminifero pedicello ovarii inserta.

Stam. sterilia nulla.

Stigma pentagonum. Placenta centralis pentagona.

Caps. quinquevalvis. Sem. compressa, apice alata.

Cedreleae. *P. Brown* jam. t. 10. f. 1.

961. *ANCHIETEA* Hilar. (1824.)

Cal. quinquepartitus.

Cor. irregularis, petalo quinto maximo calcarato.

Caps. maxima, inflata, trivalvis, polysperma. Sem. grandia, compressa, alata.

Violeae. *Hilar.* plant. us. Brasil. t. 19. [*Noisetia* *Mart.* nov. gen. 1. t. 16.]

962. *AMPHIRRHOX* * (1827.)

Cal. quinquepartitus.

Pet. quinque unguiculata, spathulata, subinaequalia, unguibus longis in tubum conniventibus.

Stam. Antherae apice longe mucronatae.

Caps. unilocularis, trivalvis, valvis placentiferis.

Violeae. [*Spathularia* *Hilar.* in mém. mus. 11. t. 24.]

963. *GLOSSARRHEN* Mart. (1822.)

Cal. pentasepalus, sepalis duobus interioribus minimis.

Cor. irregularis, petalo quinto maximo calcarato, basi bicalloso, summis minimis.

Stam. filamenta lata, appendice linguaeformi, duo processibus nectariferis. Antherae dorsales, geminae s. biloculares, loculis distinctis.

Stylus apice incrassatus. Stigma lunato-bicorne.

Caps. unilocularis, trivalvis, valvis placentiferis.

Violeae. *Mart.* nov. gen. 1. t. 15. Nov. act. nat. cur. 12. t. 8. [*Schweiggeria* *Entdeck.* 2. p. 167. *Hilar.* in mém. mus. 11. t. 22. B.]

964. *NOISETTIA* Kunth. (1821.)

Cal. pentasepalus.

Cor. irregularis, petalo quinto maximo calcarato.

Filamenta subconnata, apice membrana acuta, duo processibus filiformibus nectariferis. Antherae dorsales biloculares, loculis distinctis.

Stylus clavatus.

Caps. unilocularis, trivalvis, valvis placentiferis. Sem. ovato-subglobosa.

Violeae. Humb. nov. gen. 5. t. 499. a. b.

965. *CORYNOSTYLIS* Mart. (1822.)

Cal. pentasepalus, subaequalis.

Cor. irregularis, petalo quinto maximo.

Stam. filamentis membranaceis, apice ligula auctis, per paria connatis, quinto libero, appendiculis abbreviatis. Antherae biloculares.

Stylus clavatus.

Caps. lignescens, corticata, unilocularis, trivalvis. Sem. subquadrato-compressa, rugosa, valvis affixa.

Violeae. Mart. nov. gen. 1. t. 17. 18. [*Viola Hybanthus* Aubl. guian. t. 319. Calyptrion Ging. in Cand. prodr.]

966. *HYBANTHUS* Jacqu. (1763.)

Cal. quinquepartitus, irregularis, postice gibbus.

Petala inaequalia: inferius basi saccatum, medium elongatum bifidum.

Stam. membranacea, in tubum connata, basi exappendiculata. Antherae dorsales biloculares. Glandula conchiformis ad basin staminum inferiorum.

Stylus uncinatus, stigmate acuto.

Caps. obovata, trivalvis, valvis placentiferis.

Violeae. Jacqu. amer. t. 175. f. 24. 25. Humb. nov. gen. 5. t. 494.

967. *SOLEA* * (1800.)

Cal. pentasepalus, basi truncatus.

Cor. irregularis, ecalcarata, petalo quinto maximo, labelli specie.

Filamenta distincta, apice in ligulam producta. Glandulae nectariferae duae perigynae.

Stylus apice uncinatus.

Caps. unilocularis, trivalvis, valvis placentiferis.

Violeae. Schrad. diar. 1800. 2. t. 6. [*Jonidium Vent.* Malmais. t. 27. Humb. nov. gen. 5. t. 495—500. Hilar. plant. us. bras. t. 9.]

968. *VIOLA* Virg.

Cal. quinquepartitus, basi productus.

Petala subinaequalia, supremum cornutum.

Stam. filamenta membranacea, apice in membranulam producta, duo superiora processibus nectariferis. Antherae conniventes.

Stigma uncinatum.

Caps. unilocularis, trivalvis, valvis placentiferis.

Violeae. Schk. t. 269. Gärtn. t. 112.

969. *IMPATIENS* Dodon. (1559.)

Cal. disepalus, caducus.

Petalc quatuor inaequalia, quinto abortiente, duobus subcoactis, inferius calcaratum. *Antherae* coalitae, filamentis brevissimis.

Stigma sessile, pentagonum.

Caps. quinquelocularis, quinquevalvis: dissepimentis tenuissimis: valvis elastice dissilientibus, spiratim tortis.

Placenta centralis filiformis, tuberculis alternis seminiferis.

Geraniis affinis. *Schl.* t. 270. *Gärtn.* t. 113. [*Balsamina* *Hook. exot. fl.* t. 137. 141.]

970. *LAVRADIA* Vand. (1788.)

Cal. pentasepalus.

Petala aequalia, breviter unguiculata.

Squamae petaloideae in tubum conicum stamina includentem, demum irregulariter dehiscentem, connatae.

Antherae squamis oppositae, filamentis brevissimis.

Stylus subulatus.

Caps. subtrilocularis, trivalvis, valvis placentiferis.

Droseraeae. *Mart. nov. gen.* 1. t. 22. 23.

971. *SAUVAGESIA* Jacq. (1763.)

Cal: pentasepalus.

Petala ovata.

Corona fimbriarum subclavatarum inter petala et squamas petaloideas.

Squamae petaloideae in tubum conicum conniventes, genitalia includentes.

Stam. cum squamis alternantia. Filamenta brevissima. *Antherae* biloculares, erectae, latere dehiscentes.

Stylus simplex.

Caps. subtrilocularis, trivalvis, marginibus valvarum placentiferis.

Droseraeae. *Grundz.* t. 6. f. 13. 14. *Mart. nov. gen.* 1. t. 24. 25.

972. *BYBLIS* Salisb. (1808.)

Cal. pentasepalus.

Pet. quinque.

Antherae apice biporosae.

Stylus simplex, stigmate bilobo.

Caps. bilocularis, bivalvis, polysperma.

Droseraeae. *Salisb. parad. lond.* t. 95.

973. *GALAX* L. (1751.)

Cal. quinquepartitus.

Petala quinque.

Stam. decem, quorum alterna sterilia urceolo perigyno inserta. Filamenta subulata. *Antherae* biloculares.

Caps. trilocularis, polysperma.

Ericaceae. [*Erythorrhiza Michaux fl. bor. amer.* t. 36. *Gärtn.* t. 210. *Solanandra Vent. Malm.* t. 69. *Blandfordia Andr. rep.* 343.]

974. *ITEA* Gronov. (1743.)

Cal. quinquefidus, minimus, coloratus.

Petala lanceolata, calyci inserta.

Stam. Filamenta subulata, calyci inserta. *Antherae* subrotundae, incumbentes.

Stylus cylindricus. *Stigma* obtusum, emarginatum.

Caps. bivalvis, valvis inflexis dissepimentum formantibus.

— — *Trew.* sel. Ehret. t. 98. *Gärtn.* t. 209.

975. *BRUNIA* L. (1737.)

Flores aggregato - capitati. *Receptaculum* pilosum. *Involucrum* imbricatum.

Cal. pentasepalus.

Petala unguiculata.

Stam. filamenta capillaria, unguibus petalorum inserta.

Antherae ovato - oblongae.

Stylus bifidus.

Drupa bilocularis polysperma.

Bruniaceae. Willd. in act. acad. monach. 1809. t. 6. 7.

Gärtn. t. 30. *Wendl.* coll. t. 11. 21. 35. 45. 57. *Reichenb.*

hort. 1. t. 100. *Ann. des sc. nat.* 8. t. 36.

976. *HUMBOLDTIA* Vahl. (1794.)

Cal. quadripartitus.

Pet. quinque subaequalia, fauci calycis inserta.

Stam. calyci inserta.

Germen pedicellatum. *Stylus* filiformis.

Legumen oblongum, compressum.

Leguminosae. [Batschia Vahl. symb. 3. t. 56.]

977. *DIMORPHANDRA* H. Schott. (1827.)

Cal. urceolatus quinquefidus.

Petala quinque patentissima.

Stam. fertilia, petalis inserta: sterilia totidem petaloidea, spathulata.

Lomentum lignosum, polyspermum, latice balsameo farctum.

Leguminosae.

978. *DRYMARIA* Willd. Röm. et Schult. (1819.)

Cal. quinquepartitus.

Pet. bifida.

Stam. hypogyna, basi subconnata.

Stylus simplex. *Stigma* trifidum.

Caps. unilocularis, trivalvis, penta - polysperma.

Caryophylleae. *Humboldt* nov. gen. 6. t. 515. 516.

979. *MOLLIA* Willd. (1806.)

Cal. quinquepartitus, lacinii margine membranaceis.

Petala quinque.

Stylus simplex. *Stigma* trilobum.

Caps. unilocularis, trivalvis, polysperma.

Paronychieae. Willd. hort. berol. 1. t. 11. Schrank hort. monach. t. 25. [Polycarpaea Lam. diar. hist. nat. t. 25. Delil. fl. aegypt. t. 24. Hagea Pers. Lahaya Röm. et Schult.]

980. *EUPAREA* Banks. Gärtn. (1788.)

Cal. pentasepalus.

Petala quinque, vel decem, duodecim, angusta, patentia.
Stylus persistens, setaceus.

Caps. baccata, sphaerica, unilocularis, evalvis. Placenta globosa, libera. Sem. subangulata.

Primuleae. Gärtn. t. 50.

981. *OLDENLANDIA* Plum. (1703.)

Cal. quadri-quinquedentatus, semiinferus.

Petala quatuor, quinque, calyci inserta.

Stylus subbifidus.

Caps. bilocularis, polysperma.

Saxifrageae? [Heuchera Murr. in comm. gött. 1772 t. 1]

982. *WIBELIA* Pers. (1805.)

Cal. quinquepartitus.

Petala in tubum conniventia, apice reflexa.

Stam. disco inserta. Antherae approximatae.

Stylus simplex, stigmate bilobo.

Germen biloculare.

— — [Paypayrola Aubl. guian. t. 99.]

B. *Fructus inferi.*

a. *Corolla tripetala.*

983. *STREBLITZIA* Ait. (1789.)

Spathae monophyllae imbricatae coloratae, universales partiales involvens.

Cal. coloratus, tripetaloides, subbilabiatus. *Sepala* lanceolata, inferius naviculata.

Cor. irregularis genitalia involvens. *Petala* duo longiora acuminata, postice semisagittata, tertium brevius, ovatum.

Filamenta quinque filiformia receptaculo inserta, petalis conniventibus inclusa. Antherae lineares.

Stylus filiformis. Stigmata tria.

Caps. trilocularis, trivalvis. Placenta centralis.

Museae. Ait. hort. kew. t. 2. Rottböll in nov. act. soc. dan. 4. t. 1. Redout. lil. 77. 78. Bot. reg. 516.

b. *Corolla pentapetala.*

984. *RIBES* Fuchs. (1542.)

Cal. ventricosus, quinquefidus, coloratus.

Petala fauci calycis inserta, cum laciniis eius alternantia.

Stam. fauci calycis inserta, cum petalis alternantia. Antherae biloculares, intus dehiscentes.

Stylus bi-trifidus.

Bacca unilocularis, polysperma. Placentatio subperipherica.

Glossulariaceae. Sturm. fl. germ.

985. *HERDERA* Virg.

Cal. quinquedentatus, dentibus deciduis.

Petala patentia, lanceolata.

Stam. cum petalis alternantia; filamenta subulata. Antherae incumbentes, basi bifidae.

Germen semiimmersum calyci. *Stylus* simplex.

Bacca pentasperma, limbo calycino late coronata.

Araliaceae. Schk. t. 49.

986. *SPHAEROCARYA* Wallich. (1825.)

Cal. quinquepartitus.

Petala cum staminibus alternantia.

Squamae quinque fimbriatae inter stamina et oppositas calycis lacinias.

Stylus indivisus.

Drupa nuce monosperma.

987. *CARPODETUS* Forst. (1775.)

Cal. turbinatus, quinquedentatus.

Petala et stamina alterna calycis margini inserta.

Stigma capitatum.

Bacca globosa, quinquelocularis, pentasperma.

— — Forst. char. 17.

988. *PLECTRONIA* N. L. Burm. (1768.)

Cal. turbinatus, obsolete quinquedentatus: ore squamis villosis clauso.

Petala quinque et stamina alterna calycis fauci inserta.

Stam. filamenta brevissima. Antherae geminae sub squamis calycis.

Stylus filiformis. *Stigma* ovatum.

Bacca oblonga disperma.

— — [Rhamnus J. Burm. afr. t. 94.]

989. *GOUANIA* Jacqu. (1763.)

Flores polygami.

Cal. campanulatus quinquefidus.

Petala fauci calycis inserta, cucullata.

Stam. filamenta subulata. Antherae subrotundae, petalis tectae.

Stylus semitrifidus.

Caps. trilocularis, subulata, trisperma.

Rhamnaceae. Roxb. corom. 1. t. 98. Gärtn. t. 183.

990. *CRUMENARIA* Mart. (1822.)

Cal. campanulatus quinquefidus coloratus.

Petala cuculliformia, inter lacinias calycis inserta.

Stam. filamenta calyci inserta. Antherae petalis reconditae, biloculares.

Stylus simplex. Stigmata tria.

Caps. papyracea, tricocca, coccis margine alatis monospermis.

Rhamneae. Mart. nov. gen. 2. t. 160.

991. **STRUMPFIA** Jacqu. (1763.)

Cal. minimus quinquedentatus; persistens.

Pet. ovato-oblonga, patentia.

Stam. Antherae sessiles in corpus ovatum connatae.

Stylus subulatus, stigmate obtuso.

Bacca monosperma.

— — [Tournefortia Plum. ic. t. 251.]

992. **GRONOVIA** Houst. (1727.)

Cal. quinquepartitus, subcoloratus.

Petala linearia, cum laciniis calycis alternantia erecta.

Stam. libera, cum petalis alternantia.

Stylus simplex, stigmate capitato.

Bacca monosperma.

Cucurbitaceae. Reliqu. Houst. ed. Banks. t. 2. f. 7.

Jacqu. ic. rar. 2. t. 338.

993. **ARGOPHYLLUM** Forst. (1775.)

Cal. quinquefidus.

Petala lanceolata.

Urceolus pentagonus, apice multifido-fimbriatus, staminibus ambiens.

Stigma capitatum.

Caps. trilocularis.

— — Forst. char. 15. Gärtn. t. 210.

994. **JASIONE** L. (1737.)

Flores capitati involucri. Involucrum imbricatum. Receptaculum nudum.

Cal. quinquefidus.

Petala angusta, basi nexa.

Antherae coalitae.

Stylus elongatus. Stigma clavatum, subbilobum.

Caps. bilocularis.

Campanuleae. Schk. t. 266. Gärtn. t. 30.

995. **LIGHTFOOTIA** Herit. (1788.)

Cal. pentasepalus.

Petala lanceolata.

Stam. filamenta basi dilatata, fornicata.

Stigma tri-quinquefidum.

Caps. tri-quinquelocularis.

Campanuleae. Herit. sert. angl. t. 4. 5. Smith. exot. bot. 2. t. 69.

996. **CYPHIA** Berg. (1767.)

Cal. quinquepartitus.

Cor. pentapetala, subbilabiata: petala inferne in tubum conniventia.

Stam. filamentis in cylindrum conniventibus pilosis. Antherae distinctae, erectae, biloculares.

Stylus simplex, stigmate clavato.

Caps. bilocularis.

Campanuleae. Grundz. t. 7. f. 1 — 4. Bot. reg. 625.

997. *LAGOECIA* L. (1737.)

Flores umbellati, involucrati.

Cal. pentasepalus serratus.

Petala appendiculis bifidis capillaribus.

Stylus simplex, stigmate obtuso.

Caryopsis calyce coronata.

Umbelliferae. Schk. t. 48. Gärtn. t. 23.

** *Flores imperfecti.*

A. *Fructus superi.*

AMARANTEAE et CHENOPODEAE.

a. *Alternifoliae.*

998. *CELOSIA* Trag. (1551.)

Cal. quinquepartitus, coloratus, bracteis binis aut ternis suffultus.

Cupula quinquedentata staminifera. Antherae biloculares.

Stylus simplex. Stigma divisum.

Caps. circumscissa, polysperma.

Mart. nov. gen. 2. t. 158. f. 2.

999. *BERZELIA* Mart. (1826.)

Cal. pentasepalus, bracteis binis suffultus.

Tubus staminifer quinquepartitus, lacinii trifidis, lacinula media antheram bilocularem gerente.

Stylus simplex. Stigmata tria.

Caps. circumscissa, polysperma.

[*Celosia glauca.* Wendl. hort. herrenhus. t. 2.]

1000. *LESTIBUDESIA* Thuars. (1806.)

Cal. quinquepartitus, coloratus, bracteis binis suffultus.

Cupula decemdentata, dentibus alternis staminiferis. Antherae biloculares.

Stylus nullus. Stigmata quatuor sessilia.

Caps. circumscissa, polysperma.

Thuars. hist. plant. Afr. t. 16.

1001. *CLADOSTACHYS* Don. (1825.)

Cal. pentasepalus, demum reflexus.

Stam. basi libera.

Stylus nullus. Stigmata tria, linearia, revoluta.

Caps. polysperma.

[*Blitum.* Rumph. amboin. 5. t. 83. f. 2.]

1002. *AMARANTUS* Plin. Matthiol. (1554.)

Flores monoeci.

Cal. tri - pentasepalus.*Stam.* tria aut quinque distincta, antheris bilocularibus.*Stylus* profunde bi - tripartitus.*Utriculus* circumscissus, monospermus.*Schk.* t. 294. *Gärtn.* t. 128. [? *Polychroa* Lour.]1003. *CHAMISSOA* Kunth. (1817.)*Cal.* pentasepalus, bracteis binis suffultus.*Cupula* edentula, staminifera. Antherae biloculares.*Stylus* simplex, stigmatibus duobus revolutis.*Utriculus* circumscissus, monospermus. Sem. arillatum.*Humb.* nov. gen. 2. t. 125.1004. *AËRVA* Forsk. (1775.)*Cal.* pentasepalus, lanatus, bracteis binis suffultus.*Cupula* staminea decempartita, laciniis alternis sterilibus, alternis antheras biloculares ferentibus.*Stylus* simplex, stigmate bifido.*Utriculus* monospermus.*Gärtn.* t. 213.1005. *TRICHINIUM* R. Br. (1810.)*Cal.* pentasepalus, sepalis fructiferis patulis plumosis.

Bracteae binae ad basin.

Cupula staminifera, edentula, antheris bilocularibus.*Stylus* simplex. Stigma capitatum.*Utriculus* monospermus.1006. *PTILOTUS* R. Br. (1810.)*Cal.* pentasepalus, bracteis binis suffultus. Sepala tria interiora fructifera, medio lana cohaerentia, superne patentia, nuda.*Stam.* basi connata. Antherae biloculares.*Stylus* simplex. Stigma capitatum.*Utriculus* monospermus.1007. *DEERINGIA* R. Br. (1810.)*Cal.* quinquepartitus, bracteis tribus suffultus.*Stam.* basi dilatata, connata. Antherae biloculares.*Stylus* brevissimus vel subnullus. Stigmata tria.*Bacca* triloba, polysperma. Sem. funiculis umbilicalibus basi fructus inserta.*Hook.* in bot. mag. 2717.1008. *TRAGANUM* Delil. (1813.)*Cal.* persistens, quinquefidus, bracteis tribus suffultus.*Stam.* filamenta libera. Antherae sagittatae.*Stylus* apice bifidus.*Nux* lignosa, monosperma, calyce inclusa.*Delil.* fl. aegypt. t. 22.

009. *CORNULACA* Delil. (1813.)

Cal. tribracteatus, villis involutus, quinquefidus, lacinia una dorso demum spinifera.

Stam. filamenta libera, cum squamulis linearibus alternantia. Antherae biloculares.

Stylus bifidus.

Utriculus monospermus, calyce inclusus.

Delil. fl. aegypt. t. 22.

b. *Oppositifoliae.*010. *ACHYRANTHES* L. (1737.)

Cal. pentasepalus, bracteis binis saepe spinescentibus suffultus.

Cupula staminifera, filamentis quinque antheras biloculares ferentibus, interpositis squamulis dentatis aut fimbriatis.

Stylus simplex, stigmate capitato.

Utriculus monospermus.

Jacqu. ecl. 1. t. 74. *Gärtn.* 128.

011. *DESMOCHAETA* Cand. (1813.)

Flores subterni glomerati perfecti, interiectis totidem florum abortientium fasciculis in uncinos mutatorum.

Cal. pentasepalus.

Cupula staminifera quinquepartita. Antherae biloculares.

Stylus simplex, stigmate capitato.

Utriculus monospermus.

Jacqu. ecl. 1. t. 75. 96.

012. *PUPALIA* Mart. (1826.)

Flores gemini, bractea diphylla suffulti: altero abortiente in uncinos abeunte.

Cal. pentasepalus.

Cupula staminifera decempartita, filamentis alternis antheras biloculares ferentibus.

Stylus simplex, stigmate capitato.

Utriculus monospermus.

Mart. nov. gen. 2. t. 158. f. 1.

013. *ALTERNANTHERA* Forsk. (1775.)

Cal. pentasepalus, bracteis binis suffultus.

Cupula staminifera subdecemdentata, denticulis alternis antheras uniloculares ferentibus, alternis sterilibus subdentatis.

Stylus abbreviatus. Stigma capitatum.

Utriculus monospermus.

Brandesia Mart. l. c. t. 127.
Buchholzia Mart. l. c. t. 147. 151.]

014. *MOGIPHANES* Mart. (1826.)

Cal. pentasepalus, bracteis binis suffultus, stipitatus, basi quinqueglandulosus, stipitulo demum articulado.

Tubus cylindricus decemfidus, laciniis alternis antheras uniloculares ferentibus, alternis subdentatis.

Stylus brevissimus, stigmate capitato.

Utriculus monospermus.

Mart. nov. gen. 2. t. 134.

1015. *PAFFIA* Mart. (1826.)

Cal. pentasepalus, bracteis binis suffultus.

Tubus staminifer decemfidus, laciniis inciso-fimbriatis, alternis antheras uniloculares gerentibus.

Stylus nullus. Stigma sessile.

Utriculus monospermus.

Mart. nov. gen. 2. t. 122—124. [Sertürnera ib. t. 136—138]

1016. *GOMPHRENA* L. (1737.)

Cal. pentasepalus, bracteis binis concavis suffultus.

Tubus staminifer quinquefidus, laciniis denticulato-incis antheras uniloculares ferentibus.

Stylus simplex. Stigmata duo cylindrica.

Utriculus monospermus.

Mart. nov. gen. 2. t. 103—121. Hook. in bot. mag. 281.

1017. *HEBANTHE* Mart. (1826.)

Cal. pentasepalus, bracteis binis concavis suffultus. Sepala interiora lana stricta cincta.

Tubus staminifer quinquepartitus, laciniis trifidis, lacinulis mediis antheras uniloculares gerentibus, lateralibus simplicibus.

Stylus vix ullus. Stigma capitato-bilobum.

Utriculus monospermus.

Mart. nov. gen. 2. t. 142. [Bragantia Vandell.]

1018. *IBESINE* Patr. Brown. (1756.)

Flores diclini, tandem lanati.

Cal. pentasepalus, bracteis binis concavis suffultus.

Cupula staminifera edentula. Antherae uniloculares.

Stylus subnullus.

Stigmata duo vel tria.

Utriculus monospermus.

Mart. nov. gen. 2. t. 153. 154. [Trommsdorffia ib. t. 139. 145. f. 3. Rosea ib. t. 155.]

1019. *PHYLOXERUS* R. Br. (1810.)

Flores hermaphroditi, post anthesin non mutati.

Cal. pentasepalus, bracteis tribus suffultus.

Cupula edentula staminifera abbreviata. Antherae uniloculares.

Stylus simplex. Stigmata duo.

Utriculus monospermus.

1020. *HOPLOTHECA* Nuttall. (1818.)

Cal. semiquinquefidus, dense tomentosus, bracteis binis suffultus, post anthesin induratus, muricatus.

Tubus staminifer quinquefidus, laciniis medio antheras uniloculares cylindricas gerentibus.

Stigma capitatum, penicillato-multifidum.

Utriculus monospermus, calyce indurato inclusus.

Mart. nov. gen. 2. t. 146. [Gomphrena Herit. stirp. 1. t. 3.]

21. *AYLMERIA* Mart. (1826.)

Cal. pentasepalus, lbracteis duabus suffultus.

Tubus staminifer decemfidus: lacinae exteriores steriles petalis opposita, interiores alternantes, antheris bilocularibus.

Stylus simplex. *Stigma* capitatum.

Caps. evalvis pleiosperma.

22. *BOUSSINGAULTIA* Humb. (1825.)

Cal. membranaceus, persistens, limbo duplici, exter. bi-inter. quinquepartito.

Stam. basi calycis inserta.

Antherae biloculares, posticae.

Stylus simplex. *Stigma* trilobum.

Utriculus monospermus, stylo coronatus.

Humb. nov. gen. 7. t. 645. b.

P A R O N Y C H I E A E.

23. *ILLECEBRUM* Rupp. (1718.)

Cal. pentasepalus, basi hibracteatus. *Sepala* cucullata cornuta, dorso incrassata.

Stam. sepalis opposita: antherae biloculares: sterilia totidem fertilibus interposita.

Stigmata duo sessilia capitata.

Caps. quinquevalvis, monosperma.

Schk. t. 50. Engl. bot. 895. Gärtn. t. 184.

24. *PARONYCHIA* Clus. (1576.)

Cal. quinquepartitus, ebracteatus. *Sepala* subcucullata, dorso aequalia, subaristata.

Stam. fertilia cum sterilibus alternantia.

Stylus simplex, stigmatibus duobus.

Utriculus monospermus stylo terminatus, in calyce quinquevalvi.

Gärtn. t. 128. Delil. fl. aeg. t. 18. [Illecebrum echinatum — serpyllifolium Vill. in Schröd. diar. 1801. 2. t. 4.]

25. *ANYCHIA* Michaux. (1803.)

Cal. pentasepalus, sepalis sub apice saccatis.

Stam. filamenta distincta, fertilia, sterilibus interpositis nullis.

Stylus nullus. *Stigmata* duo aut tria capitata.

Utriculus monospermus calyce cinctus.

[Queria canadensis Orteg. dec. t. 15. f. 2. Gärtn. t. 128.]

1026. *GUILLEMINEA* Kunth. (1823.)*Cal.* campanulatus, quinquefidus.*Stam.* ori calycis inserta. Antherae uniloculares.*Stylus* apice emarginatus.*Utriculus* monospermus calyce cinctus. Sem. funiculo umbilicali longo suffultum.*Humb.* nov. gen. 6. t. 518.1027. *GYMNOCARPUS* Forsk. (1775.)*Cal.* subcampanulatus quinquefidus, coloratus.*Stam.* ori calycis inserta, cum totidem sterilibus alternantia. Antherae bilobae.*Stylus* simplex. Stigma tridentatum.*Utriculus* monospermus, calyce indurato tectus.*Vivian.* fl. lib. t. 10. f. 1.

C H E N O P O D E A E.

1028. *ANREDERA* Juss. (1789.)*Cal.* bipartitus, dorso carinatus.*Stylus* bifidus, stigmatibus duobus.*Caryopsis* calyce bialato tecta.[*Clarisia Abat.* in mem. de Sevilla, 10. p. 418. ic.]1029. *HABLIZIA* Marsch. Bieb. (1810.)*Cal.* pentasepalus, rotatus.*Stam.* sepalis opposita. Antherae globosae didymae.*Stylus* simplex. Stigma trilobum.*Utriculus* circumscissus monospermus.*Marsch. Bieb.* cent. pl. ross. 2. t. 54.1030. *CAROXYLON* Thunb. (1782.)*Cal.* quinquepartitus, bracteis duabus suffultus.*Squamae* membranaceae, ovatae, calycis laciniis insertae, persistentes.*Stam.* perigyna.*Stylus* simplex, stigmatibus duobus.*Caryopsis* calyce cincta, squamis coronata.1031. *RHAGODIA* R. Br. (1810.)

Flores polygami.

Cal. quinquepartitus.*Stylus* bifidus.*Fructus* baccatus monospermus, calyce cinctus.[*Chenopodium Labill.* nov. holl. 1. t. 96.]

R H A M N E A E.

1032. *ALZATEA* R. et P. (1794.)*Cal.* coloratus, pentagonus, quinquefidus.*Stam.* hypogyna.

Stylus simplex.

Caps. obcordata, bilocularis, bivalvis. Sem. membrana obvoluta.

Ruiz. et Pav. prodr. t. 7. fl. peruv. 3. t. 241. s.

1033. *CONDALIA* Cav. (1799.)

Cal. urceolatus, quinquefidus, basi persistens.

Stam. cum lacinis calycis alternantia. *Antherae* basi bilobae.

Discus perigynus glandulosus.

Stylus simplex, stigmatem emarginato.

Drupa monosperma.

Cav. in anal. de cienc. 1. t. 4. ic. 6. t. 525.

1034. *PENNANTIA* Forst. (1775.)

Flores polygami.

Cal. corollinus pentasepalus.

Stam. cum sepalis alternantia. *Antherae* biloculares.

Stylus nullus. *Stigma* trilobum.

Drupa triquetra subdisperma.

— — *Forst. char. 67.*

1035. *EXOCARPUS* Labill. (1798.)

Flores subpolygami.

Cal. quinquepartitus, rotatus.

Stam. basi laciniarum calycis inserta.

Stylus brevissimus.

Stigma obtusum.

Nux monosperma, pedunculo ampliato, baccato imposita.

Santaleae. Labill. it. t. 14.

1036. *CERVANTESIA* R. et P. (1794.)

Cal. quinquepartitus, in fructu auctus, carnosus.

Discus quinquelobus in fundo calycis, persistens, fructui accretus.

Stam. lacinis calycis inserta. *Antherae* biloculares.

Stylus abbreviatus, stigma emarginatum.

Nux monosperma, calyce cincta.

Santaleae. Cav. ic. 5. t. 475.

1037. *AMANOA* Aubl. (1775.)

Flores androgyno-diclini.

Cal. quinquepartitus, bracteatus.

Discus hypogynus glandulosus quinquelobus.

Stam. filamenta brevissima. *Antherae* oblongae sepalis oppositae.

Stigmata tria sessilia, crassa.

Caps. drupaeformis, tricocca. *Cocci* bivalves, monospermi.

Tricoccae. Aubl. guian. t. 101. Adr. Juss. Euphorbiae t. 2. f. 6.

1038. FLÜGGEA Willd. (1806.)

Flores dioeci.

Cal. quinquepartitus.*Stam.* filamenta basi glandulis interstincta. *Antherae* extrorsum dehiscentes.*Stylus* subnullus. *Stigmata* duo aut tria, bifida, reflexa.*Caps.* subbaccata, bi-trilocularis, loculis subdispermis.*Tricoccae.* *Adr. Juss.* *Euphorbiae* t. 2. f. 7.

1039. ANTIDESMA J. Burm. (1737.)

Flores dioeci.

Cal. parvus quinquedentatus.*Annulus* carnosus staminifer. *Filamenta* elongata. *Antherae* didymae.*Stylus* brevis. *Stigmata* quinque.*Drupa* monosperma, stylo coronata.*Rosaceae?* *Lam. ill.* t. 812. *Roxb. corom.* 2. t. 167.*Gärtn.* t. 39. [*Stilago* L.]

1040. LICANIA Aubl. (1775.)

Cal. turbinatus quinquefidus, basi bibracteatus.*Annulus* glandulosus staminifer. *Stam.* lacinii calycis opposita.*Stylus* incurvus, stigmati simplici.*Drupa* monosperma.*Rosaceae.* *Aubl. guian.* t. 45. [*Hedycra* Schreb.]

1041. CERATONIA Galen. Dodon. (1556.)

Flores polygami.

Cal. quinquepartitus.*Stam.* filamenta longissima. *Antherae* magnae didymae.*Stigma* sessile capitatum.*Legumen* quadrangulare isthmis interceptum, pulpa farctum.*Leguminosae.* *Cav. ic.* 2. t. 113. *Gärtn.* t. 146.

1042. GLAUX Dodon. (1556.)

Cal. quinquepartitus coloratus.*Stam.* hypogyna.*Stylus* conicus. *Stigma* capitatum.*Caps.* unilocularis, pentasperma. *Placenta* centralis pentagona.*Primulaeae.* *Schk.* t. 50. *Gärtn.* t. 184.B. *Fructus inferi.*

1043. PHYLICA L. (1737.)

Cal. corollinus quinquefidus.*Squamulae* cucullatae intra calycem stamina obtegentes.*Stam.* abbreviata sub squamulis.*Stylus* simplex, stigmati obtuso.*Caps.* tricocca, subbaccata.*Rhamneae.* *Berl. mag.* 8. t. 8. *Gärtn.* t. 24.

1044. *THESIUM* L. (1737.)

Cal. corollinus quinquefidus, (intus glaber vel barbatus).
Stam. extus villosa, basi laciniarum calycis inserta. Antherae biloculares.

Stylus simplex. Stigma capitatum.

Nux monosperma, striata.

Santaleae. Schk. t. 51. Reichenb. ic. 5. t. 452 — 458.
 Gärtn. t. 86.

1045. *LEPTOMERIA* R. Br. (1810.)

Cal. rotatus, quinque- (quadri-) fidus, nudiusculus.

Discus epigynus quinque- (quadri-) lobus.

Stamina basi laciniarum calycis inserta. Antherae biloculares.

Stigma emarginato - bilobum vel quinquefidum.

Drupa baccata, mono - disperma.

Santaleae. [Thesium Labill. nov. holl. t. 93.]

1046. *CHORETRUM* R. Br. (1810.)

Cal. corollinus quinquepartitus, basi bractea s. calyculo quinquedentato suffultus. Lacinae calycis fornicatae.

Stam. sub fornicibus calycis. Antherae quadriloculares, quadrivalves.

Stigma stellatum.

Drupa ?

Santaleae.

1047. *HAMILTONIA* Mühlenb. Willd. (1807.)

Flores dioeci aut polygami.

Cal. corollinus subcampanulatus, quinquefidus.

Discus perigynus quinquedentatus.

Stam. basi laciniarum calycis inserta. Antherae biloculares, (basi subbarbatae).

Stylus simplex, stigmate integro aut bilobo.

Drupa nuce monosperma.

Santaleae. [Comandra Nuttall. Pyrolaria Michaux.]

1048. *NYSSA* L. (1737.)

Flores polygami.

Cal. corollinus quadripartitus, deciduus.

Stam. in masculis decem, calyci inserta, antheris didymis, longitudine filamentorum: in hermaphroditis quinque, antheris simplicibus.

Stylus subulatus incurvus, stigmate simplici.

Drupa monosperma.

Santaleae. Lam. ill. t. 851. Michaux. arbr. forest. de l'Amér. t. 19 — 22. Gärtn. t. 216.

1049. *CEVALLIA* Lag. (1816.)

Cal. corollinus decempartitus, laciniis linearibus erectis.

Stam. filamenta basi laciniarum calycis inserta. Antherae conniventes, sagittatae, biloculares, intus longitudinali-

ter dehiscentes, appendiculo membranaceo cylindrico cavo terminatae.

Stylus filiformis. *Stigma* capitatum.

Nux globosa monosperma.

Cum *Echinope* iungenda?

1050. *MYOSCHILLOS* Ruiz et Pav. (1799.)

Cal. corollinus pentasepalus, patens.

Stam. receptaculo inserta. *Antherae* subrotundae.

Stylus trigonus, brevissimus. *Stigma* bi-trifidum.

Drupa monosperma.

Thymelaeae. Ruiz. et Pav. fl. peruv. 3. t. 242. f. 2.

1051. *CONOCARPUS* Houst. (1727.)

Flores capitati.

Cal. urceolatus, quinquefidus, deciduus.

Stam. quinque (decem) calyci inserta. *Antherae* cordatae biloculares, longitudinaliter intus dehiscentes.

Stylus stigmate simplici.

Strobilus e fructibus imbricatis, suberosis, monospermis.

Combretaeae. Jacqu. amer. t. 52. f. 1. *Catesby* carolin. 2. t. 33. *Gärtn.* 2. 216.

D I G Y N I A.

* *Flores perfecti.*

I. COROLLA MONOPETALA.

Fructus superi.

A. CONTORTAE.

a. *Carisseae: drupiferae s. bacciferae.*

1052. *RAUWOLFIA* Plum. (1703.)

Cal. quinquepartitus persistens.

Cor. infundibuliformis quinquefida.

Stam. libera, exserta. *Antherae* erectae, simplices.

Styli connati.

Drupae duae monospermae.

Plum. ic. t. 236. *Gärtn.* t. 52.

1053. *ALYXIA* Banks. R. Br. (1810.)

Cal. quinquefidus.

Cor. hypocrateriformis, fauce nuda.

Stam. inclusa, libera.

Styli subcohaerentes.

Drupae duae monospermae, pedicellatae, putamine semibiloculari. Semen semibipartitum. Albumen ruminatum, corneum.

[*Gynopogon* Forst. char. 18. *Labill.* nov. Caled. t. 34.]

1054. *COPROSMA* Forst. (1775.)

Flores polygami.

Cal. quinquepartitus.*Cor.* campanulata, quinque - septemfida.*Stam.* quinque, septem, exserta.*Styli* divaricati.*Bacca* disperma.

Forst. char. 69. Labill. nov. holl. t. 95.

1055. *MELODINUS* Forst. (1775.)*Cal.* quinquedentatus.*Cor.* tubo cylindrico, limbo hypocrateriformi, laciniis obliquis. Faux obsessa squamis bifidis, vel squamis vicenis duplici serie.*Stam.* libera inclusa.*Styli* connati.*Stigmata* capitata.*Bacca* globosa, polysperma.

Forst. char. 19. Bot. reg. 834. Labill. nov. Caled. t. 33.

b. *Apocynae. Fructus folliculares.*1056. *PLUMERIA* Tournef. (1719.)*Cal.* minimus quinquefidus.*Cor.* infundibuli - hypocrateriformis, fauce nuda, laciniis limbi obliquis.*Stam.* inclusa, libera.*Styli* breves. *Stigmata* emarginata.*Folliculi* ventricosi. Sem. basi alata.

Bot. reg. 114. 480. 510. 780.

1057. *CRYPTOSTEGIA* R. Br. (1810.)*Cal.* quinquepartitus, laciniis elongatis lanceolatis incisodentatis.*Cor.* campanulato - infundibuliformis. Squamae tubi bifidae, antheras obtegentes.*Stam.* filamenta brevissima. Antherae conniventes, incurvae, thodi modo stigma contegentes.*Stigmatibus* latera glandulis quinque spathulatis instructa, antheris respondentia.

Bot. reg. 435.

1058. *PERIPLOCA* Dodon. (1563.)*Cal.* quinquefidus.*Cor.* quinquepartita. Cornua quinque acuta ad basin laciniarum.*Stam.* Antherae plicis gynostegii (tecti stylorum) reconditae, simplices, manubriatae. Antherae succedaneae apice barbatae, filamentis distinctis.

Schk. t. 53. Von der Natur und dem Bau der Gew. t. 5. f. 26.

1059. *HEMIDESMUS* R. Br. (1810.)*Cal.* quinquefidus.*Cor.* quinquepartita. Squamae quinque obtusae sub sinibus.*Stam.* Antherae Periplocae. Antherae succedaneae imberbes.1060. *GYMNANTHERA* R. Br. (1810.)*Cal.* quinquefidus.*Cor.* hypocrateriformis. Squamae ad faucem quinque, aristatae.*Antherae* Periplocae. Antherae succedaneae imberbes.*Gynostegium* supra bimucronatum.c. *Asclepiadeae*.

Antherae geminae, filamentis brevibus glandula nexis, pollinis granulis fusiformibus e sutura antherae cartilaginea provenientes. *Gynostegium* (columna pistilla continens et obtegens) plicis lateralibus antheras recondens, apice planum peltatum. *Styli* duo conniventes. *Corona* saepius interior perigyna. *Folliculi* seminibus comosis.

α. *Antherae geminae basi unitae.*αα. *Gynostegium simplex planum.*1061. *SECAMONE* P. Alpin. (1640.) R. Br. (1810.)*Cal.* quinquefidus.*Cor.* rotata. *Corona* perigyna pentaphylla.*Antherae* quaternae erectae *gynostegio* affixae.1062. *STAPELIA* L. (1737.)*Cal.* quinquefidus.*Cor.* subcarnosa, rotata, quinquefida.*Corona* perigyna plerumque duplex, externa ligulas, interna rostra cum alis s. appendiculis formans.*Antherae* geminae basi unitae.

1. *Duvalia* Haw. Faux corollae in orbiculum solidum elevata: laciniae corollae replicatae: rostra alaeque abbreviatae obtusae: ligulae coalitae in scutum sublime. *Jacqu.* *Stapel.* t. 41 — 45.

-2. *Orbea* Haw. Faux corollae similis: laciniae planae: rostra elongata clavata: ligulae patentes subtrifidae. *Jacqu.* ib. t. 24.

3. *Tromotriche* Haw. Faux corollae simplex: rostra clavata: alae abbreviatae. *Jacqu.* ib. t. 5. 6.

4. *Podanthes* Haw. Faux corollae simplex. Alae rostraque brevissima gibbosa, *gynostegio* incumbentia: ligulae latae distinctae. *Jacqu.* ib. t. 40.

5. *Obesia* Haw. Faux corollae simplex. Alae rostraque brevissima gibbosa: ligulae coalitae in scutum sublime. *Jacqu.* ib. t. 34. 46.

6. *Gonostemma* Haw. Faux corollae simplex. Rostra gibbosa, elongata: alae nullae. *Jacqu.* ib. t. 1 — 4. 47.

7. *Caruncularia* Haw. Rostra torosa, clavata, tuberculata.
Jacqu. ib. t. 35. 36.

8. *Piuranthus* R. Br. Corona perigyna simplex, foliolis
dorso dentatis. Masson. Stapel. t. 24. 31.

1063. *HEURNIA* R. Br. (1810.)

Cal. quinquefidus.

Cor. campanulata decemdentata.

Corona perigyna duplex, externa lobos fissos, interna
cornicula subulata sistens.

Jacqu. Stapel. t. 47—55.

1064. *BRACHYSTELMA* R. Br. (1823.)

Cal. quinquefidus.

Cor. campanulata, quinquefida, sinibus angulatis.

Corona perigyna monophylla quinqueloba. Lobi antheris
oppositi.

Gynostegii membranulae plicatae nullae.

Bot. mag. 2343. Bot. reg. 722.

1065. *CARALLUMA* R. Br. (1810.)

Cal. quinquepartitus.

Cor. subrotata quinquepartita.

Corona perigyna decaphylla, foliolis quinque antheris op-
positis integris, reliquis bipartitis subulatis.

Gynostegium exsertum.

[*Stapelia* Roxb. corom. 1. t. 30.]

1066. *CEROPEGIA* L. (1747.)

Cal. quinquepartitus.

Cor. basi ventricosa, tubo infundibuliformi, limbi laciniis
conniventibus lingulatis.

Corona perigyna duplex: exter. monophylla quinqueloba;
interior pentaphylla, foliolis linearibus cum antheris
alternantibus.

Gynostegium inclusum.

Roxb. corom. 1. t. 7—10. Bot. reg. 626.

bb. *Gynostegium processibus auctum.*

1067. *HOYA* R. Br. (1810.)

Cal. quinquefidus.

Cor. rotata, quinquefida.

Processus gynostegii quinque carnosii plani, angulo in-
teriori in dentem antherae incumbentem producto.

Stigma papillosum in medio gynostegio prominulum.

Reichenb. Mag. t. 32. [Schollia Jacqu. ecl. t. 2.]

1068. *TYLOPHORA* R. Br. (1810.)

Cal. quinquefidus.

Cor. rotata, quinquepartita.

Processus gynostegii quinque carnosii, angulo interiori
edentulo.

Stigma muticum.

Colebrook. in Linn. transact. 12. t. 16.

1069. *PERGULARIA* L. (1767.)*Cal.* quinquefidus.*Cor.* hypocrateriformis, tubo urceolato.

Processus gynostegii quinque lamellosi, intus edentati vel lacinula aucti.

Stigma muticum.*Smith.* ic. pict. t. 16. Bot. reg. 412. [*Maradenia* R. Br. Bot. reg. 489.]1070. *DISCHIDIA* R. Br. (1810.)*Cal.* quinquepartitus.*Cor.* urceolata, quinquefida.

Processus gynostegii quinque, foliola bifida lacinii subulatis patentibus recurvis.

Stigma muticum.*Colebrook.* in *Linn. transact.* 12. t. 15. t.cc. *Nec corona perigyna, nec processus gynostegii.*1071. *GYMNEMA* R. Br. (1810.)*Cal.* quinquefidus.*Cor.* suburceolata, quinquefida. Faux saepe coronati denticulis aut squamulis inter lacinias.1072. *SARCOLOBUS* R. Br. (1810.)*Cal.* quinquefidus.*Cor.* subrotata, quinquefida, fauce nuda.*Folliculi* ventricosi, carnosii. Sem. marginata.*Wallich.* in *asiat. research.* 12. t. 4. 5.β. *Antherae geminae, transversim affixae et dehiscentes.*1073. *GONOLOBUS* Michaux. (1803.)*Cal.* quinquefidus.*Cor.* subrotata, quinquefida.*Corona* gynostegii quinqueloba subcarnosa.*Hook.* in *bot. mag.* 2799. Bot. reg. 252. *Humb.* nov. gen. 3. t. 238. 239.1074. *MATELEA* Aubl. (1775.)*Cal.* quinquefidus.*Cor.* tubulosa, limbo hypocrateriformi.*Corona* gynostegii lobata.*Folliculi* costati. Sem. nuda.*Aubl.* *guian.* t. 109. [*Hostea Willd.*]γ. *Antherae geminae apice unitae pendulae.*aa. *Gynostegium corona cinctum.*1075. *ASCLEPIAS* Dioscor.*Cor.* quinquepartita, revoluta.*Corona* cucullis quinque cornigeris constat.*Folliculi* glabri.*Schk.* t. 56. *Jacqu.* ecl. t. 28.

1076. *GOMPHOCARPUS* R. Br. (1810.)
Cor. quinquepartita, reflexa.
Corona cucullis intus aceratis, saepe unidentatis constat.
Folliculi ventricosi, echinati.
Reichenb. Mag. t. 28. [Acerates Elliott. Anantherix Nutt.]
1077. *ENSLENIA* Nutt. (1818.)
Cor. quinquepartita, laciniis erectis conniventibus.
Corona quinquepartita, petaloidea, laciniis truncatis, dente gemino aristato terminatis.
1078. *OXYSTELMA* R. Br. (1810.)
Cor. rotata, patens.
Gynostegium exsertum.
Corona pentaphylla, foliolis acutis indivisis.
Folliculi laeves.
Roxb. corom. 1. t. 11.
1079. *XYSMALOBIUM* R. Br. (1810.)
Cor. campanulata, basi ventricosa.
Corona prominula e laminis quinque obtusis carnosis, denticulis totidem alternis.
Folliculi paleacei.
1080. *CALOTROPIS* R. Br. (1810.)
Cor. subcampanulata, basi in angulis saccata.
Corona e laminis gynostegio appressis, lanceolatis, basi revolutis, apice bidentatis.
Folliculi laeves.
Bot. reg. 58. Franc. Hamilt. in Linn. transact. 14. p. 248.
1081. *PODOSTIGMA* Elliott. (1817.)
Cor. campanulato-rotata.
Gynostegium stipitatum.
Corona e foliolis erectis compressis, apice liberis, recurvis.
[Stylandra Nutt.]
1082. *LACHNOSTOMA* Kunth. (1818.)
Cor. hypocrateriformis, fauce barbata.
Corona e foliolis carnosis, lunato-bilobis.
Humb. nov. gen. 3. t. 232.
1083. *MACROSCOPIS* Kunth. (1818.)
Cal. quinquepartitus, elongatus, bracteatus.
Cor. carnosae, basi ventricosa, limbo patente quinquefido.
Corona e squamis quinque carnosis, intus replicatis, una cum antheris alternantibus fauci corollae insertis.
Gynostegium breve scutelliforme.
Humb. nov. gen. 3. t. 233. [Schubertia Mart. nov. gen. 1. t. 33.]
1084. *CANAHIA* R. Br. (1810.)
Cal. quinquepartitus.
Cor. campanulata, limbo quinquepartito.

Corona apici gynostegii imposita, e foliolis quinque basi incrassatis, apice subulatis.

Vahl. symb. 1. t. 7. *Delil. in Caillaud it. Meroë*, 2. t. 64.

1085. *CYNANCHUM* [Dioscor.] L. (1737.)

Cor. quinquepartita rotata.

Corona petaloidea, cum antheris alternans, vel simplex quinqueloba; vel duplex, exter. decemfida, inter. quinquepartita.

Schk. t. 54. 55.

1086. *DITASSA* R. Br. (1810.)

Cor. quinquepartita rotata, intus barbata vel glandulosa.

Corona duplex: exter. quinquepartita, laciniis acuminatis elongatis patentibus: inter. brevior pentaphylla, exteriori antherisque opposita.

Mart. nov. gen. 1. t. 31.

1087. *DIMIA* R. Br. (1810.)

Cor. subrotata.

Corona duplex: exter. brevis quinquepartita, denticulis interiectis: inter. pentaphylla erecta, foliolis basi solutis apice subulatis.

Folliculi hispidi, valvati.

[*Cynanchum Jacqu. ic. rar.* 1. t. 54. *Asclepias Pal. Beauv. fl. ovar.* 1. t. 56.]

1088. *SARCOSTEMMA* R. Br. (1810.)

Cor. rotata, quinquefida.

Corona duplex: ext. cyathiformis s. annulata crenata: inter. pentaphylla, carnosa.

Folliculi laeves.

Humb. nov. gen. 3. t. 229.

1089. *PHILIBERTIA* Kunth. (1818.)

Cor. urceolato-rotata, decemdentata, denticulis alternis minoribus.

Corona duplex: ext. in fundo corollae annulata, integra, carnosa: inter. in apice gynostegii pentaphylla.

Gynostegium apice bimucronatum.

Humb. nov. gen. 3. t. 230.

1090. *EUSTEGIA* R. Br. (1810.)

Cor. rotata, quinquefida.

Corona triplex pentaphylla: interior cum exterioribus alternans, media foliolis trilobis.

N. Entdeck. 1. t. 1. f. 5—10.

1091. *METAPLEXIS* R. Br. (1810.)

Cor. rotata, quinquefida.

Corona quinquepartita: lacinae cum antheris alternantes.

Gynostegium rostratum.

[*Cynanchum fimbriatum Kunth. in Humb. nov. gen.* 3. t. 234.

gynostegium apice capitulo stipitato.]

1092. *OXYPETALUM* R. Br. (1810.)

Cor. campanulata, laciniis elongatis angustissimis.

Corona pentaphylla carnea.

Filamenta seu retinacula antherarum sursum in appendiculam et dentes subulatos abeuntia.

Stigmata e gynostegio longe producta, divaricata.

Humb. nov. gen. 3. t. 231. *Mart.* nov. gen. 1. t. 29. 30. [*Gothofreda Vent.*]

1093. *PHYSIANTHUS* Mart. (1825.)

Cor. campanulata, basi ventricosa.

Corona pentaphylla sublibera, carnea, foliolis extrorsum apice cucullatis.

Stigmata apiculata exserta.

Mart. nov. gen. 1. t. 32.

bb. *Corona* desciscens vel nulla.

1094. *HOLOSTEMMA* R. Br. (1810.)

Cor. subrotata, quinquefida: tubo intus annulato.

Gynostegium muticum.

[*Ada-kodien Rheed.* malab. 9. t. 7.]

1095. *METASTELMA* R. Br. (1810.)

Cor. subcampanulata: faux coronata dentibus quinque exsertis, in tubum decurrentibus.

Gynostegium muticum.

[*Periploca Plum.* ic. t. 215. f. 1.]

1096. *MICROLOMA* R. Br. (1810.)

Cor. urceolata, tubo ventricoso angulato, fauce nuda.

Squamae quinque medio tubo insertae, cum totidem fasciculis villorum alternantes.

Gynostegium apiculatum.

[*Ceropegia* 2. *Lam.* ill. t. 179.]

1097. *ARAUJA* Brot. (1817.)

Cor. campanulata, basi saccis quinque nectariferis.

Filamenta latissima auriculata dentata.

Brot. in *Linn. transact.* 12. t. 4.

1098. *ASTEPHANUS* R. Br. (1810.)

Cor. subcampanulata, fauce tuboque nudis.

Gynostegium apiculatum.

B. GENTIANEA E.

1099. *GENTIANA* Dioscor.

Cal. campanulatus, quadri-quinquefidus.

Cor. subcampanulata, quadri-quinquefida: faux nuda aut barbata.

Stam. imo tubo inserta.

Styli primum connati, demum secedentes, stigmatibus obtusis.

Caps. unilocularis, apice bivalvis. Sem. non marginata.

Sturm. fl. germ. *Bunge* in act. soc. nat. mosqu. 7. t. 8—11.

1100. SWERTIA L. (1737.)

Cal. quadri-quinquepartitus.*Cor.* rotata, laciniis basi biglandulosis; aut campanulata, basi tuberculata, vel calcarata.*Styli* et *Caps.* Gentianae. Sem. margine alata.*Schk.* t. 58. *Gärtn.* t. 114. *Reichenb.* ic. 3. t. 290. vol. 5. t. 419.

1101. ROCHEFORTIA Sw. (1788.)

Cal. quinquepartitus.*Cor.* infundibuliformis, limbo quinquelobo.*Stam.* fauci inserta.*Styli* subulati pilosi.*Caps.* bilocularis, seminibus angulosis.*Swartz* fl. iud. occ. 1. t. 11.

C. CONVOLVULEAE.

1102. EVOLVULUS L. (1747.)

Cal. quinquepartitus.*Cor.* subrotata, quinquefida.*Styli* bipartiti. *Stigmata* incrassata.*Caps.* subbilocularis, loculis dispermis.*Bot. reg.* 401. *Humb.* nov. gen. 3. t. 215. *Labill.* nov. Calcd. t. 29. [*Cladostyles* Broupl. in *Humb.* plant. aequin. 1. t. 57.]

1103. DUFOUREA Kunth. (1818.)

Cal. quinquepartitus, laciniis duabus exterioribus maximis involucrentibus.*Cor.* campanulata, plicata.*Stigmata* globosa.*Caps.* bilocularis, loculis submonospermis.*Humb.* nov. gen. 3. t. 214.

1104. NEUROPELTIS Wallich. (1826.)

Cal. pentasepalus persistens; bracteis orbiculatis, membranaceis, reticulato-venosis suffultus.*Cor.* infundibuliformis quinquefida.*Stigmata* reniformia.*Caps.* monosperma, bractea dilatata suffulta.

1105. CRESSA L. (1747.)

Cal. quinquepartitus.*Cor.* infundibuli-hypocrateriformis.*Stam.* exserta.*Stigmata* capitata.*Caps.* bivalvis, monosperma.*Lam.* ill. t. 183.

1106. DICHONDRA Forst. (1775.)

Cal. quinquepartitus.*Cor.* subrotata.

Styli basilares. *Stigmata* capitata:

Utriculi gemini monospermi.

Forst. char. t. 20. *Smith.* ic. ined. t. 8. *Willd.* hort. ber. t. 81. [*Steripha Banks.* *Gärtn.* t. 94.]

1107. *CARMONEA* Cav. (1799.)

Cal. quinquepartitus, laciniis linearibus.

Cor. rotata, quinquepartita.

Stam. filamenta basi tubi inserta. *Antherae* didymae.

Styli capillares.

Drupa globosa, nuce hexasperma.

Anal. de cienc. 1. t. 3.

1108. *FALKIA* L. fil. (1781.)

Cal. quinquedentatus.

Cor. infundibuliformis, plicata.

Stam. corolla breviora.

Caryopses quatuor.

Andr. repos. t. 257.

1109. *CUSCUTA* Trag. (1551.)

Cal. quadri - quinquelobus.

Cor. subglobosa, persistens, quadri - quinquefida, basi intus squamulis obsessa.

Stamina tubo corollae accreta.

Caps. bilocularis, basi dehiscens, loculis dispermia. *Embryo* spiralis periphericus acotyledon.

Reichenb. ic. 5. t. 497 — 500. *Hook.* exot. fl. 2. t. 140.

D. H Y D R O L E A E.

110. *HYDROLEA* L. (1757.)

Cal. quinquepartitus.

Cor. subcampanulata.

Stam. inclusa, basi dilatata. *Antherae* sagittatae.

Stigmata truncata.

Caps. bi - trilocularis, polysperma: placentae e dissepimento.

Hydroleae. *Aubl.* guian. t. 110. *Bot. reg.* 566. *Gärtn.* t. 55.

111. *WIGANDIA* Kunth. (1818.)

Cal. quinquepartitus, persistens.

Cor. infundibuliformis, limbo quinquepartito.

Stam. exserta. *Antherae* sagittatae.

Stigmata subpeltato - depressa.

Caps. bilocularis. Placentae binae in quovis loculo laminiformes ad dissepimentum affixae.

Hydroleae. [*Hydrolea* Ruiz et Pav. fl. peruv. 3. t. 243 244. f. a.]

112. *NAMA* L. (1747.)

Cal. quinquepartitus, persistens.

Cor. rotato - infundibuliformis, limbo quinquefido.

Stam. subinclusa. Antherae reniformi-bilobae.

Stigmata obtusa.

Caps. bivalvis, subbilocularis. Dissepimentum medio interruptum, ibique placentiferum. Placentae laminiformes, valvis parallelae, demum liberae.

Hydroleae. Linn. fl. zeyl. t. 2. P. Brown. jam. t. 18. f. 2. Humb. nov. gen. 3. t. 218.

1113. *LONCHOSTOMA* Wikström. (1818.)

Cal. bracteatus, imbricatus, quinquepartitus.

Cor. tubulosa, limbo quinquepartito, laciniis lanceolatis, ad faucem attenuatis.

Stam. tubo inserta.

Caps. bilocularis, dissepimento utrinque placentifero, loculis subdecaspermis.

— — Wikstr. in act. acad. holm. 1818. t. 10.

II. COROLLA PENTAPETALA.

A. *Fructus superi.*

1114. *PICRAMNIA* Swartz. (1788.)

Flores dioeci.

Cal. tri-quinquepartitus.

Petala tria aut quinque, oblonga.

Stam. exserta, terna aut quina.

Styl. nulli. *Stigmata* duo.

Drupa nucleo biloculari, dispermo.

Terebintheae? Sw. fl. ind. occ. t. 4.

1115. *MENISCOSTA* Blum. (1825.)

Flores polygami.

Cal. quadri-quinquefidus.

Petala quatuor aut quinque, biserialia.

Stam. lata, petalis opposita.

Discus quinqueidentatus perigynus.

Stigmata duo.

Bacca didyma monosperma.

Menispermeis affinis.

1116. *HEUCHERA* L. (1737.)

Cal. ventricosus, quinqueidentatus.

Petala calyci inter dentes imposita.

Stamina calyci inserta. Antherae sagittatae.

Caps. semibilocularis: margines valvarum inflexi placentiferi.

Saxifrageae. Schk. t. 58. Gärtn. t. 36.

1117. *VELEZIA* Löffl. L. (1757.)

Cal. tubulosus, gracilis, quinqueidentatus.

Pet. parva, emarginata, unguibus filiformibus.

Caps. unilocularis, polysperma. Placenta filiformis.

Caryophylleae. Gärtn. t. 129.

B. *Fructus inferi.*1118. *PANAX* L. (1764.)

Flores umbellati polygami.

Involucrum polyphyllum.*Cal.* minimus, persistens.*Pet.* oblonga, revoluta.*Bacca* scrotiformis, disperma.*Araliaceae.* Jacqu. ic. 3. t. 634. Bot. mag. 1333.1119. *CUSSONIA* Thunb. (1780.)*Cal.* quinquedentatus.*Petala* trigona acuta.*Bacca* disperma.*Araliaceae.* Thunb. in nov. act. upsal. 3. t. 12. 13. Lam. ill. t. 187.1120. *VAHLIA* Thunb. (1782.)*Cal.* pentasepalus, patens.*Pet.* ovata, concava, patentia, calyci inserta.*Stam.* cum petalis alternantia. Antherae biloculares.*Caps.* unilocularis, bivalvis, polysperma.*Onagraceae.* Thunb. diss. bot. t. 2. Lam. ill. t. 183. [Rus-selia L. fil. Bistella Delil. in Caillaud it. Meroë 2. t. 63. f. 2.]1121. *LINCONIA* L. (1771.)*Cal.* quinquedentatus.*Petala* cum dentibus calycis alternantia.*Stam.* calyci inserta.*Cocci* duo membranacei, monospermi.*Brunieis* affinis. Swartz in Berl. mag. 4. t. 4.1122. *XANTHOSIA* Rudg. (1811.)*Cal.* quinquepartitus.*Petala* quinque.*Fructus* bipartibilis.*Umbelliferae.* Rudg. in Linn. transact. 10. t. 22.1123. *PHYLLIS* L. (1737.)*Cal.* minimus bifidus.*Petala* obtusa, revoluta.*Stam.* filamenta capillaria. Antherae oblongae.*Stylus* nullus. Stigmata duo reflexa.*Caps.* bilocularis, disperma.*Rubiaceae.* Dill. Eltham. t. 299. Gärtn. t. 25.1124. *DULONGIA* Kunth. (1825.)

Flores e nervo foliorum medio.

Cal. quinquedentatus.*Discus* orbicularis germen obtegens.*Petala* cum staminibus alternantia, sub ambitu disci cum iisdem inserta.

Antherae biloculares.

Stigmata duo.

Bacca semibilocularis, calyce petalis staminibusque coronata, loculis bi-trispermis.

— — Humb. nov. gen. 7. t. 623.

UMBELLIFERAE.

Familia satis naturalis. Araliis finitima, climata amans temperata, quin et arcticam incolens zonam: intra tropicos rara, ubi altissima montium iuga praefert. Herbae utplurimum, raro frutices. Folia plerumque composita, decomposita, ternatim secta. Si simplicia sunt, ut in Bupleuro, medias sistunt formas inter petiolum dilatatum et folia, cuius manifestum sistit exemplum Bolax. Foliorum compositio decrescit, quo propiora sunt floribus, ut tandem in involucri foliola abeant, Cari exemplo et Selinorum. Umbella plerumque adest, abit in capitula in Sanicula, Dondia et Eryngio: depauperata est, ut paucos exhibeat radios in Hydrocotyle, ut demum in pedunculos unifloros Bolaxis transeat. Involucrum universale et partiale; variis: umbellas excedens in Eriocalia et Bupleuris nonnullis, multo minus in plerisque, deest demum in Pimpinella.

Calyx semper superus, quinquefidus s. quinquedentatus, persistens aut caducus. Corolla pentapetala, plerumque alba, saepe flava, nonnunquam virescens, raro rubra, unico Didisci exemplo coerulea. Petala cum laciniis calycis alternantia, plerumque obcordata, involuta, lacinia media in fassura, raro plana. Saepe radium externarum umbellarum formant.

Stamina cum petalis alterna, disco nectarifero inserta, ergo altius sedent germine. Styli semper duo, stigmata capitata.

Fructus genuinus inferus, calyce coronatus, a quo et tegmen accipit. Hinc achenium dicendus; sed nonnunquam corticatus est aut utriculatus. Fructus media parte axi verticali saepissime bipartito filiformi aut superne parum dilatato iunguntur. Forma est varia, solida, globosa, ovata, prismatica, saepe compressa, marginata. Costae aut iuga prominent, valleculis interiectis. Sub tegmine vittae transparent, ductus aut conceptacula longitudinalia valleculas occupantia, succos proprios resinosos oleosove continentia. Qua fructus cum axi cohaerent, raphe adest aut commissura, planiuscula aut excavata, vittis pariter interdum obsessa. Omnis vero fructus albumine constituitur, in apice gerens embryonem, radícula sursum spectante, coryledonibus bimis simplicibus deflexis.

a. *Formae desciscentes. Umbellae plerumque capitatae.*

1125. ARCTOPUS L. (1737.)

Flores polygami dioeci.

Involucrum masculae plantae pentaphyllum, feminae tetraphyllum, demum auctum.

Fructus ovati inclusi in involucrio turgido spinoso.

Burm. afr. t. 1. Bot. reg. 705.

126. *ERYNGIUM* Nicand. Colum.

Flores capitati. Receptaculum paleaceum.

Involucrum subspinosum.

Petala inflexa.

Fructus paleaceo-squamati, calyce coronati.

Schk. t. 59. Delaroché Eryngiorum historia. 1808.

127. *EXOACANTHA* Labill. (1791.)

Capitula aggregata, involucrio longissimo polyphylo spinoso involuta. Involucella dimidiata.

Flores omnes hermaphroditii aequales.

Calyx vix ullus.

Fructus ovatus, striatus, hinc convexus, inde planus.

Labill. ic. plant. syr. t. 2. Lam. ill. t. 190.

128. *ECHINOPHORA* Tournef. (1718.)

Flores polygami, centralis sessilis femineus, germina intra pedicellos recipiens.

Involucra oligophylla.

Cal. quinquedentatus.

Pet. inaequalia.

Fructus ecostatus, striatus, raphe sulcata, in receptaculo turgido nidulans.

Lam. ill. t. 190. Engl. bot. 2413.

129. *ERIOCALIA* Sm. (1804.)

Umbella capitata, receptaculo villosa. Involucrum amplissimum lanuginosum.

Flores masculi et feminei mixti.

Cal. quinquefidus.

Cor. nulla.

Fructus ovatus, villosus quinquestriatus calyce coronatus.

Smith. exot. bot. 2. t. 78. 79. [Actinotus Labill. nov. holl. t. 92. Bot. reg. 654.]

130. *SANICULA* Brunsf. (1533.)

Umbella capitata. Involucra oligophylla.

Flosculi polygami.

Cal. persistens.

Fructus oblongus, solidus, ecostatus, setis uncinatis a. glochidatis obsessus.

Schk. t. 60. Gaud. fl. helv. 2. t. 3.

131. *DONDIA* * (1813.)

Umbella capitata. Involucrum hexaphyllum, umbella longius.

Cal. persistens.

Fructus ovatus, solidus, subtricostatus, valleculis convexis.

Sturm. fl. germ. Gaudin l. c.

1132. *ASTRANTIA* Moris. (1672.)

Umbellae fasciculiformes. Involucra umbellas aequantia subcolorata.

Cal. persistens.

Fructus oblongus, utriculis sulcatis rugosis cinctus.

Sturm. fl. germ. *Schk.* t. 60. *Gaudin* l. c.

1133. *POZOA* Lag. (1816.)

Umbella subglobosa. Involucrum polyphyllum umbellam aequans.

Fructus tetragono-prismaticus, dentibus calycis coronatus, striatus, costis binis lateralibus alatis.

[*Asteriscium Chamiss.* in *Linnaea* 1. tab. 5. f. 1.]

b. *Hydrocotylinae. Umbellae imperfectae. Involucra obsoleta. Folia simpliciuscula.*1134. *HYDROCOTYLE* Tournef. (1718.)

Umbella depauperata.

Petala plana.

Fructus compressus, orbiculatus, dorso tricostato, raphe angusta, cortice saepe reticulato.

Schk. t. 59. *Gärtn.* t. 22. *Gaudin* t. 3. *Ach. Richard* in *annual. bruxell.* 1. t. 1—18.

1135. *SPANANTHE* Jacqu. (1789.)

Umbella simplex, pauciradiata. Involucrum oligophyllum.

Fructus ovatus, solidus, glaber, dorso quinquecostatus, raphe lateribusque contractis.

Jacqu. ic. rar. 2. t. 350. [*Micropleura* *Lag.* in *oeis de Espan.* emigr. 4. p. 347. *Huanaca Cav.* ic. 6. t. 528.]

1136. *TRACHYMENE* Rudg. (1811.)

Umbella simpliciuscula. Involucrum polyphyllum.

Fructus ovatus, solidus, muriculato-rugulosus, dorso costatus, raphe planiuscula.

Rudg. in *Linn. transact.* 10. t. 21. [*Azorella* *Cav.* ic. 5. t. 485. *Labill.* nov. holl. t. 99—104.]

1137. *DIDISCUS* Cand. Hook. (1828.)

Umbella simplex. Involucrum polyphyllum.

Flores polygami (coerulei).

Petala plana.

Fructus compressus, orbiculatus, asper, bivittatus.

Hook. in *bot. mag.* 2875.

1138. *BOLAX* Commers. Juss. (1789.)

Umbella simplex aut pedunculi uniflora. Involucrum obsoletum.

Fructus ovatus solidus, tricostatus, valleculis lateribusque convexis.

[*Azorella Lam.* ill. t. 189. *Cav.* ic. 5. t. 484. *Chamitis Gärtn.* t. 22. *Selinum Cav.* ic. 486. 487. *Mulinum Pers.* *Fragosa Ruiz et Pav.* 3. t. 249. 250. *Humb.* nov. gen. 5. t. 424. *Pectophytum Kunth.* ib. t. 425.]

1139. *DRUSA* Cand. (1807.)

Pedunculi aggregati.

Cal. indivisus.*Fructus* lateribus compressis, raphe dorsoque angustissimis. Dorsum alatum, sinuato-dentatum, dentibus glochidatis.*Cand.* in ann. mus. 10. t. 38.1140. *BOWLESIA* Ruiz. et Pav. (1800.)

Umbella pauciradiata, saepe pedunculi uni-biflori.

Fructus subgloboso-ovatus, solidus, obsolete costatus pilosiusculus, raphe excavata.*Ruiz et Pav.* fl. peruv. 3. t. 251. Spec. umbellif. t. 5. f. 10.11. *Link et Otto* ic. nov. t. 4.c. *Bupleurinae. Umbellae perfectiores. Involucra latiuscula. Folia simplicia.*1141. *BUPLEURUM* Nicand.

Flores hermaphroditi, (plerumque flavi).

Cal. obsoletus.*Fructus* ovatus, prismaticus, subpentagonus (glaber aut muricatus).*Sturm.* fl. germ. *Reichenb.* ic. 2. t. 164 — 168. *Gärtn.* t. 22.[*Tenoria et Odontites* * *Diaphyllum et Isophyllum Hoffm.*]1142. *HERMAS* Thunb. L. fil. (1781.)

Flores polygami. Invol. universale breve, parziale umbellas aequans.

Fructus ovalis subcompressus, ecostatus, utriculatus.

(Caulis lignosus. Folia subtus tomentosa.)

Thunb. in nov. act. petropol. 14. t. 11. 12. *Gärtn.* t. 85.[*Lichtensteinia Schlechtend.*]d. *Pimpinelleae. Umbellae perfectae. Involucra nulla aut obsoleta. Fructus ovati solidi quinquecostati. Folia composita.*1143. *PIMPINELLA* Brunf. (1532.)

Involucra neutra.

Fructus ovatus solidus, costis quinque obtusiusculis, val-
leculis planiusculis vittatis.*Schk.* t. 78. 80. (715 — 720.) [*Trinia Hoffm. Reichenb.* ic. 5.
t. 442.]1144. *TRAGIUM* Diosc. Column. (1592.)

Involucra neutra.

Fructus ovato-oblongus ecostatus, pubescens, raphe planiuscula.*Column.* phytobas. t. 17. [*Pimpinella Loisel.* in diar. bot.
gallic. 2. t. 10. *Broter.* phytogr. lusit. t. 35. *Ledeburia Link.*]1145. *SESELI* Theophr.

Involucrum universale nullum, parziale oligo - aut polyphyllum.

Fructus ovatus solidus acute costatus.

(Rami virgati.)

Schk. t. 75. 80. (697.)

1146. **SISON** Diosc.

Involucrum nullum aut oligophyllum.

Fructus ovatus solidus obsolete tricostatus, valleculis convexis, lateribus contractis, raphe excavata.

Schk. t. 69. 80. (672.) [*Aegopodium* L. *Schk.* t. 79. *Thaspium* Nuttall. *Zizia* Koch.]

1147. **CARUM** Gul. Turn. (1551.)

Invol. universale monophyllum, partiale nullum.

Fructus prismaticus, dorso tricostatus, valleculis sulcatis striatis, raphe lata utrinque costata.

Schk. t. 77.

1148. **CNIDIUM** Cusson. (1784.)

Involucrum monophyllum.

Fructus ovatus solidus, costis quinque acutis subalatis, valleculis sulcatis striatis.

[*Peucedanum* *Schk.* t. 80. N. 638. *Silaus* Bess. *Gaud.* l. c. t. 7. *Ligusticum* Kunth. in *Humb.* nov. gen. 5. t. 422. ? *Palumbia* Bess.]

1149. **OENANTHE** Diosc.

Involucrum universale subnullum, partialia polyphylla.

Flosculi saepe radiantes polygami.

Fructus subprismaticus, angulato-costatus, calyce styloque coronatus.

Schk. t. 70. 80. (676.) *Broter.* phytogr. lusit. t. 33. *Gärtn.* t. 22. [*Rhelandrium* L. *Schk.* t. 71. *Ottosia* Kunth. in *Humb.* nov. gen. 5. t. 423. *Annesorhiza* Schlechtend. in *Linnaea* 1. t. 5. f. 3. 4.]

1150. **APIUM** Brunf. (1532.)

Involucrum neutrum.

Fructus ovatus solidus, obtuse costatus.

Schk. t. 78. 80. (722.) *Gärtn.* t. 22. *Gaudin* l. c. t. 8.

1151. **MEUM** Diosc. Tournef. (1718.)

Involucrum universale nullum, partiale oligophyllum.

Fructus prismaticus, acute angulatus.

Schk. t. 80. (681. 712.) [*Foeniculum* *Gaud.* t. 6. *Ptychotis* Koch. *Gaud.* t. 8.]

e. *Smyrnieae*. *Umbellae perfectae*. *Involucrum* universale nullum aut varium. *Fructus* solidus corticatus aut utriculatus.

1152. **SMYRNIUM** Nicand. Diosc.

Involucrum subnullum.

Fructus ovatus solidus, cortice nigro tricostato, raphe lateribusque contractis.

Schk. t. 76. *Moris.* hist. sect. 9. t. 4. *Kitab.* pl. hungar. 1. t. 23. *Gärtn.* t. 22. [*Eulophus* Nutt. *Cand.* umbell. t. 2. f. M.]

1153. *CACHRYs* Lobel. (1576.)

Involucra varia.

Fructus cortice suberoso fungoso crasso cinctus. Albumen involutum.

Reichenb. ic. 5. t. 489. *Moris.* umbellif. t. 3. sect. 9. t. 1.
 [Conium *Desfont.* atlant. 1. t. 65. ann. mus. 11. t. 29.
Crithmum maritimum L., *Rumia* et *Kruberia* *Hoffm.*
Ulopermum et *Hippomarathrum* *Link.* *Prangos pabularia*
Lindl. *Lag.* in *Ocios de espan.* emigr. 4. p. 347.]

1154. *CORIANDRUM* Plin.

Involucra nulla.

Flores radiantés.

Fructus globosus, subcostatus: raphe membrana obducta vittata. Albumen semilunare.*Schk.* t. 72. *Gärtn.* t. 22. *Engl. bot.* 67.1155. *BIFORIS* * (1820. *Bifora* M. B.)

Involucrum monophyllum foliaceum.

Fructus globosus, gibberosus, poris duobus in apice raphes.[*Coriandrum* *Moris.* sect. 9. t. 11.]1156. *SILER* *Gärtn.* (1788.)

Involucrum nullum.

Fructus compressus ovalis, costis novem obtusis, quinque prominulis.*Gärtn.* t. 22. [Agasyllis *Prodr. umb.*]1157. *CICUTA* Plin. Bessler. (1613.)

Involucrum subnullum.

Fructus ovatus solidus quinquecostatus, cortice in valleculis prominulo.*Schk.* t. 71. 80. (679.) *Gaudin.* t. 7.1158. *AETHUSA* L. (*Ethusa* 1737.)

Involucrum universale nullum, partialia dimidiata.

Fructus ovatus solidus, cortice in costas quinque obtusas turgescente, valleculis vittatis.*Schk.* t. 72. *Gaudin.* t. 7.1159. *PHYSOSPERMUM* Cusson. (1784.)

Involucrum oligophyllum.

Fructus obtuse quinquecostatus, utriculo laxo globoso vix striato vestitus.

Spec. umbell. t. 4. f. 8. [*Danaa* *Allion.* fl. pedem. t. 63.
Hänselera *Lag.*]

1160. *PLEVROSPERMUM* Hoffm. (1815.)

Involucrum utrumque polyphyllum.

Fructus ovato-oblongus, duplici tegmine membranaceo laxiusculo quinqueangulari, filis transversis cohaerente.*Hoffm.* tab. tit. umbell.161. *HASSELQUISTIA* L. (1755.)

Involucra varia. Flores radiantés.

Fructus margine compressus, planus, subrotundus: cortice ambitu turgido, dorso costas quinque obtusas producente. In media umbella fructus diffformes naviculares, margine laceri, dorso tricostrati.

Gärtn. t. 21. [*Tordylium* Lam. ill. t. 193. f. 2.]

1162. *TORDYLIUM* Diosc.

Involucra varia. Flores subradianes.

Fructus planus orbicularis, margine turgido crenulato.

Lam. ill. t. 193. f. 1. *Gärtn.* t. 21. *Schk.* t. 80. (620.)

[*Condylocarpus Hoffm.*]

1163. *THYSSELINUM* Plin. Lobel. Rivin. (1699.)

Involucra polyphylla.

Fructus planus, cortice dorsi costas tres obtusas producente, raphe plana.

Gaudin t. 5. [*Selinum Schk.* t. 63. *Callisace Fisch.*]

f. *Caucalinae.* *Fructus* armatus vel hirtus. Involucra varia.

1164. *CAUCALIS* Theophr.

Involucrum subnullum: involucella indivisa.

Flores subradianes.

Fructus oblongus, costis quinque aculeatis, angulis interpositis setosis. Albumen convolutum.

Schk. t. 61. 80. (625 — 627.) *Gärtn.* t. 20. *Gaudin.* t. 3. 4.

[*Orlaya*, *Turgenia* et *Platyspermum Hoffm.*]

1165. *DAUCUS* Galen. Lob. Clus. (1570. 1601.)

Involucri foliola pinnatifida.

Flores subradianes.

Fructus oblongus, costis quinque primariis glochidato-aculeatis, angulis interpositis setosis.

Schk. t. 61. *Gärtn.* t. 20.

1166. *TORILIS* *Gärtn.* (1788.)

Involucrum obsoletum.

Flores aequales.

Fructus ovatus subecostatus, undique aculeis setisque tectus.

Gärtn. t. 20. *Schk.* t. 79. (623. 624. 628.)

1167. *OLIVERIA* Venten. (1800.)

Involucri utriusque foliola tripartita.

Umbellae fasciculatae involucella aequantes.

Petala ad basin usque fissa.

Fructus ovatus hispidus, quinquestriatus.

Venten. hort. Cels. t. 21.

1168. *ATHAMANTA* L. (1737.)

Involucrum utrumque polyphyllum.

Fructus ovatus solidus hispidus s. villosus, quinquecostatus.

Schk. t. 79. (643.) [*Libanotis Gärtn.* t. 21.]

1169. *BUBON* L. (1737.)

Involucrum nullum.

Fructus ovatus solidus subcostatus pilosus.

[Seseli Smith. exot. bot. t. 120. Kit. pl. hung. 2. t. 146.
Bot. mag. 2073.]

1170. **BUNIAM** Diosc. * (1813.)

Involucra polyphylla.

Fructus ovatus, solidus quinquecostatus, muriculato-aculeolatus.

*Ammi Rivin. pentap. [Pituranthos Fisian. fl. lib. t. 7. f. 1.
Trachyspermum Link.]*

1171. **CAPNOPHYLLUM** Gärtn. (1791.)

Involucra varia.

Fructus ovalis, subcompressus, costis tribus muriculato-denticulatis, raphe plana vittata.

Gärtn. t. 85. [Conium Jacq. hort. vindob. t. 194. Rumia Link.]

g. *Scandicinae. Fructus pyramidati, rostrati.*

Involucrum universale nullum.

1172. **SCANDIX** Theoph.

Fructus pyramidatus, rostratus quinquecostatus, rostro nuclei longitudinem aequante vel superante.

*Gärtn. t. 85. Schk. t. 80. (684.) Venten. hort. Cels. t. 14.
[Wyllia Hoffm. umb. t. 2.]*

1173. **MYRRHIS** Theoph.

Fructus pyramidatus, quinquecostatus, stylis coronatus.

*Schk. t. 80. (683. 690—694.) Gärtn. t. 23. Rivin. pentap.
[Conopodium Koch.]*

1174. **CHAEROPHYLLUM** Colum. C. Bauh. (1596.)

Involucrum subnullum.

Fructus conicus ecostatus glaber, raphe sulcata.

Gärtn. t. 23. [Scandix Schk. t. 73.]

1175. **ANTHRISCUS** Plin. Pers. (1805.)

Involucrum subnullum.

Fructus conicus muricatus, stylis coronatus.

[Scandix Schk. t. 80. (686. 687.) Kit. plant. hung. 3. t. 224.]

h. *Amininae. Involucra varia. Fructus subovato-oblongi, prismatici costati sulcati.*

1176. **SCHULTZIA** * (1813.)

Involucra utriusque foliola bipinnata subcapillaria.

Fructus prismaticus quinquecostatus, stylis persistentibus coronatus.

[Sison crinitum. Pall. in act. petrop. 1779. P. 2. t. 7.]

1177. **AMMI** (Diosc.) Brunf. (1533.)

Involucra foliola pinnatifida.

Fructus oblongus obtuse costatus, valleculis planiusculis.

Schk. t. 61. Gärtn. t. 22. [Visnaga Gärtn. t. 21.]

1178. **CUMINUM** Theophr.

Involucrum utrumque oligophyllum.

Fructus ovato-prismaticus, glabriusculus, obtuse quinquecostatus, subutriculatus, vallecule vittatis.

Schk. t. 80. (714.)

1179. SIUM Diosc.

Involucrum utrumque foliolis integris.

Fructus subprismaticus, angulato-costatus.

Schk. t. 69. 80. (666. 670.) *Gärtn.* t. 23. [*Bunium* L. *Brignolia* Bertol. *Sebast.* fasc. pl. rom. 2. t. 2. *Critamus* Bess. *Koch.* *Drepanophyllum* Hoffm. *Helosciadium* Koch. *Reichenb.* ic. 3. t. 218.]

1180. CONIUM Theophr.

Involucrum utrumque polyphyllum.

Fructus ovatus solidus, costis quinque obtusis, immaturis crenulatis.

Schk. t. 62. *Gaudin* t. 7. *Hook.* exot. fl. 2. t. 152. [*Arracacia* *Bancroft.* *Pentacrypha* *Lehm.*]

1181. LIGUSTICUM Diosc.

Involucrum utrumque polyphyllum.

Fructus ovato-oblongus, costis quinque acutis, vallecule sulcatis.

Schk. t. 68. [*Molopospermum* Koch. *Levisticum* Gaud. t. 7. *Trochiscanthes* Koch. *Gaud.* l. c. *Gaya* Gaud. t. 6]

i. *Selineae.* *Involucra* varia. *Fructus* compressi plani, marginati, vel alati.

1182. SELINUM (Theophr.) L. (1737.)

Involucrum utrumque polyphyllum.

Fructus planissimus ovalis, margine alatus, costis dorsalibus tribus subprominulis, vallecule rapheque vittatis.

Schk. t. 79. (639.) *Gaudin* t. 5. [*Cervaria* *Gärtn.* t. 21. *Gaudin.* t. 4. *Melanoselinum* Koch. in nov. act. nat. cur. 12. t. 9. f. 6—8.]

1183. PEUCEDANUM Theophr.

Involucrum oligophyllum aut nullum.

Fructus planus marginatus (non alatus), costis dorsalibus (tribus aut quinque) obtusis, vallecule rapheque parallele vittatis.

Schk. t. 63. *Gärtn.* t. 21.

1184. HERACLEUM L. (1737.)

Involucrum subnullum.

Fructus compressus membranaceo-marginatus, costis dorsalibus tribus obtusis, vittis clavatis.

Schk. t. 67. [*Zosimia* Hoffm.]

1185. PASTINACA Plin. Dodon. (1563.)

Involucrum nullum.

Fructus ovalis compressus, marginatus, dorso tri-, raphe bistriatus.

Schk. t. 76. 80. (707.) [*Anethum* ib. t. 77. *Malabaila* Hoffm.]

186. *FERULA* Plin.

Involucra varia.

Flores polygami.

Fructus compressus, planus, margine incrassato, costis tribus dorsalibus obtusis.*Moris*, sect. 9. t. 15. *Lam.* ill. t. 205. *Desfont.* fl. atl. t. 67.*Kit.* pl. hung. 1. t. 60.187. *ANGELICA* Braunsch. (1500.)

Involucrum universale nullum.

Fructus subcompressus margine alatus, costis tribus dorsalibus acutis subulatis, valleculis sulcatis.*Lam.* ill. t. 198. *Gärtn.* t. 85. [*Selinum* *Schk.* t. 79. (636.)*Archangelica* *Gaudin.* t. 5. *Ostercicum* *Hoffm.* *Reichenb.*ic. 5. t. 402. *Conioselinum* *Fisch.* *Wimmer et Grabowski*fl. siles. 1. p. 266. *Mylinum* *Gaudin.* t. 5.]188. *IMPERATORIA* Matth. (1560.)

Involucrum universale nullum.

Fructus compressus planus, margine alatus, costis tribus dorsalibus obtusis, valleculis sulcatis.*Lam.* ill. t. 199. *Schk.* t. 71. *Gaudin.* t. 5. [*Angelica* mon-tana *ib.* *Peucedanum* *Chabraei* *ib.* t. 4.]189. *THAPSIA* Theophr.

Involucrum universale subnullum.

Fructus angustus subcompressus ecostatus, alis dorsalibus marginalibusque.*Lam.* ill. t. 206. *Gouan.* ill. t. 10. [*Laserpitium* *Desf.* fl. atlant. t. 68. 72.]190. *LASERPITIUM* Plin. Lobel. (1572.)

Involucrum utrumque polyphyllum.

Fructus ovalis subcompressus, costis primariis tribus acutis, secundariis et margine alatis.*Schk.* t. 67. 80. (651. 652.) *Gärtn.* t. 85.191. *ARTEDIA* L. (1736.)

Involucrum foliola pinnatifida.

Flores radiantes.

Fructus oblongus subcompressus, alis marginalibus sinuatis, striis dorsalibus quinque, raphe squamata.*Gärtn.* t. 85. *Lam.* ill. t. 139.* * *Flores imperfecti.*192. *GUMILLEA* Ruiz et Pav. (1800.)*Cal.* campanulatus quinquefidus.*Cor.* nulla.*Caps.* bilocularis, birostris, polysperma.— — *Ruiz et Pav.* fl. peruv. 3. t. 24.193. *CHAILETTIA* Cand. (1811.)

Petioli floriferi.

Cal. quinquepartitus.

Squamae cum staminibus alternantes.

Drupa bilocularis.

— — Ann. mus. 17. t. 1.

1194. *CELTIS* Tournef. (1718.)

Flores dioeci.

Cal. quinque - sexpartitus.

Styli nulli. Stigmata divaricata bifida.

Drupa baccata monosperma.

Amentaceae. Tournef. t. 383. Schk. t. 354. Gärtn. t. 77. [Mertensia Kunth. in Humb. nov. gen. t. 103.]

1195. *ULMUS* Colum. Virgil.

Cal. quadri - quinque - octolobus campanulatus.

Stamina quatuor, quinque; octo.

Samara.

Amentaceae. Schk. t. 57. Gärtn. t. 49.

C H E N O P O D E A E.

1196. *CHENOPODIUM* Tabernaem. (1613.)

Cal. quinquepartitus, demum haud mutatus.

Stam. receptaculo inserta.

Utriculus monospermus. Embryo periphericus.

Gärtn. t. 75. [Chenolea L. Cochliospermum Lag.]

1197. *EUCHYLAENA* R. Br. (1810.)

Cal. quinquepartitus, demum baccatus.

Stam. basi calycis inserta.

Fructus baccatus monospermus. Embryo periphericus.

[Suaeda Pall. illustr. t. 41. 43.]

1198. *ANABASIS* L. (1750.)

Cal. pentasepalus, tribracteatus, demum baccatus.

Fructus monospermus. Embryo periphericus.

Pall. illustr. t. 8. 9. Gärtn. t. 77.

1199. *SALSOLA* Caesalp. (1583.)

Cal. quinquepartitus, demum appendiculatus.

Stamina basi calycis inserta.

Embryo vel spiralis exalbuminosus, vel albuminosus curvatus.

Pall. illustr. t. 25. 30. Gärtn. t. 75.

1. *Kochia* Roth., calycibus demum alatis. Sturm. fl. germ.

2. *Sclerochlaena* R. Br. calycibus demum spinoscentibus. [Suaeda Pall. illustr. t. 35. Bassia Allion. misc. taurin. 3. t. 4. f. 2.]

1200. *HERNIARIA* Dodon. (1563.)

Cal. pentasepalus.

Stam. decem, quorum alterna sterilia.

Utriculus monospermus.

Schk. t. 56. Gärtn. t. 213.

201. MICROTEA Sw. (1788.)

Cal. quinquepartitus.*Stam.* quinque vel octo.*Styli* duo vel quatuor.*Achenium* echinatum.*Swartz fl. ind. occ. t. 12. [Ancistrocarpus Kunth. in Humb. nov. gen. 2. t. 122.]*

1202. ADENOGRAMMA Reichenb. (1827.)

Cor. calycina, pentapetala, caduca, petalis striatis.*Stam.* cum petalis alternantia, hypogyna.*Stigmata* duo.*Germen* trilobum glandulosum monospermum. Embryo inversus curvatus.*Reichenb. hort. 2. t. 109.*

T R I G Y N I A.

* *Flores perfecti.*I. *Fructus superi.*

A. COROLLA MONOPETALA.

1203. REICHELIA Schreb. (1789.)

Cal. quinquepartitus.*Cor.* calyce minor, campanulata, quinquefida.*Stam.* basi tubi corollae inserta. Antherae utrinque bifidae, incumbentes.*Styli* setacei. *Stigmata* capitata.*Caps.* trilocularis circumscissa polysperma. Placenta centralis trigona.*Convolvuleae. [Sagonea Aubl. guian. t. 111.]*

B. COROLLA PENTAPETALA.

1204. STACKHOUSIA Smith. (1798.)

Cal. quinquepartitus.*Petala* unguibus coalita.*Stam.* inaequalia, cum petalis alternantia.*Stigmata* acuta simplicia.*Caps.* tricoeca.*— — Labill. nov. holk. t. 104.*

1205. SALMASIA Schreb. (1789.)

Cal. quinquepartitus, persistens.*Pet.* hypogyna, cum laciniis calycis alternantia.*Stylus* nullus. *Stigmata* tria.*Caps.* trilocularis, trivalvis, polysperma, valvis medio septi- et placentiferis.*Cisteae. [Tachibota Aubl. guian. t. 112.]*

1206. *MERCURIA* Fisch. Schlechtend. (1826.)*Cal.* pentasepalus persistens.*Petala* unguiculata, integra.*Stam.* cum petalis alternantia. Antherae subglobosae biloculares.*Caps.* inflata, depresso-globosa, umbilicata, sulcata, subtrilocularis. Placenta centralis. Sem. pyriformia.*Carthophylleae.*1207. *DRYPIS* Theophr. Anguill. (1561.)*Cal.* tubulosus, quinquedentatus.*Petala* fauce bidentata, laminis bipartitis.*Caps.* circumscissa monosperma.*Schk.* t. 86. *Gärtn.* t. 123.1208. *XANTHOXYLON* Colden. (1743.)

Flores diœci.

Cal. brevis, tri-quinquepartitus.*Petala* longiora (raro nulla).*Stamina* hypogyna.*Styli* tres, quinque (interdum connati).*Cocci* bini, terni, quini, bivalves, monospermi.*Rutaceis* affinis. *Adr. Juss.* in mœm. mus. 12. t. 25. f. 38. *Jacqu. ecl.* 1. t. 70. *Hilar. pl. us. bras.* t. 37. [*Fagara* L., *Aubertia* Bory., *Tenorea* Rafin., *Langsdorfia* Leand., *Pohlana* Mart.]1209. *AILANTHUS* Desfont. (1786.)

Flores polygami.

Cal. quinquedentatus.*Pet.* patentia.*Stam.* in floribus masculis subduplicata.*Styli* connati, stigmatate peltato coronati.*Samaræ* tres aut quinque.*Rutaceis* et *Terebinthaceis* affinis. *Desfont.* in act. acad. paris. 1786. t. 3. *Roxb. corom.* 1. t. 23.1210. *CASSINE* Pluknet. (1700.)*Cal.* quinquesidus minimus.*Pet.* patentia.*Stam.* cum petalis alternantia.*Styl.* nulli. Stigmata tria.*Drupa* trilocularis trisperma.*Rhamneae.* *Lam. ill.* t. 340. *Schk.* t. 84. *Gärtn.* t. 92.1211. *STAPHYLEA* L. (1737.)*Cal.* quinquepartitus coloratus.*Stam.* receptaculo glanduloso inserta.*Styli* duo aut tres.*Caps.* binae aut ternae, membranaceae inflatae: seminibus quaternis; quinis aut senis osseis.*Rhamneae.* *Schk.* t. 84. *Gärtn.* t. 69. [*Bumalda* Thunb. e *Cand.* huc pertinet.]

1212. *PALIURUS* Theophr. Tournef. (1718.)*Cal.* quinquefidus.*Pet.* unguiculata.*Discus* carnosus perigynus.*Styli* breves.*Drupa* suberosa alata trilocularis trisperma.*Rhamneae.* Lam. ill. t. 210. Gärtn. t. 43. Bot. mag. 2535.1213. *SEMECARPUS* L. fil. (1781.)

Flores subdioeci.

Cal. quinquefidus.*Pet.* oblonga.*Discus* urceolatus perigynus.*Styli* recurvati.*Nux* reniformis, receptaculo carnosso insidens.*Terebinthaceae.* Roxb. corom. 1. t. 12. [*Anacardium*
Gärtn. t. 40.]1214. *RHUS* Theophr.

Flores subdiclini.

Cal. quinquefidus.*Petala* ovata patentia, aut nulla.*Styli* tres aut stigmata tria.*Bacca* monosperma.*Terebinthaceae.* Schk. t. 82. Ann. mus. 5. t. 30.
[*Lobadium* Rafin. in diar. phys. 89. t. 98. glandulis bi-
lobis hypogynis differt.]1215. *ASTRONIUM* Jacq. (1763.)

Flores dioeci.

Cal. pentasepalus.*Pet.* ovata.*Glandulae* quinque nectariferae.*Achanium* calyce acuto tectum.*Terebinthaceae.* Jacq. amer. t. 181.1216. *SPATHELIA* L. (1764.)*Cal.* quinquepartitus coloratus.*Pet.* oblonga.*Stam.* filamenta basi dilatata, utrinque dentata, villosa.

Antherae incumbentes, biloculares.

Styli nulli. Stigmata tria.*Drupa* trilocularis, alata, trisperma.*Terebinthaceae.* Bot. reg. 670. Gärtn. t. 58.1217. *TRICEROS* Lour. (1790.)*Cal.* quinquepartitus.*Pet.* oblonga.*Styli* basi distantes.*Bacca* coriacea tricornis, trilocularis.*Terebinthaceae.*1218. *LACEPDEA* Kunth. (1821.)*Cal.* pentasepalus, subimbricatus.

Pet. brevissime unguiculata, obovata.

Discus perigynus annularis decemlobus.

Stam. filamenta basi dilatata. Antherae erectae, biloculares.

Fructus baccatus, tricuspidatus, trilocularis, loculis subtrispermis.

Hippocrateae. *Humb.* nov. gen. 5. t. 444.

1219. *TURNERA* Plum. (1703.)

Cal. infundibuliformis quinquefidus.

Pet. patentia, cum staminibus calyci inserta.

Styli tres simplices, vel bipartiti. Stigmata multifida.

Caps. unilocularis, trivalvis, valvis placentiferis. Sem. arillo membranaceo unilaterali instructa.

Loasais affinis. *Aubl.* guian. t. 113. 114. *Plum.* ic. 150. *Gärtn.* t. 76. [*Piriqueta* *Aubl.* t. 117.]

1220. *SCHUMACHERIA* *

Cal. tubulosus, quinquedentatus, deciduus.

Pet. quinque receptaculo inserta.

Stam. hypogyna. Filamenta capillaria. Antherae oblongae, erectae.

Styli capillares.

Caps. linearis, subtridentata unilocularis, trivalvis. Sem. rugosa, valvarum marginibus affixa.

— — [*Wormsköldia* *Schumach.* guin. p. 165. *Raphanus pilosus* *W.*, *Cleome raphanoides* *Cand.*]

1221. *MALESHERBIA* Ruiz. et Pav. (1794.)

Cal. tubulosus, ventricosus, decemstriatus, quinquedentatus.

Pet. ori calycis inserta, cum dentibus alternantia, basi squamis subdenis denticulatis.

Stam. hypogyna, exserta. Antherae incumbentes.

Styli laterales, filiformes, stigmatibus capitatis.

Caps. unilocularis, trivalvis, polysperma, valvis placentiferis.

Malesherbiae *Don.* [*Gynopleura* *Cav.* ic. 4. t. 375. 376.]

1222. *PORTULACARIA* Jacqu. (1786.)

Cal. disepalus, coloratus.

Pet. hypogyna, obovata, persistentia.

Styl. 0. Stigmata muricata.

Fructus triqueter, alatus, monospermus.

Portulacaeae. *Jacqu.* coll. 1. t. 22. [*Häükea* *Salisb.*]

1223. *TELEPHIUM* Diosc.

Cal. quinquepartitus persistens.

Pet. integra.

Stam. receptaculo inserta.

Stigmata sessilia recurva.

Caps. trivalvis, basi trilocularis, polysperma.

Paronychiaae. *Schk.* t. 85.

224. CORRIGIOLA Euric. Cord. (1534.)*Cal.* pentasepalus, margine membranaceus.*Petala* integra, cum sepalis alternantia.*Stigmata* sessilia.*Nucula* monosperma.*Paronychiaea*. Schk. t. 85.II. *Fructus inferi.***225. HOLIGARNA** Buchan. Roxb. (1814.)*Flores* polygami.*Cal.* quinquepartitus.*Pet.* villosa, subconcreta.*Nux* monosperma.*Terebinthaceae*. Roxb. corom. 3. t. 282.**226. GRISELINIA** Forst. (1786.)*Flores* polygami.*Cal.* quinquedentatus.*Petala* obovata.*Stam.* cum petalis alternantia. *Antherae* biloculares.*Styli* tres lanceolati.*Fructus* monospermus?

— — [Scopolia Forst. char. 70.]

227. SAMBUCUS Plin.*Cal.* quinquedentatus.*Cor.* rotata.*Stigmata* tria.*Bacca* subtrisperma.*Caprifolia*. Schk. t. 83. Gärtn. t. 27.**228. VIBURNUM** Virgil.*Cal.* quinquedentatus.*Cor.* quinqueloba.*Stigmata* sessilia, tria.*Bacca* monosperma.*Caprifolia*. Sturm. fl. germ. Schk. t. 81. Gärtn. t. 27.**229. ZANONIA** L. (1764.)*Flores* dioeci.*Cal.* tripartitus.*Cor.* quinquepartita.*Stam.* filamenta plana, basi subconnata. *Antherae* uniloculares.*Styli* bifidi.*Fructus* carnosus, trilocularis, loculis dispermis. *Sem.* alata.*Cucurbitaceae*. Lam. ill. t. 816.**230. PASCHANTHUS** Burch. (1824.)*Flores* polygami.*Cal.* tubulosus, quinquefidus.

Pet. quinque linearia calyci inserta.

Stam. Antherae lineares, erectae.

Stigmata tria brevissima.

Bacca stipitata hexasperma. Sem. arillo carnoso ornata.

Passifloreae.

** *Flores imperfecti.*

1231. *PHARNACEUM* L. (1737.)

Cal. pentasepalus margine scariosus.

Stam. hypogyna, abortu saepe tria.

Styli tres.

Caps. trilocularis, trivalvis, polysperma.

Caryophylleae. Lam. ill. t. 214. Gärtn. t. 130. [Molugo *L.* floribus triandris differt.]

1232. *GINGINSIA* Cand. (1828.)

Cal. quinquepartitus margine scariosus.

Stam. imo calyci inserta.

Discus perigynus carnosus quinquelobus.

Stigmata sessilia cristaeformia.

Caps. unilocularis, trivalvis, polysperma. Placenta centralis.

Portulacaeae. Act. soc. hist. nat. paris. 4. t. 17-18.

1233. *BASSELLA* Rheed. (1688.)

Cal. corollinus quinque-septemfidus, coloratus, subconnivens, bracteatus.

Stam. calyci adnata, filamentis basi dilatatis, antheris bilocularibus.

Stigmata lateralialia.

Sem. unicum in calyce baecato.

Chenopodeae. Lam. ill. t. 215. Jacqu. ecl. 2. t. 164. Gärtn. t. 128.

1234. *BETA* Colum.

Cal. quinquepartitus.

Stigmata sessilia.

Sem. unicum in calyce durescente.

Chenopodeae. Schk. t. 56. Gärtn. t. 75.

1235. *BOSEA* L. (1737.)

Cal. quinquepartitus, imbricatus.

Stigmata sessilia.

Drupa monosperma.

Chenopodeae. Jacqu. ecl. t. 25. Gärtn. t. 77.

1236. *SECURINEGA* Commers. Juss. (1789.)

Flores dioeci.

Cal. quinquepartitus.

Stam. disco glanduloso inserta, cincta glandulis totidem.

Stigmata tria, subsessilia, biloba.

Caps. tricocca. Cocci bivalves, monospermi.

Tricoccae. Adr. Juss. Euphorb. t. 2. f. 4.

1237. *PORANTHERA* Rudg. (1811.)

Cal. pentasepalus, corollinus.

Stam. curvata. Antherae quadriloculares, apice glandulosae.

Caps. tres, polyspermae.

— — Rudg. in Linn. transact. 10. t. 22.

1238. *KERAUDRENIA* Gay. (1821.)

Cal. quinquepartitus persistens.

Stam. filamenta basi dilatata, conniventia. Antherae erectae biloculares, longitudinaliter dehiscentes.

Styli subulati, apice cohaerentes.

Caps. trilocularis, loculis dispermis.

Lasiopetaleae. Mém. mus. 7. t. 23.

T E T R A G Y N I A.

1239. *PARNASSIA* Tournef. (1718.)

Cal. quinquepartitus.

Pet. subrotundo-ovata.

Squamae glandulis stipitatis cinctae, petalis oppositae.

Stigmata quatuor sessilia.

Caps. unilocularis quadrivalvis, valvis medio septiferis.

Sem. membrana obvoluta.

Dioscoreae. Sturm. fl. germ.

P E N T A G Y N I A.

* *Flores perfecti.*

I. *Fructus superi.*

1240. *GALIPEA* Aubl. (1775.)

Cal. quinquefidus.

Pet. oblongo-linearis, conglutinata.

Stam. quinque, quorum duo, tria aut quatuor abortiunt.

Antherae basi bilobae.

Styli subcoacti.

Cocci quinque bivalves.

Cuspariae. Aubl. guian. t. 269. *Hilar.* plant. us. brasil. t. 12. 13. 14. *Cand.* mém. mus. 9. t. 10. *Adr. Juss.* mém. mus. 12. t. 23. f. 34. [*Raputia* Aubl. guian. t. 212. *Sciuris* Schreb. *Bonplandia Willd.* *Humb.* plant. aequin. 2. t. 57. *Richard.* act. instit. gall. 1811. P. 2. t. 10. *Bavia* Nees in act. acad. nat. cur. 11. t. 19. 23. 24. *Concho-carpus* Mikan del. brasil. t. 2. *Erythrochiton* Nees in act. acad. nat. cur. 11. t. 22. *Diglossia* ib. t. 25.]

1241. *PILOCARPUS* Vahl. (1796.)

Cal. quinquefidus.

Petala apice uncinata.

Stam. subulata, antheris subrotundis

Styli quinque laterales apice in stigma quinquelobum coaliti.

Cocci quinque coaliti, bivalves monospermi.

Rutaceae. Vahl. eclog. 1. t. 10. *Nees* in act. nat. cur. 11. t. 30. *Hilar.* fl. brasil. t. 17.

1242. *COMMERSIONIA* Forst. (1775.)

Cal. quinquepartitus.

Pet. basi saccata, apice longe mucronata.

Stam. tricuspidata: interpositis sterilibus petaloideis, stelatim patentibus.

Caps. tri-quinquevalvis, echinata, setis villosis. Cotyledones planae.

Büttneriae. Forst. char. 22. *Andr.* repos. t. 519.

1243. *SPONDYLOCOCCUS* Willd. Röm. et Schult. (1820.)

Cal. pentasepalus.

Pet. obtusa.

Stigmata ciliata.

Caps. quinquelocularis polysperma.

1244. *DROSERA* L. (1737.)

Cal. quinquepartitus persistens.

Stam. filamenta hypogyna, apice dilatata. Antherae loculis duobus distantibus.

Styli emarginati, bipartiti aut multifidi.

Caps. tri-quinquevalvis, polysperma valvis placentiferis.

Sem. nuda, membrana obvoluta.

Droserae. Schk. t. 87. *Gärtn.* t. 61. *Humb.* nov. gen. 5. t. 490. *Labill.* nov. holl. t. 105. 106.

1245. *ALDROVANDA* Caj. Monti. (1747.)

Cal. quinquepartitus persistens.

Styli indivisi capitati.

Caps. unilobularis, quinquevalvis, decasperma.

Droserae. Monti in comment. bonon. 2. P. 3. t. 12.

1246. *SIBBALDIA* L. (1737.)

Cal. campanulatus, semidecemfidus, laciniis alternis minoribus.

Pet. calycis ori inserta, cum laciniis maioribus alternantia.

Stam. filamenta abbreviata, calyci inserta. Antherae biloculares.

Styli sublaterales decidui.

Caryopses nudae quinae, denae aut plures.

Potentillae. Engl. bot. 897. *Sturm.* fl. germ. [Horkelia Schlechtend. Hook. bot. mag. 2880. staminum numero duplicato differt.]

1247. *STATICE* Plin. Dalech. (1586.)

Flores spicato - paniculati.

Cal. monophyllus, plicatus, scariosus.

Petala quinque subconnata.

Stam. basi petalorum inserta.

Utriculus monospermus calyce inclusus.

Plumbagineae. Reichenb. ic. 2. t. 162. 163. 172. 173.
186 — 188. vol. 3. 225. 226. 237. 291.

1248. *ARMERIA* Willd. (1809.)

Flores fasciculato - capitati.

Involucrum imbricatum, basi tubulo productum. Receptaculum paleaceum.

Cetera Statices.

Plumbagineae. [Statica *Armeria* Schk. t. 87.]

1249. *LINUM* Virg. Plin.

Cal. subpentasepalus.

Petala quinque cum staminibus annulo perigyno inserta: filamenta sterilia cum fertilibus alterna.

Caps. decemlocularis, loculis monospermis, valvis introflexis.

Lineae. Schk. t. 87. Sturm. fl. germ.

1250. *CRASSULA* Dillen. (1732.)

Cal. tubulosus, quinquepartitus.

Petala basi connexa, staminifera.

Squamae quinque perigynae nectariferae.

Caps. quinque acuminatae, introrsum dehiscentes, polyspermae.

Sedae. Andr. repos. 56. Bot. reg. 320.

1. *Globulea* Haw. Cand., petalis apice subglobuliferis. Cand. pl. succul. t. 61. 133.

2. *Cyrtogyns* Haw. Cand., ovariis gibberosis, stylis elongatis. Jacqu. hort. schönbr. t. 433.

3. *Grammanthes* Caud., squamis deficientibus. Herm. hort. lugduno-bat. p. 533.

4. *Rochea* Cand., petalis cohaerentibus, sed ita, ut facile secedant.

1251. *EURYCOMA* Jack. (1825.)

Flores polygami.

Cal. quinquepartitus.

Petala glanduloso - pilosa.

Glandulae decem geminae inter stamina.

Fructus quinque monospermi, basi coaliti.

Terebintheae.

1252. *GOUPIA* Aubl. (1775.)

Cal. quinquedentatus persistens.

Petala patentia, intus ligula acuta appendiculata.

Stam. disco glanduloso inserta.

Stigmata quinque.

Bacca pentasperma.

Rhamneae. Aubl. guian. t. 116. [Glossopetalum Schreb.]

CLASSIS VI. H E X A N D R I A.

CONSPECTUS

CLASSIS SEXTAE.

M O N O G Y N A E.

- I. *Floribus perfectis.*
 - A. *Corolla monopetala.*
 - a. *Fructibus superis* 1260 — 1266.
 - b. *Fructibus inferis* 1267 — 1277.
 - B. *Corolla pleiopetala.*
 - a. *Fructibus superis.*
 - α. *Corolla dipetala* 1278 — 1284.
 - β. *Corolla tripetala* 1285 — 1308.
 - γ. *Corolla tetrapetala* 1309 — 1312.
 - δ. *Corolla pentapetala* 1313 — 1314.
 - ε. *Corolla hexapetala* 1315 — 1319.
 - b. *Fructibus inferis* 1320 — 1325.
- II. *Floribus imperfectis.*
 - A. *Fructibus superis.*
 - a. *Perianthio duplici* 1326 — 1328.
 - b. *Perianthio simplici.*
 - α. *corollino monopetalo* 1329 — 1377.
 - β. *corollino pleiopetalo* 1378 — 1415.
 - B. *Fructibus inferis.*
 - a. *Spathaceae, spatha uni-multiflora.*
 - α. *Perianthium duplex* 1416 — 1420.
 - β. *Perianthium simplex* 1421 — 1438.
 - b. *Affines* 1437 — 1453.

D I G Y N A E.

1454 — 1460.

T R I G Y N A E.

1461 — 1517.

H E X A G Y N A E.

1518 — 1519.

P O L Y G Y N A E.

1520.

M O N O G Y N I A

I. *Flores perfecti.*

A. COROLLA MONOPETALA.

a. *Fructus superi.*

1260. *ASTROCARYON* Meyer. (1818.)

Flores monoeci, spatha simplici.

Flores masculi, rhachi alveolatae immersi.

Cal. tripartitus.

- Cor. campanulata tripartita.*
Flores feminei infra spicas masculas solitarii.
Cal. urceolatus, ore subtridentato.
Cor. campanulata, ore subtridentato, intus duplicata.
Stylus conicus: stigmata simplicia.
Drupa fibrosa, monosperma: putamine poris tribus verticalibus stellatis. Albumen cavum, aequabile. Embryo intra porum.
Palmae. Mart. palm. bras. t. 58—65.
1261. *MNASIUM* Schreb. (1789.)
Involucrum amplissimum, bifidum, basi cordatum, acuminatum. Flores numerosi congesti.
Cal. pedunculatus, bracteis imbricatis suffultus, tripartitus, laciniis concavis lanceolatis, acuminatis.
Cor. tripetala, petalis membranaceis obtusis.
Stam. filamenta brevissima. Antherae elongatae, lanceolato-lineares, tetragonae, rimis longitudinalibus dehiscentes, apice appendiculatae (appendicula ovata, concava, nigra).
Styl. filiformis, stigmatibus tribus spiratim convolutis.
Fructus capsularis orbicularis, trivalvis, trispermus.
Juncea. [Rapatea Aubl. guian. t. 118. Desvoux in ann. scient. nat. 13. t. 4. f. 2. Rudg. guian. t. 2. Spathanthus Desv. l. c. f. 1.]
1262. *TRIPHASIA* Lour. (1790.)
Cal. urceolatus, tridentatus.
Cor. campanulata, tripartita.
Stam. (quandoque quina, sexto abortiente). Filamenta disco hypogyno inserta, inferne applanata, subulata.
Antherae erectae, ovatae.
Stylus crassus, stigmate trigono.
Fructus aurantii, trilobularis, trispermus.
Aurantiaecae Corr. [Limonia Jacqu. ic. rar. 3. t. 463. Andr. bot. rep. 143.]
1263. *PRINOS* Gronov. (1743.)
Flores subdiclini, polygami.
Cal. semisexfidus, persistens.
Cor. rotata, sexpartita.
Stam. Filamenta subulata, perigyna, erecta.
Stylus brevis, stigmate obtuso.
Bacca hexapyrena.
Rhamnae. Duhamel. arbr. 1. t. 23.
264. *MABA* Forst. (1775.) Juss. (1804.)
Flores dioeci.
Cal. tripartitus.
Cor. tubulosa, trifida.
Stam. hypogyna (quandoque terna). Filamenta abbreviata, alterna gemina. Antherae erectae, oblongae.
Germen trilobulare, loculis dispermis.

Bacca ovalis, calyce cupulato suffulta.

Ebenaceae Juss. *Forst.* char. 61. *Labill.* nov. Caled. t. 36. [*Perreola Roxb.* corom. 1. t. 45.]

1265. *ACHRAS* P. Brown. (1756.)

Cal. hexasepalus, sepalis ovatis concavis, interioribus coloratis.

Cor. tubulosa, limbo erecto sexfido. *Squamas* intra faucem corollae, emarginatae.

Stam. Filamenta tubo corollae inserta, brevissima. *Antherae* cordato-sagittatae, inclusae, cum squamis alternantes.

Stylus subulatus, corolla longior. *Stigma* obtusum.

Fructus pomaceus, carnosus, decem - aut duodecimlocularis, loculis ternis plerumque seminiferis. *Sem.* solitaria, compressa, ovalia. *Embryo* evolutus. *Albumen* consumtum.

Sapoteae. P. Brown. *jam.* t. 49. f. 3. *Gärtn.* t. 104.

1266. *ANOPTERUS* Labill. (1804.)

Cal. sexfidus, persistens.

Cor. tubo brevissimo, limbo sexfido.

Stam. Filamenta subulata, basi corollae affixa. *Antherae* erectae, ovatae, biloculares.

Stylus brevis, apice bifidus.

Caps. oblonga, unilocularis, bivalvis. *Sem.* margini varum affixa, apice alata.

Gentianeae? *Labill.* nov. holl. 1. t. 112.

b. *Fructus inferi.*

1267. *AMAJOUA* Aubl. (1775.)

Cal. campanulatus, sexdentatus.

Cor. infundibuliformis, limbo sexfido.

Stam. inclusa. *Antherae* oblongae, erectae.

Stylus filiformis. *Stigma* incrassatum.

Bacca ovalis unilocularis, polysperma. *Placenta* centralis columnaris.

Rubiaceae. Desfont. in *Mém. du mus.* 6. t. 4. 5. *Humb.* nov. gen. 3. t. 294. *Aubl.* *guian.* t. 357. [*Melanopsidium?* *Coll. hort. ripul.* t. 35. *Ehrenbergia* * *syst.* n. 1318.]

1268. *ISERTIA* Schreb. (1789.)

Cal. coloratus, sexdentatus, persistens.

Cor. infundibuliformis, limbo sexfido.

Stam. Filamenta brevissima, intra faucem corollae. *Antherae* lineares, dorso adnatae, erectae.

Stylus filiformis, basi glandula cinctus. *Stigma* sexfidum.

Pomum subglobosum, sexloculare, loculis intus crustaceis.

Sem. plura, angulata.

Rubiaceae. [Guettarda *Aubl.* *guian.* t. 123.]

1269. *CASSUPA* Bonpl. (1808.)

Cal. integer, coloratus, subcarnosus.

Cor. tubulosa, superne verrucosa, interne sericea; limbo sexpartito.

Stam. exserta. Filamenta fauci affixa, brevissima, complanata. Antherae dorso affixae, oblongo-lineares, longitudinaliter dehiscentes.

Stylus stigmatе bifido.

Bacca sphaerica, bilocularis, oculis polyspermis.

Rubiaceae. Humb. plant. aequinoct. t. 12.

1270. *RICHARDIA* Houst. L. (1737.)

Cal. sex-octopartitus, laciniis lanceolatis aequalibus.

Cor. infundibuliformis, limbo sex-octofido.

Stam. Filamenta subulata, fauci inserta. Antherae incumbentes, versatiles, biloculares.

Stylus filiformis. Stigmata tria capitata.

Caps. tricocca, coccis monospermis.

Rubiaceae. Banks. reliq. Houst. t. 9. [*Richardsonia* Kunth. in Humb. nov. gen. 3. t. 279.]

1271. *STEVENSIA* Poit. (1804.)

Cal. integer, basi globulosus; limbo bifido, caduco.

Cor. tubulosa; limbo sex-septemfido, plano.

Stam. Antherae sessiles, biloculares, summo corollae tubo adnatae.

Styl. stigmatе bilamellato.

Caps. sphaerica, bivalvis; valvis introrsum divisis ut quadrilocularis videatur esse capsula. Placenta hemisphaerica valvis affixa. Sem. numerosa, minuta, compressa, sursum alata.

Rubiaceae. Poit. in ann. mus. 4. t. 60. Gärtn. t. 197.

1272. *COUTAREA* Aubl. (1775.)

Cal. bracteis duabus aristatis suffultus, turbinatus, limbo deciduo sexpartito; laciniis aristatis.

Cor. infundibuliformis, sursum sensim ampliata, incurva; limbo sexpartito, laciniis abbreviatis, recurvis.

Stam. exserta. Filamenta basi tubi inserta. Antherae oblongae, erectae.

Styl. filiformis, stamina aequans, stigmatе capitato.

Caps. coriaceo-chartacea, obovata, bilocularis, subquadri-
valvis, polysperma. Placenta libera, fungosa, scabra.

Sem. compressa, alata.

Rubiaceae. Aubl. guian. t. 122. Gärtn. t. 194. [*Portlandia* Jacq. amer. t. 182. f. 20.]

1273. *HILLIA* Jacq. (1763.)

Cal. hexasepalus; sepalis oblongis, erectis.

Cor. tubulosa, longissima, sexsulcata; limbo sexfido.

Stam. Filamenta brevissima, fauci corollae inserta. Antherae oblongae, erectae.

Styl. filiformis. Stigma capitatum.

Caps. compressa, bilocularis, siliquaeformis. *Sem.* minuta, numerosa, alata, ala apice in lacinias capillares abeunte.

Rubiaceae. *Jacqu.* amer. t. 66. *Swartz.* ind. occ. t. 11. *Gärtn.* t. 197. [*Fereiria Vandell.* in *Römer.* script. hisp. t. 6. f. 8.]

1274. *PSATHYRA* Commers. Juss. (1789.)

Cal. minimus, sexdentatus.

Cor. subcampanulata, sexloba, intus barbata.

Stam. intra faucem. Filamenta brevissima. *Antherae* ovatae, erectae.

Styl. capillaris. Stigma trilamellatum.

Bacca subglobosa, sulcata, hexapyrrena.

Rubiaceae. *Lam.* ill. t. 260. *Gärtn.* t. 194.

1275. *SCHRADERA* Vahl. (1796.)

Flores capitati. Involucrum coriaceum, tetra-hexasepalum.

Cal. campanulatus, subinteger.

Cor. campanulata, fauce hirsuta, limbo sexfido, (interdum quinque-septem-octofido).

Stam. inclusa, numero incerta.

Styl. incrassatus, stigmatate tri-quadripartito.

Bacca sexlocularis, polysperma.

Caprifolia. *Vahl.* eclog. t. 5.

1276. *CANARIA* L. (1771.) (*Canarina* Reich.)

Cal. sexfidus, lacinii lanceolatis, patulis, trinerviis, persistentibus.

Cor. campanulata, limbo sexfido, patente.

Stam. Filamenta basi dilatata, fornicata. *Antherae* erectae, biloculares, subconnatae.

Stylus simplex, stigmatate clavato.

Caps. hexagona, sexlocularis; loculis polyspermis. *Placentae* binae in quovis loculo, membranaceae.

Campanuleae. *Bot. mag.* 444. *Lam.* ill. t. 269. *Gärtn.* t. 211. [*Campanula* *L.* hort. *Cliffort.* t. 8.]

1277. *LEPTODERMIS* Wallich. (1825.)

Cal. sexdentatus, bractea membranacea, tubulosa, superne bifida inclusus.

Cor. sexfida.

Antherae mediis filamentis adnatae, lineares.

Stigma sexfidum.

Caps. hexasperma.

B. COROLLA PLEIOPETALA.

a. *Fructus superi.*

a. *Corolla dipetala.*

Gramineae.

1278. *BAMBUSA* Schreb. (1789.)

Spiculae paniculatae, distichae, multiflorae.

Flores inferiores hermaphroditi, superiores masculi.

Cal. valvae plures, infimae minutae, superiores valvae corollae inferiori similes.

Cor. bivalvis; valva inferior coriacea, superior membranacea, marginibus inflexis.

Squamae hypogynae binae.

Stylus bifidus.

Caryopsis valvis corollae involuta.

Roxb. corom. 1. t. 79. [*Nastus Pal. Beauv.* agrost. t. 25. f. 3.]

79. *GUADUA* Kunth. (1822.)

Spiculae fasciculatae, paniculatae, distichae, multiflorae.

Flores inferiores masculi vel neutri univalves.

Cal. bivalvis.

Cor. bivalvis: valva inferior concava, superior bicarinata.

Squamae hypogynae tres.

Stylus tripartitus.

Caryopsis valvis corollae involuta.

[*Bambusa Humb.* plant. aequin. 1. t. 20. 21.]

80. *NASTUS* Juss. (1789.)

Spiculae paniculatae, uniflorae.

Cal. imbricatus, valvis distichis.

Cor. bivalvis, valva infer. subtridentata. Rudimentum floris abortientis pedicellati.

Squamae hypogynae tres.

Stylus tripartitus.

[*Stenotaphrum Pal. Beauv.* agrost. t. 25. f. 5.]

81. *MELOCANNA* Trin. (1822.)

Spiculae paniculatae, multiflorae. Flores aequales, inferiores masculi aut neutri.

Cal. bivalvis.

Cor. bivalvis, valvis acuminatis.

Stylus tripartitus.

Pericarpium maximum, carnosum, acuminatum, monospermum.

[*Bambusa baccifera Roxb.* corom. 3. t. 243. *Besha Rheed.* malab. 5. t. 60.]

82. *ENRHARTA* Thunb. (1779.)

Spiculae paniculatae.

Cal. bivalvis, triflorus, valvis corolla minoribus.

Cor. lateralis neutra univalvis; altera bivalvis, valva superiore transversim rugosa, mutica aut aristata: cor. intermedia hermaphrodita bivalvis, valva infer. naviculari, quandoque mucronata, super. falcata.

Squamae hypogynae truncatae, dentatae.

Stam. quandoque quaterna.

Stylus bifidus.

Caryopsis bicornis.

Thunb. in act. holm. 1779: t. 8. *Smith.* ic. ined. t. 9 32. 33.
Swartz in *Lin.* transact. 6. t. 3. 4. *Palis. Beauv.* agrost.
 t. 12. f. 3. 4. [*Trochera Rich.* in diar. Rozer. 13. t. 3.
Tetradleuca R. Brown. *Ehrharta distichophylla Labill.*
 nov. holl. t. 117.]

1283. *GAHNIA* Forst. (1775.)

Spiculae paniculatae, uniflorae.
Cal. squamae undique imbricatae, numerosae.
Cor. bivalvis, valvis inaequalibus.
Stam. filamentis elongatis, persistentibus.
Stylus tridus. Stigmata bifida.
Nux ossea, nucleo transversim inciso.

Cyperaceae Forst. char. 26. *Labill.* nov. holl. t. 115.

1284. *LAMPOCARYA* R. Brown. (1810.)

Spiculae subracemosae uniflorae.
Cal. squamae numerosae, undique imbricatae.
Cor. sexvalvis, valvis aequalibus acuminatis.
Stam. filamenta elongata, persistentia.
Stylus trifidus; stigmata indivisa.
Nux ossea, basi styli persistentis cuspidata, nucleo laevi.
Cyperaceae. [*Gahnia trifida Labill.* nov. holl. t. 116.]

β. *Corolla tripetala.*

1285. *FLÖRKEA* Willd. (1801.)

Cal. tripartitus, persistens; laciniis lanceolatis.
Petala calyce minora, ipsius basi inserta, cum laciniis
 calycis alternantia.
Stam. minuta. Filamenta capillaria, basi calycis inserta,
 alternatim articulata. Glandulae ternae ad basin. An-
 theras subrotundae.
Styl. lateralis, stigmatibus duobus subrotundis.
Utriculi duo monospermi.

Portulacaceae? [*Nectris pinnata Pursh.*]

1286. *AMOORA* Roxb. (1819.)

Flores polygami.
Cal. tridentatus, bibracteatus.
Petala ovalia, concava.
Urceolus interne antheras oblongas, trigonas gerens.
Stylus nullus. Stigma dilatatum, trilobum, lobis quan-
 doque bifidis.
Caps. obovata, corticata, trivalvis, trilocularis, valvis
 medio septiferis. Placenta centralis. Sem. arillo caroso
 obvoluta. Cotyledones crassissimae.

Meliaceae. *Roxb. exorn.* 3. t. 258. [*Aphanamixis Blum.*]

1287. *FLOSCOPA* Lour. (1790.)

Cal. inferus, infundibuliformis, coloratus, persistens; limbo
 trifido patente.

Petala ovata, erecta.

Stam. Filamenta subulata, corolla longiora. Antherae bilobae, subrotundae.

Stylus subulatus, inflexus. Stigma crassiusculum.

Caps. biloba, bilocularis, monosperma.

1288. *GALANDRINIA* Kunth. (1823.)

Cal. disepalus, persistens.

Pét. tria (vel quinque) aequalia.

Stam. basi petalorum inserta (interdum quatuor) aut plura.

Filamenta basi dilatata. Antherae bilocularae.

Stylus simplex. Stigma tripartitum.

Caps. elliptica, calyce persistente oblecta, unilocularis, trivalvis, polysperma. Placenta centralis.

Portulacaceae. Humb. nov. gen. 6. t. 526. Bot. reg. 1194.

[*Talinum* Hook. exot. fl. 1. t. 82. *Cosmia* Domb. Juss.]

1289. *CANARIUM* Rumph. (1741.)

Flores dioeci aut polygami.

Cal. inferus, tridentatus.

Pét. concava, conniventia.

Discus urceolatus, sexcrenatus.

Stam. Filamenta subulata, basi subcoalita. Antherae erectae, oblongae, biloculares.

Stylus brevis. Stigmata tria.

Drupa nucleo triloculari, loculis duobus saepe abortientibus. Sem. bina in quovis loculo, pendula, exalbuminosa. Cotyledones tripartitae, contortae.

Terebintheae. Rumph. amboin. 2. t. 47. 49. 54. Gärtn. t. 102. Ann. of bot. 1. t. 7. f. 1. 2.

1290. *TRADESCANTIA* Rupp. (1718.)

Cal. inferus, trisepalus, sepalis ovatis, concavis, persistentibus.

Petala subrotunda, aequalia.

Stam. Filamenta villosa, villis articulatis. Antherae reniformes, biloculares, loculis discretis.

Stylus filiformis. Stigma capitatum, papillosum, vel trigonum, tubulosum.

Caps. trilocularis, trivalvis, valvis medio septigeris. Sem. in quovis loculo bina, valvarum parieti adhacrentia, albuminosa. Embryo dorsalis, in cavea exigua, e granulis duobus constans.

Commelineae. Schk. t. 88. [Ephemerum Tourn. t. 193. Cyanotis Don.]

1291. *CAMPEDIA* Rich. (1808.)

Cal. inferus, trisepalus, post anthesin incrassatus.

Pét. subrotunda, aequalia.

Stam. Filamenta villosa. Antherarum loculi discreti.

Stylus filiformis, stigmate trilobo.

Caps. baccata, trilocularis, calyce incrassato cincta.

Commelineae. Nov. act. nat. cur. vol. II. t. 1. f. 4.
[*Commelina Zanonia* L. *Redout.* lil. t. 192. *Zanonia Plum.*]

1292. *DICHORISANDRA* Mikan. (1820.)

Cal. inferus, trisepalus; sepalis concavis, intus coloratis, supremo latiori.

Pet. subrotunda, inferius angustius.

Stam. Filamenta basi petalorum inserta, nuda, brevissima. Antherae ovatae, erectae, tres superiores cum media inferiore contiguae: duae laterales inferiores divergentes.

Stylus columnaris, stigmatibus trigono.

Caps. trilocularis, polysperma.

Commelineae. Mikan del. flor. bras. 1. Nov. act. nat. cur. vol. II. t. 1. Bot. reg. 682. Hook. in Bot. mag. 2721.

1293. *CARTONEMA* R. Br. (1810.)

Cal. trisepalus, bracteatus; sepalis lanceolatis, acuminatis. *Pet.* obovata.

Stam. tria petalorum basi inserta, tria perigyna. Filamenta abbreviata, nuda. Antherae oblongae, biloculares.

Stylus conicus, stigmatibus capitato.

Caps. trilocularis, trivalvis; valvis medio septiferis; loculis dispermis.

Commelineae. Bauer ill. nov. Holl. t. 7.

1294. *DASYPOGON* R. Br. (1810.)

Flores capitati, paleis distincti.

Cal. tripartitus, pilis rigidis asperis cinctus: limbo subarticulato.

Pet. longe unguiculata, hyalina, laminis denticulatis.

Stam. Filamenta subulata, petalorum basi inserta. Antherae oblongae, incumbentes, biloculares.

Stylus conicus subulatus.

Utriculus monospermus.

Junceae? R. Brown ad Flinders it. t. 8.

1295. *ACANTHOSPORA* * (1817.)

Cal. inferus disepalus, sepalo altero maiore bifido.

Pet. convoluta, basi nuda.

Stam. basi perianthii inserta.

Stylus filiformis, stigmatibus tribus fimbriatis.

Caps. trilocularis. Placenta centralis triala. Sem. caudata.

Bromeliceae. [Bonaparte Ruiz et Pav. fl. peruv. 3 t. 262. 263.]

1296. *TILLANDSIA* L. (1737.)

Cal. inferus, bracteatus, tripartitus.

Pet. convoluta, basi nuda.

Stam. basi petalorum inserta. Antherae oblongae, erectae, biloculares.

Stylus simplex. Stigma trifidum, obtusum.

Caps. trilocularis, loculis polyspermis. Sem. papposa.

Bromeliace. Hook. exot. fl. t. 154. 173. 205. ? 218. Bot. reg. 105. 749. Bot. mag. 2841.

297. GUZMANNIA Ruiz et Pav. (1802.)

Cal. inferus, trifidus, convolutus.

Pet. convoluta, basi nuda, unguibus membranaceis.

Stam. receptaculo inserta. Antherae lineares biloculares.

Stylus triangularis, stigmatibus tribus acutis.

Caps. trilocularis, trivalvis, polysperma. Sem. acuminata.

Bromeliace. Ruiz et Pav. fl. peruv. 3. t. 261.

298. POURRETIA Ruiz et Pav. (1802.)

Cal. inferus, tripartitus.

Pet. convoluta, basi nuda.

Stam. petalorum basi inserta.

Stylus filiformis. Stigmata tria, linearia, convoluta.

Caps. trilocularis, polysperma. Sem. alata.

Bromeliace. Ruiz et Pav. fl. peruv. 3. t. 256. 257.

299. PITCARNIA Herit. (1788.)

Cal. semiinferus, trifidus.

Pet. ascendens, subunilabiata, basi squamis instructa.

Stam. basi petalorum inserta, declinata. Antherae lineares.

Stylus filiformis. Stigmata tria, linearia, convoluta.

Caps. trilocularis, polysperma. Sem. caudata.

Bromeliace. Herit. sert. angl. t. 11. Bot. reg. t. 1011. 1069. 1092.

300. LICUALA Rumph. (1741.)

Flores hermaphroditi. Spatha communis nulla, partiales vagae.

Cal. tripartitus.

Pet. tria, basi subcohaerentia.

Stam. basi in urceolum connata.

Stylus simplex. Stigmata tria.

Drupa monosperma. Albumen ruminatum. Embryo dorsalis.

Palmae. Rumph. amboin. 1. t. 9. Gärtn. t. 139.

301. CHAMAEDOREA Willd. (1808.)

Flores dioeci, spadici adnati, ebracteati. Spathae plures.

Cal. tripartitus.

Pet. in ♀ squamis praemunita.

Stylus trigonus.

Bacca monosperma. Albumen acquabile. Embryo lateralis.

Palmae. Mart. palm. brasil. t. 3. [Borassus Jacq. Schönbrunn. 2. t. 247. 248. Nunezharia Ruiz et Pav.]

1302. ACROCOMIA Mart. (1824.)

• Flores androgyni, foveis immersis.

Cal. trisepalus.

Pet. convoluta, prismatica.

In ♀ cupula annularis sexdentata perigyna.

Stylus brevis. Stigmata tria.

Drupa monosperma, putamine lenticulari, lateribus triporoso. Albumen aequabile. Embryo lateralis.

Palmae. Mart. palm. brasil. t. 56. 57. [Bactris minor Gärtn. t. 9.]

1303. KUNTHIA Humb. (1814.)

Flores monoeci, spadici adnati. Spathae plures.

Cal. tripartitus.

Pet. ovata.

Bacca globosa. Embryo basilaris.

Palmae. Humb. plant. aequinoct. 2. t. 122.

1304. ELATE Ait. (1789.)

Flores androgyni sessiles. Spatha duplex.

Cal. cupularis, tridentatus.

Cor. tripetala.

Stam. Antherae sessiles.

Styl. subulatus.

Stigmata tria.

Drupa putamine basi sulco exarato et triporo. Albumen aequabile. Embryo basilaris.

Rheed. malab. 3. t. 22—25.

1305. GEONOMA Willd. (1805.)

Flores monoeci, foveolis spadicis immersis. Spatha duplex.

Cal. trisepalus.

Pet. in ♀ coalita.

Urceolus sexdentatus, dentibus singulis antheras didymas gerentibus.

Stylus lateralis, vel basilaris. Stigmata revoluta.

Bacca monosperma. Albumen aequabile. Embryo basilaris vel lateralis.

Palmae. Mart. palm. brasil. t. 4—20. [Gynestum Poir. in mém. du mus. 5. t. 16—20.]

1306. MAURITIA L. fil. (1781.)

Flores dioeci, polygami, in amentis teretibus. Spadix spathis pluribus articulato-vaginatus.

Cal. campanulatus vel urceolatus, trifidus vel tridentatus.

Pet. in hermaphroditis coalita.

Stam. antheris linearibus erectis.

Stigma sessile, trilobum.

Bacca tessellato-corticata, monosperma. Albumen aequabile. Embryo lateralis.

Palmae. Mart. palm. bras. t. 38—44.

1307. *TALIERA* Mart. (1824.)

Flores hermaphroditi, spadici terminali ramosissimo adnati. Spathae vagae.

Cal. trifidus.

Stam. in cupulam hypogynam connata.

Styl. brevis.

Baccae tres, monospermae. Albumen aequabile, iptus cavum. Embryo verticalis.

Palmace. [Corypha *Taliera* Roxb. corom. 3. t. 255. 256.]

 1308. *CALAMUS* L. (1764.)

Flores hermaphroditi, in spadice ramoso. Spatha communis nulla, partiales vagae.

Cal. trisepalus.

Pet. calyce longiora, acuminata.

Stam. Filamenta capillaria. Antherae rotundae.

Stylus conicus, apice trifidus.

Bacca globosa, squamoso-loricata, trisperma. Embryo basilaris.

Palmace. *Gärtn.* t. 139. [Palmijuncus *Rumph.* amboin. 5. t. 51 — 58.]

γ. Corolla tetrapetala.

 1309. *CLEOME* L. (1737.)

Cal. tetrasepalus, sepalo quarto inferiori.

Pet. inaequalia, ascendentia, basi glandulis tribus nectariferis.

Stam. utplurimum sena, nonnunquam abortu quaterna, vel coasitu octona, duodena vel plura. Filamenta vel stipiti germinis adnata, submonadelpha, vel calyci inserta, semper declinata. Antherae ovatae, erectae.

Styl. simplex. Stigma capitatum.

Siliqua stipitata, vel sessilis, unilocularis, bivalvis, placentis marginalibus filiformibus.

Capparideae. *Schk.* t. 189. b. *Gärtn.* t. 76. [Gynandropsis *Cand.* Polanisia *Rufin. Cand.*, Schepperia *Neck.* Macromerium *Burch.*]

 1310. *PERITOMA* Cand. (1824.)

Cal. cupulatus, quadridentatus, basi circumscissus.

Pet. ovata, subsessilia, aequalia.

Stam. stipiti inserta, basi monadelpha, aequalia. Antherae oblongae, recurvae.

Siliqua stipitata, oblonga, bivalvis, stylo persistente coronata.

Capparideae. [Atalanta *Nuttall.*]

 1311. *STERIPHOMA* * (1827.)

Cal. campanulatus bilobus.

Pet. erecta.

Stam. duo inferiora longiora. Antherae incumbentes.

Stylus nullus. Stigma capitatum.

Germen stipitatum, oblongum.

Capparideae. [Capparis paradoxa Jacq. schönbr. 1.
t. 111. Stephania Willd.]

1312. PHYSOSTEMON Mart. (1823.)

Cal. tetrasepalus.

Petala ascendentia.

Stam. hypogyna, declinata, inaequalia, Filamenta subulata, breviora infra antheram inflato-incrassata. Antherae erectae lineares, biloculares, demum recurvae.

Styl. subulatus, ascendens.

Siliqua bivalvis. Sem. uniseriata, rugoso-echinata.

Capparideae. Mart. nov. gen. t. 45—47.

δ. Corolla pentapetala.

1313. FRANKENIA L. (1740.)

Cal. tubulosus, canaliculatus, quinquedentatus.

Pet. unguiculata.

Stamina hypogyna. Filamenta capillaria, nuda. Antherae subrotundae.

Stylus trifidus.

Caps. unilocularis, tri-quadrivalvis; valvis placentiferis.

Sem. pauca subrotunda.

Frankenieae Hilar. Engl. bot. 205. Bot. mag. 2896.
Gärtn. t. 184. [Frauca Michel. nov. gen. t. 22. Beatsonia Roxb.]

1314. COSSINIA Commers. Juss. (1789.)

Cal. quinquepartitus, persistens.

Pet. quatuor aut quinque, unguiculata.

Stam. disco hypogyno inserta.

Styl. brevis, stigmati simplici.

Caps. tomentosa, trigona, trilocularis, apice dehiscens: loculis di-trispermis: placenta centrali.

Sapindeae. Lam. ill. t. 256.

ε. Corolla hexapetala.

1315. BERBERIS Brunsfels. (1532.)

Cal. hexasepalus, subcoloratus, tribracteatus.

Pet. basi biglandulosa, sepalis opposita.

Stam. hypogyna, petalis opposita. Filamenta vel elastico-articulata, vel dentata. Antherae bi-quadriloculares, loculis discretis, a basi ad apicem valvula elastica dehiscens.

Stigma subsessile, peltatum.

Bacca di - trisperma. Sem. testa crustacea.

Berberideae. Schk. t. 99. Ruiz et Pav. fl. peruv. 3. t. 280—283. Lam. ill. t. 253. Humb. nov. gen. 5. t. 430—434. Bot. reg. 702. 729. Pursh. fl. bor. amer. t. 4. 9. [Nahonia Nutt. Cand. Lessert ic. sel. 2. t. 3. 4.]

1316. N. ANDINA Thunb. (1781.)

Cal. sexfariam imbricatus, caducus.

Pet. calyce longiora, basi eglandulosa.

Stam. Antherae oblongae, subsessiles.

Styl. trigonus, brevissimus, persistens.

Bacca exsucca, disperma. Sem. hinc convexa, illinc concava.

Berberideae. Lam. ill. t. 261. Gärtn. t. 92.

1317. LEONTICE L. (1737.)

Cal. hexasepalus, ebracteatus, caducus.

Pet. sepalis opposita: unguibus petalorum squamatis.

Stam. petalis opposita, hypogyna: filamenta brevia: antherae biloculares.

Styl. brevissimus, stigmatē obtuso.

Caps. vel inflata, non dehiscens, unilocularis, subtrisperma; vel carnosa, dein rupta. Sem. globosa.

Berberideae. Lam. ill. t. 254. [Caulophyllum Michaux fl. bor. am. t. 21.]

1318. DIPHYLLIA Mx. (1803.)

Cal. trisepalus, deciduus.

Pet. sepalis maiora, patentia.

Stam. hypogyna, petalis opposita. Filamenta brevina, plana. Antherae biloculares, loculis a basi ad apicem, membranula soluta, dehiscentibus.

Stigma subsessile, capitatum.

Bacca subglobosa, unilocularis, di - trisperma.

Berberideae. Michaux fl. bor. amer. t. 19. 20.

1319. PEPLIS L. (1737.)

Cal. duodecimfidus, lacinis alternis angustioribus.

Pet. fugacia, aut nulla.

Stam. hypogyna. Antherae erectae, biloculares.

Stigma sessile capitatum.

Caps. bilocularis, polysperma.

Salicariae. Schk. t. 99.

b. *Fructus inferi.*

1320. LORANTHUS Vail. (1720.)

Cal. margo obsolete dentatus.

Pet. plerumque sena (interdum quaterna, raro octona.)

Stam. totidem, alterna breviora, petalorum basi affixa.

Antherae dorso adnatae, biloculares.

Styl. filiformis, stigmatē obtuso.

Bacca calyce persistente umbilicata, monosperma.

Loranthaceae. Jacqu. fl. austr. t. 30. Mirbel in ann. mus.
16. t. 21. Humb. nov. gen. 3. t. 299. 300.

1321. MYRRHINIUM Schott. (1827.)

Cal. quadrifidus, laciniis obtusis.

Pet. quatuor.

Stam. longissima: filamenta linearia, in aestivatione versus petala duplicata. Antherae oblongae.

Bacca oblonga, tri-quadrilocularis, polysperma, calyce coronata et bracteis binis suffulta.

— —

1322. BROMELIA Plum. (1703.)

Cal. tripartitus, corolla brevior.

Pet. convoluta, basi nuda.

Stam. basi petalorum inserta.

Styl. brevissimus. Stigmata tria carnosa, abbreviata.

Bacca trilocularis, polysperma.

Bromeliaceae. Trew. ic. Ehret. t. 51. Gärtn. t. 11.

1323. ANANASSA Plum. (1703.) Lindl. (1827.)

Cal. tripartitus, corolla brevior, carnosus, bracteatus.

Pet. convoluta, basi squamata, squamis geminis fimbriatis, tubulosis.

Stam. disco epigyno inserta.

Stylus filiformis. Stigmata tria, recta, carnosa.

Fructus carnosus baccatus, ex ovariiis bracteisque concretus.

Bromeliaceae. Trew. ic. Ehret. t. 2. Bot. reg. 1081.

1324. BILLBERGIA Thunb. (1823.)

Cal. tripartitus, corolla brevior, bracteatus.

Pet. convoluta, basi squamata.

Stam. basi sepalorum et petalorum inter squamas inserta.

Styl. filiformis. Stigmata tria linearia vel petaloidea.

Caps. baccata.

Bromeliaceae. Bot. reg. 1068. 1130. 1181. Bot. mag.
2892. [Bromelia. Hook. exot. fl. 41. 42. 143. Bot. mag.
1732. 2686. Bot. reg. 766.]

1325. AECHMEA Ruiz et Pav. (1802.)

Cal. duplex: exterior campanulatus coriaceus, tripartitus, laciniis duabus obtusis, tertia mucronata: interior exteriore longior, corolla brevior, tripartitus, laciniis revolutis.

Pet. convoluta, apice reflexa, basi squamis geminis munita.

Stam. basi petalorum inserta.

Stylus filiformis. Stigmata linearia, convoluta.

Caps. baccata.

Bromeliaceae. Ruiz et Pav. fl. peruv. 3. t. 264.

II. *Flores imperfecti.*

A. *Fructus superi.*

a. *Perianthium duplex.*

326. *POLLIA* Thunb. (1781.)

Perianthium bracteatum, utrumque tripartitum: exter. ovatum, concavum, obtusum; intern. acutum, reflexum, nervosum.

Stam. filamenta receptaculo inserta, capillaria, apice incurva. Antherae subrotundae, didymae.

Styl. subulatus, stigmate obtuso.

Bacca globosa, perianthio bracteaque persistentibus cincta, polysperma.

Smilacaceae. Thunb. ic. plant. jap. t. 25.

327. *TULBAGHIA* L. (1771.)

Perianthium extern. infundibuliforme, sexpartitum; internum trilobum, lobis bifidis, vel triphyllum, sexdentatum.

Stam. Antherae sessiles, biseriales ad internam perianthii interni paginam.

Styl. cylindricus, brevis. Stigma turbinatum, cavum.

Caps. trigona, trilocularis, loculis dispermis. Sem. triquetra.

Liliaceae. Lam. ill. t. 243. Bot. mag. 806. Gärtn. t. 16.

328. *ALBUCA* L. (1764.)

Perianthium utrumque tripartitum: externum calycinum, patens: internum foliolis forpiciatis, conniventibus, apice glandulosis.

Stam. alterna breviora (quandoque abortientia): filamenta crassiuscula, erecta. Antherae versatiles.

Stylus triqueter, stigmate capitato.

Caps. trilocularis, trivalvis. Sem. plana.

Asphodelaceae. Jacqu. ic. rar. 1. t. 63. 64. vol. 2. t. 440 — 446. Bot. reg. 277.

b. *Perianthium simplex.*

α. *Corollinum monopetalum.*

329. *THRINAX* L. fil. (1781.)

Spathae plures, vaginantes. Flores sessiles bracteati.

Cal. corollinus cupularis, sexdentatus.

Filamenta capillaria. Antherae lineares.

Styl. crassiusculus. Stigma subclavatum, oblique infundibuliforme.

Bacca monosperma sicca. Albumen solidum, aequabile.

Embryo subbasilaris.

Palmae. Swartz fl. Ind. occid. t. 13.

1330. *LOPHIOLO* Ker. (1814.)

Cal. corollinus persistens, regulariter sexpartitus, dense lanatus.

Stam. hypogyna, filiformia. Antherae ovatae erectae.

Styl. subulatus, tripartibilis.

Caps. trilocularis, polysperma.

Haemodorea? Bot. mag. 1596. [*Conostylis Pursh.*]

1331. *PONTEDERIA* L. (1737.)

Flores spatha suffulti.

Cal. corollinus tubulosus, sexpartitus, bilabiatus.

Stam. tria inferne, tria superne calycis tubo affixa; illa inclusa, haec exserta. Antherae oblongae, erectae.

Stylus simplex, declinatus. Stigma crassiusculum.

Caps. conica, apice lato reflexo, trilocularis, polysperma, axi centrali affixa.

Pontederaceae Kunth. Trew. ic. Ehret. t. 83. Bot. mag. 1156.

1332. *AGAPANTHUS* Herit. (1788.)

Cor. basi ventricosa, limbo regulariter sexpartito, lacinii oblongis, patentibus, unguiculatis.

Stam. declinata. Filamenta fauci inserta, capillaria, corollam aequantia. Antherae incumbentes.

Stylus declinatus, filiformis.

Caps. trigona, trilocularis, polysperma. Sem. hinc alata.

Liliaceae. Redout. lil. t. 6. [*Crinum Gärtn.* t. 83. *Mauhlia linearis* Thunb. Sed altera, *ensifolia*, habitu, corollae forma antherarumque, staminum directione, stylo et nectario differt. Neque huc, neque ad *Massoniam* (*Andr.*), neque ad *Polianthen* (*Jacqu.*) neque ad *Hyacinthum* (*Car.*) trahenda.]

1333. *BULBOCODIUM* L. (1737.)

Cor. infundibuliformis, tubo longissimo, lacinii limbi unguiculatis.

Stam. unguibus petalorum inserta. Antherae incumbentes.

Stylus filiformis, stigmate trifido.

Caps. trigona, trilocularis, polysperma.

Liliaceae. Bot. mag. 153. Redout. lil. t. 197. [*Meredera Ramond.* differt stylis ternis discretis. Genus peculiare esse nequit, cum et in *Bulbocodio* stylorum connatorum vestigia adsint. *Lapeyroux.* hist. abrég. des plant. des Pyrén. p. 201. 202. *Marschall.* Bieberst. cent. pl. ross. t. 50.]

1334. *BLANDFORDIA* Sm. (1804.)

Cor. tubulosa, ore quinquelobo.

Stam. tubo inserta. Antherae dactylethraeiformes, basi impositae.

Stylus subulatus, stigmate simplici.

Caps. prismatica, tripartibilis. Sem. biseriata, marginibus suturae inserta, testa laxa pubescente.

Liliaceae. Smith. exot. bot. 1. t. 4. [*Aletris Labill.* t. 111.]

335. *FUNKIA* * (1817.)

Cor. campanulata, basi ventricosa, super ovario contracta; limbo sexfido, subringente.

Stam. Filamenta basi corollae inserta, declinata. Antherae incumbentes.

Styl. declinatus. Stigma obtusum.

Caps. trilocularis, polysperma. Sem. membrana obvoluta, embryonibus pluribus feta.

Liliaceae. [Hemerocallis Andr. repos. 6. 194. Hosta Trattin. arch. 2. t. 89.]

336. *YUCCA* Dalech. (1586.)

Cor. campanulata, regulariter sexpartita.

Stam. basi corollae inserta; filamentis superne crassioribus. Antherae minores, biloculares.

Styl. nullus. Stigma trilobum.

Caps. prismatica, trigona, carnosa, sexlocularis. Sem. compressa.

Liliaceae. Bot. mag. 1260. Gärtn. t. 85.

337. *LILIUM* Virg. Plin.

Cor. campanulata, regulariter sexpartita, laciniis patentireflexis, sulco nectarifero longitudinali instructis (qui interdum deest.)

Stam. basi corollae inserta. Antherae versatiles.

Styl. columnaris. Stigma trilobum.

Caps. trigona, sexsulca, trilocularis, trivalvis; valvis reticulo fibroso nexis. Sem. compressa.

Liliaceae. Bot. mag. 30. 36. 259. 278. 519. 798. 800. 858. 893. 936. 1165. 1237.

338. *HEMEROCALLIS* L. (1737.)

Cor. campanulata, regulariter sexpartita, laciniis patentireflexis.

Stam. basi corollae inserta, declinata. Antherae subincumbentes, biloculares.

Stylus declinatus. Stigma trigono-capitatum.

Caps. trigona, trilocularis. Sem. globosa.

Liliaceae. Schk. t. 98. a.

339. *ALOË* Diosc.

Cor. tubulosa, fundo nectarifero; limbo patulo vel regulariter sexfido, vel subringente.

Stam. receptaculo inserta. Antherae oblongae incumbentes.

Styl. simplex. Stigma trilobum.

Caps. trilocularis, trivalvis: placenta centralis. Sem. triquetra, aut compressa, alata.

Liliaceae. Gärtn. t. 17. [Apicra Willd. Haworthia Duval., corollis subringentibus aut bilabiatis. Gasteria Duval., corollis incurvis.]

340. *HYACINTHUS* Colum.

Cor. infundibuliformis vel campanulata, limbo regulari sexfido patente.

Stam. basi corollae inserta.

Germen apice glandulas tres nectariferas gerens. *Stylus* brevis. *Stigma* obtusum trilobum.

Caps. trilocularis, loculis oligospermis.

Liliaceae. Schk. t. 96. Gärtn. t. 12.

1341. *MUSCARI* Clus. (1578.)

Cor. globoso-ovata vel subcylindrica, fauce constricta, limbo abbreviato sexdentato.

Stam. Antherae subsessiles, intra tubum.

Germen eglandulosum. *Stylus* brevissimus, trisulcus.

Stigma trigonum.

Caps. trilocularis, loculis dispermis.

Liliaceae. Schk. t. 96.

1342. *ADAMSIDIA* Willd. (1812.)

Cor. campanulata, tubo brevi; limbo patulo sexfido.

Dentes brevissimi emarginati sex ad faucem corollae.

Stam. Antherae oblongae subsessiles inter dentes faucis.

Styl. columnaris. *Stigma* crassiusculum.

Caps.?

Liliaceae. [Puschkinia MB. cent. 2. pl. ross. t. 91.]

1343. *TRITOMA* Ker. (1805.)

Cor. tubulosa, subcylindrica; limbo sexdentato.

Stam. exserta, alterna longiora. Filamenta receptaculo inserta. Antherae incumbentes.

Stylus filiformis. *Stigma* simplex.

Caps. triquetra, trilocularis, polysperma.

Liliaceae. Bot. mag. 744. 758. 764. [Alettris Andr. repot. 54. Veltheimia Uvaria et pumila W.]

1344. *ALETTRIS* L. (1754.)

Cal. corollinus, tubulosus, rugosus; limbo regulari sexfido.

Stam. erecta. Filamenta brevissima, laciniarum basi inserta. Antherae subsagittatae.

Styl. triqueter, stigmatе trigono.

Caps. semiinfera, calyce vestita styloque coronata, trilocularis, apice trivalvis. Placenta centralis. Sem. numerosa, minutissima, arcuata.

Haemodoreae? Bot. mag. 1418. [Wurmbea Willd. hort. berol. t. 8.]

1345. *VELTHEIMIA* Gleditsch. (1769.)

Cor. tubulosa, superne ampliata; limbo sexdentato, erecta.

Stam. tubo inserta, corolla breviora.

Stylus filiformis, exsertus.

Caps. membranacea, trialata, subtrisperma.

Liliaceae. Jacqu. hort. schönbr. 1. t. 77. 78. Bot. mag. 501. 1091. [Alettris capensis L. A. glauca Ait.]

1346. *DRIMIA* Jacqu. (1790.)

Cor. campanulata, sexfida, laciniis revolutis.

Stam. basi corollae inserta. Filamenta corollam aequantia.

Antherae subglobosae.

Stylus exsertus. Stigma capitatum.

Caps. trilocularis, polysperma.

Liliaceae. Jacqu. ic. rar. 2. t. 373 — 377. Bot. mag. 1380. 1444.

1347. MILLEA Cav. (1793.)

Cor. tubo longissimo, limbo patente sexfido; laciniis alternis angustioribus, apice uncinulatis.

Stam. fauci corollae inserta. Filamenta brevissima. Antherae oblongae erectae.

Germen pedicellatum. Stylus filiformis.

Stigmata tria, globulosa, pilosa.

Caps. trilocularis, trivalvis, polysperma.

Liliaceae. Cav. ic. 2. t. 196. [Bravoa Lexarz. fueritne species altera huius generis: sed antherae dicuntur versatiles. Tertia species, etiam magis differens, habetur ab Hookerio, ad bot. mag. 2877. Brodiaea grandiflora Pursh; non Smith. Hookeria coronaria Salisb. parad. lond. t. 98.]

1348. POLIANTHES L. (1737.)

Cor. calycina, infundibuliformis: limbo nutante, aequaliter sexfido, patente.

Stam. subexserta. Filamenta tubo adnata. Antherae oblongae erectae.

Stylus triqueter, inclusus. Stigma trilobum.

Caps. basi calyce tecta, trilocularis, trivalvis. Sem. compressa, biserialia.

Liliaceae. Bot. reg. 63.

1349. UROPETALUM Ker. (1816.)

Cal. corollinus campanulatus; limbo duplici, exteriori profundius diviso; laciniis longioribus, angustioribus, reflexis; interioribus erectis, conniventibus.

Stam. tubo inserta: filamenta subulata. Antherae mucronatae, inclusae.

Stylus filiformis, subtriangularis. Stigma trilobum.

Caps. membranacea, trilocularis, trivalvis, valvis septifidis. Sem. plana.

Liliaceae. Asphodeleae. Bot. reg. 156. 974. [Hyacinthus serotinus L. Bot. mag. 859., H. viridis Jacqu. ic. rar. 1. t. 66. Lachenalia Thunb. Ait. Zuccagnia Thunb. Willd. enum. Polemannia Berg. Schlechtend.]

1350. EUCOMIS Herit. (1788.)

Cor. subrotata, regularis, sexpartita.

Stam. Filamenta basi dilatata, subconnata. Antherae ovaes.

Styl. subulatus, stigmate simplici.

Caps. trilocularis, polysperma.

Asphodeleae. Andr. repos. 369. Bot. mag. 913. 1539. [Basilaea Lam. ill. 239.]

1351. *CHLOROPHYTUM* Ker. (1808.)

Cal. corollinus, campanulato - rotatus, tubo perbrevis, limbo sexpartito: laciniis exterioribus persistentibus.

Stam. hypogyna: filamenta filiformia: antherae oblongae, erectae.

Styl. filiformis, persistens. Stigma capitellatum.

Caps. triloba, trilocularis, loculis bi - trispermis. Placenta centralis. Sem. reniformia nigra.

Asphodeleae. Bot. mag. 1071. Bot. reg. 813.

1352. *NARTHECIUM* Möhring. (1778.)

Cal. corollinus, sexpartitus, patens, persistens.

Stam. hypogyna, laciniis calycis opposita. Filamenta dense pilosa. Antherae oblongae, demum tortae.

Styl. brevissimus, conicus.

Caps. prismatica, trilocularis, polysperma. Sem. utrinque processibus filiformibus.

Asphodeleae. Engl. bot. 535. [Alabama Adans. Cand.]

1353. *LAXMANNIA* R. Br. (1810.)

Cal. corollinus sexpartitus, persistens; laciniis alternis longioribus.

Stam. laciniis inserta. Filamenta subulata, glabra. Antherae peltatae, subrotundae.

Styl. stigmatate obtuso.

Caps. trilocularis, trivalvis; valvis medio septiferis. Sem. subbina, peltata.

Asphodeleae.

1354. *BAUMGARTENIA* * (1817.)

Flores capitati, bracteati.

Cor. infundibuliformis; limbo sexpartito aequali.

Stam. fauci corollae inserta. Filamenta filiformia. Antherae ovato - oblongae, erectae.

Stylus filiformis. Stigma obtusum.

Caps. bracteis persistentibus inclusa, trilocularis, trivalvis; valvis medio septiferis: loculis subdispermis.

Asphodeleae. [Borya Labill. nov. holl. t. 107.]

1355. *XANTHORRHOEA* Smith. (1798.)

Flores spicati, confertissimi, bracteis imbricatis muniti.

Cal. corollinus sexpartitus, persistens: laciniis interioribus concavis.

Stam. basi calycis affixa, exserta. Filamenta linearis, glabra. Antherae versatiles.

Stylus columnaris, trisulcus.

Caps. lignea, ovata, trilocularis, trivalvis: loculis mono-dispermis. Sem. compressa, atra.

Asphodeleae.

1356. *HERRERIA* Ruiz et Pav. (1800.)

Cor. rotata, profunde sexpartita, laciniis alternis angustioribus.

Stam. basi corollae inserta, laciniis opposita. Filamenta subulata. Antherae erectae, subrotundae, biloculares.

Stylus triangularis. Stigma trigonum.

Caps. trialata, trilocularis, trivalvis; loculis di-tetraspermis. Sem. plana, orbicularia, marginata.

Asphodeleae. Ruiz et Pav. fl. per. 3. t. 103. [Salsa Feuill. 2. t. 7.]

1357. *DIANELLA* Lam. (1786.)

Cor. rotata, regularis, sexpartita, decidua.

Stam. hypogyna. Filamenta curvata, apice incrassata, stuposa. Antherae lineares, erectae.

Styl. filiformis. Stigma simplex.

Bacca globosa, polysperma.

Asphodeleae. Bot. mag. 505. Bot. reg. 734. 1120.

1358. *DRACAENA* Vandell. (1762.)

Cor. patens, regularis, sexpartita, laciniis subrevolutis.

Stam. basi corollae inserta. Filamenta basi membranacea, medio incrassata, apice subulata.

Styl. subangulatus. Stigma trifidum.

Bacca sessulca, trilocularis, loculis plerumque monospermis.

Asphodeleae. Vandell. in Römer script. t. 2. Jacqu. schübr. 1. t. 95. 96. Bot. mag. 1081. 1180.

1359. *CORDYLINE* Commers. R. Br. (1810.)

Cal. campanulata, regulariter sextida, marcescens.

Stam. fauci inserta. Filamenta subulata, glabra. Antherae versatiles, basi bifidae.

Styl. filiformis. Stigma trilobum.

Bacca globosa, trilocularis, loculis pleiospermis. Sem. umbilico strophiolato.

Asphodeleae. [Dracaena australis Forst. Hook. in Bot. mag. 2835. Drac. stricta Bot. mag. 2575. Bot. reg. 956. Charlwoodia Sweet. flor. austr. t. 18.]

1360. *SANSEVIERA* Thunb. (1794.)

Cor. tubulosa, basi subventricosa marcescens; limbo sexpartito, laciniis revolutis.

Stam. fauci imposita, subulata. Antherae oblongae, incumbentes.

Styl. exsertus. Stigma capitatum.

Bacca trilocularis, loculis monospermis.

Asphodeleae. Bot. reg. 160. Bot. mag. 2631. [Salmia Cav. ic. 3. t. 246.]

1361. *LUZURIAGA* Ruiz et Pav. (1800.)

Cor. sexpartita, (hexapetaloidea) patens, decidua; laciniis patentibus.

Stam. basi lacinarum inserta. Filamenta filiformia, glabra, apice curvata. Antherae conniventes, sagittatae, filamentis longiores.

Styl. filiformis, trisulcus. Stigma simplex.

Bacca trilocularis; loculis submonospermis.

Asphodelaceae. Ruiz et Pav. fl. peruv. 3. t. 298.

1362. EUSTREPHUS R. Br. (1810.)

Cor. sexpartita, (hexapetaloidea) patens, decidua: laciniis interioribus fimbriatis.

Stam. hypogyna. Filamenta brevissima; plana, subcoacta. Antherae erectae.

Styl. filiformis. Stigma trigonum.

Caps. baccata, trilocularis, trivalvis, valvis medio septiferis: loculis pleiospermis. Sem. strophiolata.

Asphodelaceae. Bot. mag. 1245.

1363. ASPARAGUS Colum.

Cor. subcampanulata, sexpartita, decidua.

Stam. basi corollae inserta. Filamenta subulata. Antherae erectae, subcordato-ovatae.

Styl. brevis. Stigma trilobum.

Bacca globosa, trilocularis, oligosperma.

Sarmentaceae. Schk. t. 96.

1364. CONVALLARIA L. (1737.)

Cor. tubulosa, limbo sexfido.

Stam. tubo adnata. Filamenta subulata. Antherae erectae, subcordato-ovatae.

Styl. columnaris. Stigma capitatum.

Bacca trilocularis, loculis oligospermis.

Sarmentaceae. Schk. t. 97. Sturm. fl. germ.

1365. SMILACINA Desf. (1807.)

Cor. hexapetaloidea, rotata s. stellata.

Stam. perigyna.

Stylus filiformis.

Bacca trilocularis, loculis oligospermis.

Sarmentaceae. Desfont. in ann. mus. 9. t. 8. 9. [Convallariae spec. L.]

1366. STREPTOPUS Michaux. (1803.)

Cor. hexapetaloidea, campanulata, laciniis basi nudis.

Stam. hypogyna. Filamenta abbreviata. Antherae cordato-lanceolatae, erectae.

Stylus conicus, trisulcus, abbreviatus.

Bacca trilocularis, oligosperma. Sem. nuda.

Sarmentaceae. Michaux flor. bor. amer. t. 18. Sturm. fl. germ. Bot. mag. 1489. 1490. [Uvulariae spec. L.]

Pedunculi distorti vel geniculati.

1367. UVULARIA L. (1737.)

Cor. hexapetaloidea, campanulata; laciniis basi fovea nectarifera instructis.

Stam. hypogyna. Filamenta brevissima.

Styl. abbreviatus. Stigmata tria revoluta.

Caps. trivalvis, valvis septiferis. Sem. membrana obvoluta.

Sarmentaceae. *Smith* exot. bot. 1. t. 49 — 52.

1368. PHILESIA Commers. Juss. (1789.)

Cor. calycina, bracteata, campanulata: lacinae exteriores breviores acutae, interiores elongatae, obtusae.

Stam. basi corollae affixa. Filamenta basi connata. Antherae versatiles.

Styl. simplex. Stigmata tria.

Bacca subtrigona, polysperma.

Sarmentaceae. *Lam.* ill. t. 248. [*Lapageria Ruiz et Pav.* fl. peruv. 3. t. 297.]

1369. RHIPOGONUM Forst. (1775.)

Cor. sexpartita, patens, decidua, bracteis binis suffulta.

Stam. basi laciniarum inserta. Filamenta subulata. Antherae erectae, cordato-lanceolatae.

Stylus brevissimus. Stigma trigonum.

Bacca mono-disperma.

Sarmentaceae. *Forst.* char. 25.

1370. ONCUS Lour. (1790.)

Cor. campanulata, pilosa; tubo oblongo, hexagono; limbo sexfido, reflexo; laciniis subulatis. Bractae binae corollam amplexantes.

Stam. basi laciniarum inserta. Filamenta brevissima.

Antherae subrotundae, minimae.

Styl. brevis, tripartitus. Stigmata tria, reflexa, apice bifido.

Bacca oblonga, trilocularis, polysperma.

Sarmentaceae.

1371. SMILAX Theophr.

Flores dioeci.

Cal. corollinus, hexasepaloides, patens, persistens.

Stam. basi laciniarum inserta. Filamenta subulata. Antherae erectae.

Styl. brevissimus. Stigmata tria.

Bacca mono-trisperma. Embryo umbilico oppositus.

Sarmentaceae. *Schk.* t. 328. Bot. mag. 1846. *Gärtn.* t. 16.

1372. OPHIOPOGON Ker. (1801.)

Cal. corollinus, semisuperus; limbo sexpartito, patente.

Stam. epigyna. Filamenta brevissima. Antherae erectae, cordato-lineares.

Styl. triqueter. Stigma trilobum.

Bacca subtrisperma.

Sarmentaceae. Bot. mag. 1063. Bot. reg. 593. [*Convallaria japonica Thunb.* *Liriope spicata Lour.* *Flüggea Rich.* *Slatertia Desv.*]

1373. *PELIOSANTHES* Andr. (1808.)

Cal. corollinus, semisuperus, sexlobus: fauce fornicata, antheras fovens.

Antherae sessiles, didymae.

Styl. subnullus. Stigmata tria obtusa.

Baccae tres clavatae, monospermae.

Sarmentaceae? Andr. repos. 605. Bot. mag. 1302. 1532. (Pedicelli articulati.)

1374. *MIEERSIA* Lindl. (1826.)

Bracteeae patentes, sex exteriores petaloideae, interiores bifidae, coloratae, depauperatae.

Perianth. urceolatum, carnosum, ore sexdentato.

Stam. minima, fauci perianthii inserta.

Stigma capitatum.

Caps. trilocularis, apice trivalvis, polysperma.

Gillisiaeae Lindl.

1375. *CALECTASIA* R. Br. (1810.)

Cal. corollinus, tubulosus, persistens; limbo patente, sexfido.

Stam. fauci inserta. Filamenta brevissima. *Antherae* erectae oblongae, apice biporosae.

Styl. filiformis.

Utriculus monospermus, tubo calycis indurato inclusus.

— — R. Brown ad Flinders it. t. 9.

1376. *TUPISTRA* Ker. (1814.)

Cor. bracteata, campanulata, subcarnosa; limbo sexfido, subpatente, subirregulari.

Antherae subsessiles, tubo summo adnatae, biloculares.

Styl. crassus, trisulcus, tubulatus. *Stigma* trilobum, plicato-crenatum.

Bacca trilocularis.

— — Bot. mag. 1655. Bot. reg. 704.

1377. *RONDEA* Roth. (1821.)

Spadix cylindricus, floribus tectus: spatha basilaris, triphylla.

Cal. corollinus, subrotundo-hexagonus, subcoriaceus, ore sexdentato.

Stam. calyci adnata. *Antherae* didymae, fauci calycis inhaerentes.

Styl. brevis, crassus. *Stigma* trifidum.

Bacca monosperma, stylo coronata.

Aroideae. [Orontium japonicum Banks. ic. Kämpfer. t. 12. Bot. mag. 898.]

β. *Perianthium pleiopetalum.*

1378. *ORNITHOGALUM* Nicand. Dioscor.

Cal. corollinus, hexasepalus, patens, marcescens.

Stam. receptaculo inserta. Filamenta vel basi dilatata, apice attenuata; vel plana, fere petaloidea, emarginata, bi- tricuspidata. Antherae versatiles, biloculares.

Styl. columnaris, stigmatе trigono.

Caps. trilocularis, trivalvis, polysperma. Sem. ovato-globosa.

Asphodelaceae. Schk. t. 94. Sturm. fl. germ. Reichenb. ic. t. 117. 142. [Gagea Salisb. Sweet.]

1379. *MASSONIA* Thunb. (1781.)

Cor. hexapetala, patens, vel reflexa.

Stam. subulata, erecta, basi in annulum unita. Antherae biloculares.

Germen sexangulare, inter angulos alternos poris nectariferis instructum.

Styl. filiformis, trisulcus.

Caps. triquetra, trilocularis, polysperma. Sem. globosa.

Asphodelaceae. Ait. hort. kew. ed. 1. t. 3. Bot. reg. 694. 998. Gärtn. t. 182.

1380. *ALLIUM* Pin.

Flores subumbellati. Spatha umbellam fulciens.

Cor. calycina, hexapetala, patens, marcescens.

Stam. petalorum basi inserta. Filamenta vel aequalia subulata; vel basi dilatata subconnata; vel tricuspidata.

Antherae incumbentes, biloculares.

Styl. filiformis. Stigma capitatum.

Caps. trilocularis, polysperma.

Asphodelaceae. Schk. t. 91. Sturm. fl. germ. Reichenb. ic. 5. t. 415 — 418. 427 — 430.

1381. *SOWERBAEA* Sm. (1798.)

Flores umbellati, bracteati, pedicellis articulatis.

Cor. calycina, hexapetala, patens, marcescens.

Stam. basi petalorum inserta, tria fertilia, petalis opposita, tria abortientia. Antherae bilobae, lobis distinctis.

Styl. filiformis, stigmatе simplici.

Caps. trilocularis, trivalvis; valvis medio septiferis. Sem. subsolitaria, peltata.

Asphodelaceae. Smith. in Linn. transact. 5. t. 6. Andr. repoa. 81. Bot. mag. 1104.

1382. *SCILLA* Theophr.

Cor. hexapetala, vel subcampanulata, vel patentissima, decidua.

Stam. basi petalorum inserta. Filamenta subulata, aequalia. Antherae incumbentes, biloculares.

Styl. simplex.

Caps. trilocularis, trivalvis, polysperma. Sem. subglobosa.

Asphodeleae. Schk. t. 94. Engl. bot. 377. Bot. mag. 1574.

1383. *BARNARDIA* Lindl. (1826.)

Cor. hexapetala, patens, persistens.

Stam. basi petalorum inserta. Filamenta basi dilatata, fimbriata. Antherae erectae, oblongae.

Stylus simplex.

Germen triloculare, loculis monospermis.

Asphodeleae. Bot. reg. 1029.

1384. *CYANELLA* L. (1740.)

Cor. hexapetala, patens, petalis subinaequalibus, interioribus longioribus.

Stam. hypogyna. Filamenta basi in urceolo carnoso connata. Antherae erectae: quinque aequales, poro verticali dehiscentes: sexta duplo maior, oblonga.

Styl. declinatus.

Caps. trilocularis, trivalvis, polysperma.

Asphodeleae. Bot. mag. 568. 1252. Bot. reg. 1111.

1385. *LACHENALIA* Jacqu. (1787.)

Cor. hexapetala, vel hexapetaloidea, campanulata: petala exteriora breviora, saepe apice callosa; interiora longiora, acuta.

Stam. basi petalorum adnata. Filamenta subulata. Antherae oblongae.

Styl. subulatus.

Caps. triquetra, trilocularis, pleiosperma. Sem. globosa.

Asphodeleae. Jacqu. ic. rar. 2. t. 382. 384. 391. 394. 395. 401. Bot. mag. 1497. 1517. 1611. Bot. reg. 287. 314.

1386. *PHORMIUM* Forst. (1775.)

Cor. hexapetala, vel hexapetaloidea, campanulata: petala exteriora breviora, acuta; interiora longiora, rotundata.

Stam. basi corollae adnata, ascendentia, exserta. Antherae erectae, subtriquetrae.

Styl. filiformis, ascendens.

Caps. trilocularis, polysperma.

Asphodeleae. Forst. char. 24. Cook it. 2. t. 96. Redout. lil. t. 448. 449.

1387. *ERIOSPERMUM* Jacqu. (1790.)

Cor. calycina hexapetala, patentiuscula.

Stam. basi petalorum inserta; filamenta planiuscula conniventia. Antherae sagittato-didymae.

Styl. triquetra, stigmatibus subtrilobo.

Caps. trigona, trilocularis, loculis oligospermis. Sem. papposa.

Asphodeleae. Jacqu. ic. rar. 2. t. 420—422. Bot. mag. 1382. Bot. reg. 578. 795.

1388. *ASPHODELUS* Theophr.

Cor. hexapetala, patens.

Stam. Filamenta e fornicibus cum basi petalorum connexis surgentia, arcuata, subulata. Antherae incumbentes.

Styl. subulatus, ascendens.

Caps. coriacea, trilocularis, pleiosperma. Sem. triquetra.

Asphodeleae. Schk. t. 95. Bot. mag. 964.

1389. *EREMURUS* Marsch. Bieb. (1818.)

Cor. hexapetala, patens, post anthesin involuta.

Stam. basi petalorum affixa, primum intra corollam conduplicata, post anthesin valde elongata ac rigentia.

Stylus demum reflexus.

Caps. trilocularis, pleiosperma. Sem. trigona.

Asphodeleae. Marsch. Bieb. cent. pl. ross. 2. t. 61.

[*Asphodelus altaicus* Pall. in act. petrop. 1779. t. 10.]

1390. *ANTHERICUM* L. (1764.)

Cor. hexapetala, patens.

Stam. hypogyna. Filamenta subulata, glabra. Antherae incumbentes, biloculares.

Germen glandulis tribus nectariferis.

Styl. subulatus. Stigma subclavatum.

Caps. trilocularis. Sem. trigona.

Asphodeleae. Schk. t. 95. Bot. reg. 567. 877. [Phalangium Tourn. Lam. ill. t. 240.]

1391. *BULBINE* Theophr. L. (1737.)

Cor. hexapetala, patens.

Stam. hypogyna. Filamenta subulata, barbata. Antherae incumbentes, biloculares.

Styl. subulatus.

Caps. trilocularis. Sem. trigona.

Asphodeleae. [Anthericum Jacqu. ic. var. 2. t. 403. 407. 408. Lam. 240. Bot. mag. 1317. 1339. 1454.]

1392. *ARTHEPODIUM* R. Br. (1810.)

Cor. hexapetala, patens; petala interiora margine undulata s. fimbriata. Pedicelli articulati.

Stam. hypogyna. Filamenta barbata. Antherae erectae, basi emarginatae.

Styl. filiformis. Stigma hispidulum.

Caps. trilocularis, trivalvis. Sem. angulata.

Asphodeleae. Bot. mag. 1421. [Anthericum Radout. lil. 58.]

1393. *STYPANDRA* R. Br. (1810.)

Cor. hexapetala, patens: petala aequalia. Pedicelli articulati.

Stam. hypogyna. Filamenta basi attenuata, curvata, inferne glabra, superne stuposo-barbata. Antherae erectae, basi emarginatae.

Styl. filiformis. Stigma simplex.

Caps. trilocularis, trivalvis, oligosperma.

Asphodeleae.

1394. CAESIA R. Br. (1810.)

Cor. hexapetala, patens: petala aequalia. Pedicelli articulati.

Stam. hypogyna. Filamenta glabra, utrinque attenuata. Antherae erectae, basi emarginatae.

Styl. filiformis.

Caps. clavata, trilocularis: loculis dispermis. Sem. ventricosa, strophiolata.

Asphodeleae.

1395. TRICORYNE R. Br. (1810.)

Cor. hexapetala, patens: petala aequalia. Pedicelli articulati.

Stam. hypogyna. Filamenta penicillato-barbata. Antherae erectae, basi emarginatae.

Styl. filiformis.

Nuces tres, clavatae, monospermae.

Asphodeleae.

1396. THYSANOTUS R. Br. (1810.)

Cal. corollinus, hexasepalus, patens: petala interna latiora, colorata, fimbriata.

Stam. hypogyna, declinata; alterna saepe longiora, nonnunquam tantum terna. Antherae erectae, lineares; alternae saepe longiores.

Styl. filiformis, declinatus.

Caps. trilocularis, trivalvis, loculis dispermis: semine altero erecto, altero pendulo.

Asphodeleae. Bot. reg. 655. 656.

1397. CZACKIA Andr. (1818.)

Cor. hexapetala, campanulato-infundibuliformis.

Stam. hypogyna, declinata, corollam aequantia. Filamenta filiformia. Antherae incumbentes.

Stylus exsertus. Stigma trilobo-capitatum.

Caps. subhexagona, trilocularis, trivalvis. Sem. angulata.

Asphodeleae. Andrzejowski libellus singularis cum ic. Crzeminiac. 1818. 4. [Anthericum Liliastrum L. spec. ed. 2., Heimerocallis Liliastrum L. spec. ed. 1.]

1398. TULIPA O. Gesner. (1560.)

Cor. hexapetala, campanulata, decidua. Petala basi nuda.

Stam. hypogyna. Filamenta brevia. Antherae erectae, oblongae.

Styl. nullus. Stigma trilobum.

Caps. trigona, trilocularis, trivalvis. Sem. compresso-plana, gemino ordine sibi incumbentia.

Liliaceae. Schk. t. 93.

1399. *FRITILLARIA* C. Gesner. (1576.)

Cor. hexapetala, campanulata, (plerumque pendula aut nutans.) Petala inferne foveola nectarifera.

Stam. hypogyna. Filamenta subulata. Antherae erectae, oblongae.

Styl. simplex, stigmatе trifido.

Caps. oblonga, triloba, trilocularis, trivalvis. Sem. plana, gemino ordine sibi incumbentia.

Liliaceae. Schk. t. 92. [Imperialis Juss.]

1400. *ERYTHRONIUM* Dioscor.

Cor. hexapetala, basi campanulata, a medio reflexa, patens. Petala lanceolata, acuta: alterna callis binis ad basin instructa.

Stam. hypogyna. Filamenta subulata, brevissima. Antherae erectae, oblongae.

Stylus rectus. Stigma trifidum.

Caps. obovata, trilocularis, trivalvis, valvis placentiferis. Sem. ovata acuta.

Liliaceae. Schk. t. 92.

401. *GLORIOSA* L. (1737.)

Cor. hexapetala, a basi inde reflexa. Petala lineari-lanceolata, acuminata, undulata, longissima.

Stam. hypogyna. Filamenta subulata, elongata, patentia. Antherae incumbentes.

Styl. filiformis, inclinatus. Stigma trifidum.

Caps. trilocularis, trivalvis. Sem. baccata, globosa, duplici serie.

Liliaceae. Bot. reg. 77. Andr. rep. 129. Gärtn. t. 18. [Methonica Herm. hort. lugd. bat. 669. Redout. lîk 26.]

402. *LOMATOPHYLLUM* Willd. (1811.)

Cor. hexapetala, in tubuli formam connivens: petalis lanceolatis obtusis.

Stam. hypogyna. Filamenta subulata. Antherae erectae.

Styl. filiformis. Stigma simplex.

Caps. baccata, trisulca, trilocularis; loculis monotrispermis.

Liliaceae. [Aloë marginalis Cand. succul. t. 31.]

403. *LEDEBOURIA* Roth. (1821.)

Cal. corollinus, hexasepalus. Sepala oblongo-linearîa, basi conniventia, erecto-patula.

Stam. basi sepalorum inserta. Filamenta subulata. Antherae dorso affixae.

Styl. capillaris, simplicissimus. Stigma acuminatum.

Utriculi tres monospermi obovati.

Sarmentaceae.

404. *CALLIXENE* Commers. Juss. (1789.)

Cor. hexapetala, patens. Petala alterna basi biglandulosa.

Stam. basi petalorum inserta. Filamenta basi latiora.

Antherae versatiles, oblongae.

Styl. simplex. Stigma trigonum.

Bacca trilocularis, loculis subtrispermis.

Sarmentaceae. Lam. ill. t. 248.

1405. *COMPSANTHUS* * (1827.)

Cor. hexapetala, campanulata. Petala apice incrassata, exteriora basi gibboso-saccata.

Stam. aequalia. Filamenta basi dilatata, subconnata.

Antherae incumbentes, ovali-oblongae, biloculares.

Styl. brevis. Stigmata tria, revoluta, apice bifida.

Caps. elongata, triquetra, trilocularis: loculis polyspermis.

Sem. angulata, simplici serie disposita.

Sarmentaceae. [Compositae Don.]

1406. *DRYMOPHILA* R. Br. (1810.)

Cor. hexapetala, patens.

Stam. hypogyna.

Styl. tripartitus. Stigmata revoluta.

Bacca trilocularis, polysperma.

Sarmentaceae.

1407. *BURCHARDIA* R. Br. (1810.)

Cor. hexapetala, patentissima: in petalorum unguibus foveola nectarifera.

Stam. basi petalorum inserta. *Antherae* peltatae, posticae.

Styl. tripartitus. Stigmata acuta.

Caps. tripartibilis, polysperma.

Melanthaceae R. Br.

1408. *SCHELHAMMERA* R. Br. (1810.)

Cor. hexapetala, campanulato-patens. Petala aequalia, unguiculata.

Stam. basi petalorum inserta. Filamenta subulata. *Antherae* oblongae, dorso affixae.

Styl. simplex. Stigmata tria revoluta.

Caps. trilocularis, trivalvis; valvis medio septiferis. *Sem.* globosa.

Melanthaceae. Hook. in Bot. mag. 2712.

1409. *APHYLLANTHES* Lobel. (1570.)

Cal. corollinus, hexasepalus, patens; bracteis scariosis, imbricatis suffultus. Petala unguiculata.

Filamenta setacea, petalorum laminis inserta. *Antherae* oblongae.

Styl. filiformis. Stigmata tria, oblonga.

Caps. turbinata, trilocularis, polysperma.

Juncea. Lam. ill. t. 252.

1410. *JUNCUS* Plin.

Cal. corollinus, hexasepalus, bibracteatus.

Stam. hypogyna (quandoque terna). *Antherae* oblongae erectae.

Styl. simplex. *Stigmata* tria pilosa, vel barbata.

Caps. trivalvis polysperma; unilocularis, septis persistentibus columnam centralem constituentibus, raro placentis parietalibus; subinde trilocularis. *Testa* seminis, vel conformis nucleo, vel utrinque alata.

Junceae. *Sturm.* fl. germ.

α. *Cephaloxys* Desv. (1808.) triandra. *Caps.* prismatica, trivalvis, valvis septiferis, septis columellam tri-alatam mentientibus. (*Journ. de botan.* 1. t. 11.)

β. *Rostcovia* Desv. (1808.) hexandra. *Caps.* unilocularis, placentis tribus parietalibus. (*Journ. de bot.* 1. t. 12. f. 2.)

γ. *Marsippospermum* Desv. (1808.) hexandra. *Caps.* oblonga, unilocularis: testa seminum utrinque producta, acuminata. (*Journ. de bot.* 1. t. 12. f. 1.) Cf. *Laharpe* in act. soc. hist. nat. paris. 3. p. 104.

1411. *LUZULA* Cand. (1805.)

Cal. corollinus, hexasepalus, bracteatus.

Stam. hypogyna. *Antherae* erectae, oblongae.

Styl. simplex. *Stigmata* tria.

Caps. unilocularis, trivalvis, trisperma. Sem. subglobosa, strophiole longitudinali.

Junceae. *Michel.* nov. gen. t. 31. (*Juncoides*) *Sturm.* fl. germ.

† 1412. *LINDEREA* Thunb. (1784.)

Cor. hexapetala, patens.

Stam. germiini (?) inserta, corolla breviora. *Antherae* minutae.

Styl. brevis. *Stigmata* duo. (?)

Caps. bilocularis. (?)

— — *Thunb.* fl. jap. t. 21.

1413. *ORONTIUM* L. (1754.)

Spadix multiflorus. *Spatha* nulla.

Cal. corollinus, hexasepalus, carnosus, connivens.

Filamenta brevissima, plana. *Antherae* biloculares, loculis discretis.

Styl. nullus. *Stigma* obtusum, bifidum.

Utriculus monospermus.

Aroideae. *Linn.* amoen. acad. 3. t. 1. f. 3. *Lam.* ill. t. 251.

1414. *ACORUS* Diosc.

Spadix lateralis multiflorus. *Spatha* nulla.

Cal. corollinus hexasepalus.

Stam. sepalis opposita, hypogyna. *Filamenta* subulata.

Antherae peltatae.

Styl. nullus. *Stigma* obtusum.

Caps. hexagona, trilocularis. Sem. ?

Aroideae. Schk. t. 97.

1415. PHARUS P. Brown. (1756.)

Flores monoeci, feminei spicati, masculi superiores pedicellati.

Cal. masculi floris bivalvis, basi bracteatus. Filamenta brevissima. Antherae oblongae erectae.

Cal. feminei floris bivalvis, basi subnudus, acuminatus, persistens.

Stylus tripartitus, basi persistens.

Caryopsis acuminata, calyce involuta.

Gramineae. P. Brown. jam. t. 38. f. 3. *Palis. Beauv.* agrost. t. 22. f. 8. N. Etd. 1. t. 1. f. 1—4.

B. *Fructus inferi.*

a. *Spathaceae: spatha uni-multiflora.*

a. *Perianthium duplex.*

1416. NARCISSUS Theophr.

Perianthium externum basi tubulosum; limbo regulari, sexfido, patente; internum tubulosum, limbo erectiusculo, crenulato, dentato aut lobato.

Stam. longitudine saepe inaequalia, tubo adnata. Filamenta filiformia. Antherae oblongae biloculares.

Styl. filiformis. *Stigma* trilobum, obtusum.

Caps. subrotunda, trigona, trilocularis. Sem. globosa, appendiculata.

Schk. t. 90.

a. *Hermione Salisb.* Corona laciniis corollae brevior, campanulata, lobata, plicata, crenata. Spatha tri-multiflora. Bot. mag. 15. 1687. 925. 948. 1299.

β. *Narcissus Salisb.* Corona brevis patellaris subintegra. Tubus laciniis brevior. Spatha subuniflora. Schk. t. 90. Engl. bot. 276.

γ. *Queltia Salisb.* Corona repanda, lobata, laciniis corollae brevior. Tubus lacinias aequans. Spatha unipauciflora. Bot. mag. 48. 121. 934. 945.

δ. *Ajax Salisb.* Corona subrepanda, lacinias corollae aequans vel superans. Spatha uniflora. Engl. bot. 17. Bot. mag. 6. 88. 1300.

1417. PANCRACTIUM Dalech. (1586.)

Perianthium corollinum, inferne tubulosum, limbo hexapetaloides, regulari, patente. Corona interior dentata, staminifera.

Filamenta subulata, (interdum inflexa). Antherae incumbentes, versatiles.

Styl. filiformis. Stigma obtusum.

Caps. trilocularis, trivalvis, polysperma. Sem globosa.

Bot. reg. 43. 161. 174. 221. 265. 413. 479. 600. 715. 927. 940.

[*Hymenocallis Salisb.* et *Herbert.* *Ismene Herb.* corona magna lobata, staminibus inflexis. *Liriope Herb.* flore ringente. *Leperiza* et *Carpodetes Herb.*]

418. *CALOSTEMMA* R. Br. (1810.)

Perianthium infundibuliforme sexpartitum. Corona interior tubulosa duodecim - dentata, dentibus alternis subulatis antheriferis.

Antherae versatiles.

Styl. filiformis. Stigma obtusum.

Bacca globosa, mono - disperma.

Bot. mag. 2100. 2101. Bot. reg. 421. 422.

419. *CHRYSIPIHALA* Lamb. (1822.)

Perianthium tubulosum, tubo medio constricto; limbo erecto, infundibuliformi sexfido. Corona interior abbreviata, dentata, staminifera.

Filamenta subulata, stricta. *Antherae* incumbentes.

Stylus inferne fusiformi - incrassatus. Stigma clavatum.

Caps. trilocularis, trivalvis, polysperma.

Bot. reg. 778. *Hock.* exot. fl. 2. t. 132. [*Stenomesson Herb.* in Bot. mag. 2640. *Pancratium Ruiz.* et *Pav. fl.* per. 3. t. 284. 285.]

1420. *GALANTHUS* L. (1737.)

Perianthium utrumque tripetalum; externum petalis lanceolatis patulis; internum petalis erectis, brevioribus, emarginatis.

Stam. Filamenta epigyna, brevissima. *Antherae* acuminatae, apice setaceae.

Styl. filiformis. Stigma simplex.

Caps. trigona, trilocularis: valvis medio septiferis. Sem. ovalia, solida.

Schk. t. 89. Bot. reg. 545.

β. *Perianthium simplex.*

1421. *LEVCÖIUM* Dodon. (1557.)

Perianthium campanulatum, hexasepalum, petalis apice crassiusculis.

Stam. epigyna. Filamenta abbreviata, subulata. *Antherae* oblongo - lineares.

Styl. clavatus. Stigma attenuatum. (In L. autumnali stylus filiformis.)

Caps. obovata, trilocularis, trivalvis: valvis medio septiferis. Sem. ovalia, solida.

Schk. t. 89. [παλαιοτέρων et νεωτέρων *Rengalm.* specim. p. 100. ic.]

1422. *CHLIDANTHUS* Lindl. (1821.)

Flores umbellati.

Perianthium tubuloso-infundibuliforme: tubo elongato, gracili: limbo sexfido, subirregulari.

Stam. tubo adnata, erecta. Filamenta alterna longiora, e latiore basi subulata, alterna breviora, bidentata. Antherae oblongae, erectae, adnatae.

Styl. filiformis. Stigma trilobum.

Caps. subcartilaginea, trivalvis. Sem. membranacea, marginata.

Lindl. collect. 34. Bot. reg. 640. [*Clinanthus* et *Proiphys* *Herb.* *Pancratium amboinense* Bot. mag. 1419. *Herb.* append. fig. lithograph.]

1423. *EUSTEPHIA* Cav. (1794.)

Flores umbellati.

Perianthium tubulosum, sexpartitum; laciniiis inferne conniventibus, superne in limbum patentiusculis.

Stam. basi laciniarum inserta: sterilia sex basi, exteriorum adnata. Antherae ovaes, incumbentes.

Stylus longe exsertus. Stigma simplex.

Caps.?

Cav. ic. 3. t. 238. *Phycella Lindl.* [*Amaryllis ignea* Bot. reg. 809.]

1424. *EUCROSIA* Ker. (1817.)

Flores umbellati.

Perianthium sexpartitum, ringens; lacinia infima declinata.

Stam. longissima, declinata. Filamenta subulata, basi connata. Antherae incumbentes.

Styl. declinatus. Stigma simplex.

Caps. trilocularis. Sem. duplici ordine disposita.

Bot. reg. 207. Bot. mag. 2490. *Hook.* exot. fl. 3. t. 209.

1425. *CLIVIA* Lindl. (1828.)

Flores umbellati.

Perianthium tubulosum, sexpartitum, deciduum; laciniiis omnibus conniventibus, exterioribus brevioribus.

Stam. tubo inserta, aequalia, tubum aequantia. Antherae versatiles.

Styl. filiformis. Stigma trifidum.

Bacca monosperma.

Bot. reg. 1182.

1426. *ZEPHYRANTHES* Herb. (1821.)

Spatha uni-biflora.

Perianthium infundibuliforme, erectum; limbo patente, regulariter sexfido.

Stam. tubo inserta, aequalia, erecta. Filamenta filiformia. Antherae lineares incumbentibus - erectae.

Styl. filiformis. Stigma trifidum.

Caps. triloba, trilocularis, polysperma. Sem. compressa.
Bot. reg. 821. 902. Bot. mag. 2583. 2593. 2594. [Amaryllis
caudata Bot. reg. 724.]

1427. *GRIFFINIA* Lindl. (1820.)

Spatha multiflora.

Perianthium campanulatum, subbilabiatum, laciniis limbi
patentibus subinaequalibus.

Stam. summo tubo inserta. Filamentum unum assurgens,
reliqua declinata. Antherae incumbentes.

Stylus filiformi - subulatus.

Caps. trilocularis, loculis dispermis. Sem. obovata, ven-
tricosa.

Bot. reg. 511. 990. [Amaryllis hyacinthina Bot. reg. 163.]
Folia ovali - oblonga, nervis ope venarum parallelarum
unitis.

1428. *BRUNSVIGIA* Heist. (1753.)

Spatha multiflora.

Germen longe pedicellatum, clavatum.

Perianthium sexpartitum, subringens, laciniis revolutis.

Stam. tubo inserta, declinata, exserta.

Stylus filiformis declinatus.

Caps. subdiaphano - membranacea, triquetra, trilocularis,
loculis mono - dispermis.

Bot. reg. 192. 193. 567. 954. 1153. Bot. mag. 1443. [Boophane
et Imhofia Herb. Amaryllis laticoma Bot. reg. 497. Am.
radiata ib. 596. Nerine Herb. app. p. 51. ic. Lycoris Bj.]

1429. *AMARYLLIS* L. (1737.)

Spatha pauci - multiflora.

Germen pedicellatum.

Perianthium sexpartitum, subringens. Faux squamis
(rarissime membrana) coronata, vel nuda. Limbus
patens.

Stam. tubo inserta, declinata. Antherae incumbentes.

Styl. filiformis. Stigma trifidum.

Caps. triloba, trilocularis, loculis polyspermis. Sem. com-
pressa, marginato - alata.

Bot. reg. 23. 38. 164. 172. 199. 226. 234. 426. 444. 531.
719. 849.

[Hippeastrum, Sprekelia et Habranthus Herb. in Bot. mag.
2464. 2475. 2485. 2597. 2639. Coburgia Herb. aut Ama-
ryllis reticulata Ait., semina habet globosa, baccata. Bot.
reg. 352.]

1430. *CRINUM* Theophr. L. (1737.)

Spatha multiflora.

Germen sessile vel pedicellatum.

Perianthium basi tubulosum, limbo subregulari, patente,
sexpartito, laciniis alternis apice uncinatis.

Stam. tubo inserta, recta. Filamenta subulata. Antherae
incumbentes.

Caps. trilocularis. Sem. triquetro-compressa.

Bot. reg. 52. 171. 1049. [Amaryllis longifolia 303., purpurea 552., insignis 579., revoluta 615. solandraefolia Lindl. coll. 11. Bot. reg. 876. Vallota Herb.]

1431. **HAEMANTHUS** Hermann. (1687.)

Spatha quadri-sexvalvis, multiflora, persistens.

Flores dense fasciculati.

Cor. basi tubulosa, limbo sexpartito, regulari, erecto.

Stam. tubo corollae imposita, exserta. Filamenta subulata.

Antherae incumbentes.

Styl. simplex.

Bacca trilocularis. Sem. solitaria triquetra.

Jacqu. schönb. 1. t. 56—61. vol. 4. t. 407—412. Bot. mag. 1075. 1315. 1523. Bot. reg. 181. 382. 509.

1432. **CYRTANTHUS** Ait. (1789.)

Spatha pauci-multiflora. Flores umbellati.

Cor. tubulosa, curvata, sursum ampliata, limbo erectiusculo sexfido.

Stam. tubo inserta. Filamenta filiformia. Antherae incumbentes.

Styl. exsertus. Stigma trifidum.

Caps. trilocularis. Sem. membrana laxa obvoluta.

Bot. mag. 271. 1183. Jacqu. schönb. 1. t. 75. 76. Bot. reg. 162. 167. 503. [Monella Herbert.]

1433. **HIMANTOPHYLLUM** * (Imatophyllum Hook. 1828.)

Spatha multiflora. Flores umbellati.

Corolla tubulosa, subcurvata, sexpartita, laciniis subaequalibus, apice conniventibus.

Stam. basi connata, tubo imposita, exserta. Filamenta filiformia. Antherae incumbentes.

Styl. filiformis, staminibus longior. Stigma trifidum.

Bacca globosa, trilocularis, loculis trispermis.

Bot. mag. 2856.

1434. **STERNBERGIA** Kit. (1806.)

Spatha uniflora.

Cor. sexpartita, subregularis.

Stam. basi laciniarum inserta. Antherae quadriloculares.

Styl. filiformis. Stigma trigonum.

Caps. trilocularis. Sem. strophiolata.

Kit. pl. hung. 2. t. 159. [Amaryllis lutea L. Bot. mag. 290.]

1435. **GETHYLLIS** L. (1737.)

Spatha uniflora, radicalis.

Cor. tubo gracili, elongato; limbo sexpartito, patente.

Stam. subtriadelphæ, sæpe senarium numerum excedentia, tubo corollae imposita. Filamenta subulata. Antherae elongatae, apice tortae.

Styl. filiformis. Stigma capitato-trigonum.

Bacca trilocularis, clavata, polysperma. Sem. globulosa.

Jacqu. schönbr. 1. t. 79. Bot. mag. 1068. Bot. reg. 1016.
[*Papiria Thunb. in physiogr. soc. act. 1. p. 110. a.*]

436. *STRUMARIA* Jacqu. (1790.)

Spatha pauci-multiflora. Flores umbellati.

Cor. sexpartita, patens, regularis, tubo brevissimo.

Stam. basi corollae adnata, patentia. Filamenta subulata, interdum alterna connata. Antherae oblongae, incumbentes,

Styl. basi strumosus, aut fusiformis, triquetro-sulcatus, aut subulatus. Stigma trilobum.

Caps. membranacea trivalvis, valvis medio septigeris.

Sem. duplici ordine disposita.

Jacqu. ic. rar. 2. t. 356—361. schönbr. 1. t. 71. Bot. mag. 1363. 1383. 1620. Bot. reg. 440. [Hassia Berg. in Schlechtend. Linnaea 1: p. 252.]

b. *Affines.*

437. *COERTOCAPNIA* Link. (1828.)

Flores racemosi.

Cor. tubulosa, curvata, superne ampliata, sexdentata, dentibus apice glandulosis.

Stam. basi tubi adnata. Filamenta filiformia. Antherae incumbentes.

Styl. filiformis. Stigma concavum, margine fimbriato.

Caps. trilocularis.

Link. et Ott. nov. ic. t. 18.

438. *HYPOXIS* L. (1774.)

Scapus uni-pauciflorus.

Cal. corollinus persistens, sexpartitus, regularis, patens.

Stam. disco glanduloso inserta. Filamenta brevissima. Antherae erectae, basi emarginatae.

Stylus brevissimus, crassus. Stigmata tria.

Caps. trilocularis, polysperma.

Asphodeleae. Jacqu. ic. 2. t. 307. 367. Bot. mag. 662. 710. 711. Bot. reg. 159. 663.

439. *CURCULIGO* Gärtn. (1788.)

Flores fasciculati, subspicati, bracteati.

Corollae tubus filiformis, cum stylo concretus; limbus regularis sexpartitus patens.

Stam. corollae inserta. Filamenta filiformia, recta, Antherae oblongo-lineares, adnatae.

Styl. teres. Stigma capitatum, aut trilobum.

Bacca oblonga, oligosperma. Sem. strophiole rostrato.

Asphodeleae. Roxb. corom. 1. t. 13. Jacqu. schönbr. 1. t. 80. Bot. mag. 1076. Bot. reg. 345. 754. 770. Labill. nov. Caled. t. 24.

1440. XEROPHYTA Juss. (1789.)

Flores solitarii.

Cal. corollinus tubulosus, limbo sexpartito regulari, persistente; laciniiis tribus exterioribus angustioribus, acuminatis.*Stam.* basi laciniarum inserta. Antherae elongatae, subsessiles.*Styl.* simplex. Stigma oblongum, incrassatum.*Caps.* scabra, trilocularis, polysperma.— — *Lam.* ill. t. 225.

1441. AGAVE L. (1754.)

Cal. corollinus campanulato-infundibuliformis, limbo sexpartito.*Stam.* calyci imposita, exserta. Antherae elongatae, versatiles.*Caps.* trigona, utrinque attenuata, trilocularis, polysperma.*Sem.* plana, gemino ordine disposita.— — *Lam.* ill. t. 235. *Andr. repos.* 438. *Bot. mag.* 1157. 1522. *Bot. reg.* 1145. [*Littaea Brignol.*] *Fourcroya Vent.* filamentis initio inclusis differt.

1442. ALSTRÖMERIA L. (1762.)

Cal. corollinus subcampanulatus, sexpartitus, subbilabatus. Sepala interiora angustiora, duo basi tubulosa.*Stam.* basi calycis inserta, declinata. Antherae oblongae, erectiusculae.*Styl.* declinatus. Stigma trifidum.*Caps.* subrotunda, sexcostata, trilocularis, trivalvis: valvis dissepimento contrariis. *Sem.* subglobosa.— — *Bot. mag.* 125. 139. 1613. *Hook. exot. fl.* t. 64. 65. 181. *Bot. reg.* 731. 1008.

1443. DORYANTHES Corr. (1802.)

Flores fasciculati, bracteati.

Cor. infundibuliformis, sexpartita.*Stam.* basi laciniarum inserta. Filamenta subulata. Antherae dactylothecae forma, erectae, tetragonae.*Styl.* trisulcus. Stigma trigonum.*Caps.* trilocularis, trivalvis: valvis medio septiferis. *Sem.* duplici ordine disposita.— — *Corr. in Linn. transact.* 6. t. 23. 24.

1444. CONANTHERA Ruiz et Pav. (1800.)

Cal. corollinus campanulatus, sexpartitus, subregularis, patens.*Stam.* basi corollae inserta. Filamenta brevissima, annulo iuncta. Antherae in conum conniventes, apice processu bifido, acuminato.*Styl.* subulatus.

- Caps. trilocularis, trivalvis, polysperma.*
 — — Ruiz et Pav. fl. peruv. 3. t. 304. Bot. mag. 2496.
 Hook. fl. exot. t. 214. Bot. reg. 1193. [Echeandia Orteg.
 Anthericum reflexum Cav. ic. 3. t. 241.]
445. **BARBACENIA** Vandell. (1788.)
Cal. corollinus, germi adnatus, tubulosus, extus glanduloso-pilosus; limbo erecto, sexfido.
Stam. imo tubo adnata. Filamenta brevia, plana, apice bifida. Antherae inter dentes adnatae, erectae, lineares, biloculares.
Styl. triquet. Stigma trigonum.
Caps. trilocularis, polysperma. Placentae singulorum loculorum conicae.
Haemodoreae. Vand. in Römer. script. t. 6. f. 9. Mart. nov. gen. t. 10—13. Hook. in Bot. mag. 2777.
446. **SCHWÄGRICHENIA** * (1817.)
Cal. corollinus, germi adnatus, tubulosus, extus plumoso-hispidulus; limbo sexfido; laciniis superioribus secundis.
Stam. fauci inserta, ascendentia. Antherae erectae.
Styl. filiformis.
Caps. trilocularis, polysperma, apice dehiscens.
Haemodoreae. [Anigozanthos Labill. it. t. 22. Bot. mag. 1151. Salisb. parad. 97.]
447. **CONOSTYLIS** R. Br. (1810.)
Cal. corollinus campanulatus, extus plumoso-lanatus, sexpartitus, regularis, persistens.
Stam. basi calycis inserta. Antherae erectae.
Styl. conicus, cavus. Stigma breve.
Caps. stylo coronata, apice dehiscens, polysperma. Placentae centralis, triquetra.
Haemodoreae.
448. **LANARIA** Ait. (1789.)
Cal. corollinus subcampanulatus, extus plumoso-lanatus; tubo brevi; limbo sexpartito, regulari.
Stam. basi laciniarum inserta. Filamenta filiformia. Antherae ovatae, subincumbentes.
Styl. filiformis. Stigma trifidum.
Caps. trilocularis.
Haemodoreae. [Argolasia Juss.]
449. **PHLEBOCARYA** R. Br. (1810.)
Cal. corollinus, imberbis, sexpartitus, persistens.
Stam. basi laciniarum inserta. Antherae subsessiles, tetragonae.
Styl. filiformis. Stigma simplex.
Nux. corticata, coronata, monosperma.
450. **TACCA** Rumph. (1754.)
 Flores umbellati. Involucrum polyphyllum.

Cal. corollinus, regularis, sexpartitus, persistens.

Stam. basi calycis inserta. Filamenta dilatata, superne cucullata. Antherae adnatae, biloculares, loculis discretis, apicibus solutis.

Stylus trisulcus. Stigmata tria dilatata.

Bacca polysperma.

— — *Rumph.* amboin. 5. t. 113. 114. *Bot. mag.* 1488.
Lam. ill. t. 232. *Roxb.* corom. 3. t. 257. *Gärtn.* t. 11.
 [Leontice Leontopetaloides L.]

1451. *MUSA* Dodon. (1557.)

Flores spathacei, androgyni, polygami.

Cor. dipetala: petalum alterum erectum, quinquedentatum, alterum concavum, basi nectariferum.

Stam. sex epigyna, in floribus hermaphroditis fertilibus quinque abortiunt, in floribus masculis sex perfecta adsunt, in hermaphroditis sterilibus quinque perfecta, sextum abortit. Filamenta planiuscula. Antherae adnatae, lineares, erectae.

Stylus columnaris. Stigma in fertilibus capitatum, in sterilibus bifidum.

Fructus oblongus, angulatus, baccatus, polyspermus.

Sem. albuminosa: embryo fungilli forma nidulans.

Musae. *Trew.* ic. Ehret. t. 18 — 23. *Andr.* repos. 47.
Roxb. corom. 3. t. 223. 275. 300. *Colla* in act. acad.
 taurin. 25. t. 14. *Bot. reg.* 706. *Gärtn.* t. 11. *Mirbel* in
 ann. mus. 16. t. 18.

1452. *HELICONIA* L. (1771.)

Flores spathacei hermaphroditi. Spathae distichae communes coloratae, partiales flores fasciculatos includentes.

Cor. sexpartita; exterior bilabiata, revoluta; interior convoluta.

Stam. basi corollae inserta: sextum plerumque abortiens. Filamenta subulata erecta. Antherae lineares.

Styl. triqueter. Stigma obtusum.

Caps. oblonga, trigona, trilocularis, pleiosperma. *Sem.* strophiole lanato.

Musae. *Andr.* repos. 124. *Bot. reg.* 706. *Hook.* exot.
 fl. t. 190. *Gärtn.* t. 139.

1453. *URANIA* Schreb. (1789.)

Flores spathacei, hermaphroditi. Spathae distichae.

Perianthium imbricatum, subpentasepalum. Foliola erecta, acuta, intimum bifidum, basi nectariferum.

Stam. basi perianthii inserta, exserta. Filamenta filiformia. Antherae lineares, longissimae.

Styl. filiformis. Stigma sexfidum.

Caps. triquetra, trilocularis, apice trivalvis. Sem. duplici ordine disposita, strophiole lanato.

Muscag. Ravenala Adans. Sonnerat. it ind. 2. t. 124—126.

Lam. ill. t. 222. Mirbel. in ann. mus. 16.]

D I G Y N I A.

454. SOULAMEA Lam. (1783.)

Cal. minimus, trisepalus, caducus.

Cor. tripetala: petala oblonga, concava, cum sepalis alterna.

Filamenta brevia. Antherae globulosae.

Styl. nullus. Stigmata duo obtusa.

Caps. supera, compressa, apice emarginata, bilocularis, loculis monospermis.

Polygaleae. [Rex amaroria Rumph. amb. 2. t. 41.]

455. NECTRIS Schreb. (1789.)

Cal. corollinus sexpartitus, laciniis interioribus utrinque coloratis.

Stam. basi calycis inserta. Filamenta subulata. Antherae biloculares, quadrangulae.

Styli conici, persistentes. Stigmata capitata.

Caps. duae monospermae. Sem. lenticularia.

Alismaceae? [Cabomba Aubl. guian. t. 124. Rich. in ann. mus. 17. t. 5. f. 23.]

456. OXYRIA Hill. (1772.)

Cal. tetrasepalus, sepalis basi subcohaerentibus; exterioribus marcescentibus, interioribus obovatis, post anthesin auctis.

Stam. quatuor exterioribus, duo interioribus opposita.

Filamenta brevia subulata. Antherae oblongae, biloculares.

Styli simplices. Stigmata penicillata.

Achenium compressum alatum. Embryo centralis.

Polygoneae. Campder. Rumex, t. 3. [Rumex digynus L. Engl. bot. 910. Rheum digynum Wahlb. fl. lapp. t. 9. f. 2.]

457. ATRAPHAXIS L. (1735.)

Cal. tetrasepalus, persistens; sepalis interioribus maioribus.

Filamenta capillaria, basi sepalorum inserta. Antherae subpeltatae, biloculares.

Stigmata capitata.

Achenium compressum. Embryo lateralis.

Polygoneae. Herit. stirp. 1. t. 14. Gärtn. t. 119.

458. OXYZA Theophr. Diosc.

Gramen paniculatum.

Cal. uniflorus, bivalvis, minimus.

Cor. bivalvis, persistens; valvis navicularibus; infer. dorso gibba, aristata.

Squamae hypogynae, emarginato - dentatae.

Caryopsis corolla vestita, striata.

Palis. Beauv. agrost. t. 7. f. 7.

1459. *ZIZANIA* Gronov. (1743.)

Flores monoeci, paniculati, vel (raro) subspicati.

Cal. ♂ bivalvis coloratus, valvis muticis, vel infer. aristata.

Squamae hypogynae ovatae, obtusae.

Filamenta brevissima. Antherae lineares, bifidae.

Cal. ♀ bivalvis, clausus, valva inferiore aristata, (quandoque mutica).

Squamae hypogynae ovatae, obtusae.

Caryopsis calyce vestita.

Schreb. gram. t. 29. *Palis. Beauv.* agrost. t. 22. f. 6.

1460. *POTAMOPHILA* R. Br. (1810.)

Flores monoeci, polygami.

Cal. bivalvis, uniflorus.

Cor. bivalvis, calycem excedens; valva exteriore quinque-interiore trinervia.

Squamae hypogynae duae.

T R I G Y N I A.

P A L M A E.

1461. *SABAL* Adans. (1763.)

Flores hermaphroditi sessiles. Spathae vagae.

Cal. campanulatus, tridentatus.

Cor. tripetala.

Stam. distincta.

Styli connati.

Bacca vel tri - vel monopyrena. Embryo lateralis.

[*Corypha* minor *Jacqu.* hort. vindob. 3. t. 8. *Chamaerops* acaulis *Mx.* *Rhapis* acaulis *Willd.*]

1462. *MORENIA* Ruiz et Pav. (1794.)

Flores dioeci sessiles. Spathae vagae.

Cal. trifidus.

Cor. tripetala.

Stigmata tria.

Baccae tres. Embryo basilaris.

Ruiz et Pav. prodr. t. 32.

1463. *RHAPIS* L. fil. Ait. (1789.)

Flores dioeco - polygami sessiles. Spathae vagae.

Cal. cupularis trifidus.

Cor. tripetala.

Styli tres.

Bacca monosperma.

464. **CHAMAEROPS** Plin. Dalech. (1586.)
 Flores polygami. Spatha duplex, coriacea, interior oblique aperta. Flores sessiles bracteati.
Cal. tripartitus.
Cor. tripetala.
Filamenta basi dilatata, connata.
Stigmata tria subulata.
Baccae tres monospermae. Embryo lateralis.
 [Phoenix humilis Cav. ic. 2. t. 115.]
465. **LIVISTONIA** R. Br. (1810.)
 Flores hermaphroditi.
Cal. tripartitus.
Cor. tripartita.
Styli connati.
Bacca monosperma. Embryo dorsalis.
 [Latania chinensis Jacqu. fragm. t. 11. f. 1. ?]
466. **CORYPHA** L. (1737.)
 Flores hermaphroditi, sessiles. Spathae vagae.
Cal. urceolatus, trifidus.
Cor. trifida.
Stam. distincta, corollae inserta.
Styli connati.
Bacca monosperma. Embryo basilaris.
 Mart. palm. bras. t. 49. 50. Gärtn. t. 7. Rheed. malab. 3. t. 1—12. [Saribus Rumph. amb. 1. t. 8.]
467. **PHOENIX** Theophr.
 Flores dioeci, sessiles. Spatha simplex.
Cal. tridentatus.
Cor. tripetala.
Stigmata uncinata.
Drupa monosperma. Albumen reticulatum. Embryo dorsalis.
 Kämpfer amoen. exot. p. 697. ic. Roxb. corom. 1. t. 74. vol. 3. t. 273.
468. **LEPIDOCARYON** Mart. (1824.)
 Flores dioeci, distichi, spathis duplicibus.
Cal. campanulatus, obsolete tridentatus.
Cor. tripartita.
Antherae ovato-oblongae, dorso adnatae.
Stigmata connata, linearia.
Bacca tessellato-corticata, monosperma. Albumen aequabile. Embryo lateralis, in fossa umbilicali circulari.
 Mart. palm. bras. t. 45—47.
469. **METROXYLON** Rottböll. (1788.)
 Flores androgyni, in amentis compressis distichi. Spathae duplices.
Cal. campanulatus, subtridentatus.
Cor. tripartita.

Antherae ovato-oblongae, erectae.

Stigmata acuta, connata.

Bacca squamis tessellatis oorticata, monosperma. Albumen ruminatum. Embryo lateralis, super fossa umbilicali.

Ann. of bot. 1. t. 4. [Sagus Rumph. Gärtn. t. 10. Raphia Pal. Beauv. fl. ovar. t. 44—46.]

1470. *HYPHAENE* Gärtn. (1791.)

Flores dioeci, radiatum in amentis dispositi, spathis setarumque fasciculis interstincti,

Cal. tripartitus.

Cor. tripetala.

Stigmata sessilia.

Drupa mono- (raro tri-) sperma; putamine coriaceo, apice pervio. Albumen aequabile, cavum. Embryo verticalis.

Gärtn. t. 82. [Cucifera Delil. flor. aegypt. t. 1. 2.]

1471. *LEOPOLDINIA* Mart. (1824.)

Flores androgyni in spadice ramosissimo sessiles, bracteati. Spathae nullae.

Cal. trisepalus, imbricatus.

Cor. tripetala.

Stigmata sessilia.

Bacca drupaeformis, sicca, e fibris reticulatis, monosperma. Albumen aequabile. Embryo subbasilaris.

Mart. palm. bras. t. 52. 53.

1472. *HYOSPATHE* Mart. (1824.)

Flores androgyni, sessiles, ebracteati. Spatha duplex.

Cal. tripartitus.

Cor. tripetala.

Stigmata sessilia.

Bacca monosperma. Albumen aequabile. Embryo basilaris.

Mart. palm. bras. t. 1. 2.

1473. *ABECA* Raj. (1688.)

Flores sessiles, in eodem spadice, superiores masculi, inferiores feminei. Spatha duplex, membranacea.

Cal. tripartitus.

Cor. tripetala.

Stigmata sessilia.

Bacca vel drupa fibrosa, monosperma. Albumen ruminatum. Embryo basilaris.

Gärtn. t. 7. Roxb. corom. 1. t. 75. [Pinanga Rumph. amb. 1. t. 4.]

1474. *OENOCARPUS* Mart. (1824.)

Flores sessiles ebracteati, androgyni. Spatha duplex, lignosa.

Cal. tripartitus.

Cor. tripetala.

Stigmata tria.

Bacca grumosa, fibrosa, monosperma. Albumen ruminatum. Embryo basilaris.

Mart. palm. bras. t. 22 — 27. [Oreodoxa W. et Kunth.]

475. EUTERPE Gärtn. (1788.)

Flores androgyni in scrobiculis spadiciis sessiles, bracteati.

Spathae duae membranaceae.

Cal. trisepalus.

Cor. tripetala.

Stigmata tria.

Bacca grumosa, fibrosa, monosperma. Albumen ruminatum. Embryo lateralis.

Gärtn. t. 9. Mart. palm. bras. t. 28 — 32. [Areca oleracea Jacq. Alphaeus Humb. Martinezia ensifolia R. et P. Pinauga Rumph. amb. 1. t. 5.]

476. WALLICHIA Roxb. (1819.)

Flores androgyni sessiles. Spathae perfectae plures.

Cal. campanulatus, tridentatus, in ♀ bracteis binis suffultus.

Cor. tripetala.

Stigmata sessilia.

Bacca disperma. Albumen aequabile. Embryo lateralis.

Roxb. corom. 3. t. 295.

1477. GULIELMA Mart. (1824.)

Flores androgyni, sessiles, bracteolati. Spatha duplex.

Cal. ♂ trifidus, ♀ annuliformis.

Cor. ♂ tripetala, ♀ campanulata.

Stigmata sessilia.

Drupa, cuius putamen in vertice poros tres habet stellatos. Albumen aequabile. Embryo intra porum.

Mart. palm. bras. t. 66. 67. [Palma Pirijao Humb. nov. gen. 1. p. 315.]

1478. ELAEIS Jacqu. (1763.)

Flores monoeci, foveolis immersi. Spatha duplex.

Cal. trisepalus.

Cor. tripetala.

Stam. basi in urceolum connata.

Stigmata sessilia, aut styli tres, aut stylus triqueter, stigmatibus tribus.

Drupa oleosa, cuius putamen in vertice tres poros habet.

Albumen aequabile. Embryo intra porum.

Jacqu. amer. t. 172. Gärtn. t. 6. Mart. palm. bras. t. 54 — 56. Alphonsia Humb.]

1479. SYAGRUS Mart. (1824.)

Flores androgyni sessiles. Spatha duplex.

Cal. trisepalus.

Cor. tripetala.

Stigmata sessilia.

Drupa fibrosa, putamine basi poro unico pervio, duorumque rudimentis. Albumen aequabile. Embryo basilaris, intra porum.

Mart. palm. brasil. t. 89. 90.

1480. *Cocos* L. (1764 *Coccus* 1737.)

Flores androgyni sessiles. Spatha simplex.

Cal. trisepalus.

Cor. tripetala.

Stigmata sessilia.

Drupa fibrosa, putamine basi tribus poris instructo. Albumen aequabile, cavum. Embryo intra porum, basilaris.

Gärtn. t. 4—6. Roxb. corom. 1. t. 73. Mart. palm. bras. t. 78—88. [Calappa Rumph. amb. 1. t. 12.]

1481. *MAXIMILIANA* Mart. (1824.)

Flores androgyni sessiles. Spatha simplex.

Cal. trisepalus.

Cor. tripetala.

Stigmata tria revoluta.

Drupa monosperma, putamine poris tribus lateralibus. Albumen aequabile. Embryo basilaris, intra porum.

Mart. palm. bras. t. 91—93.

1482. *MARTINEZIA* Ruiz et Pav. (1794.)

Flores androgyni sessiles.

Cal. trisepalus.

Cor. tripetala.

In floribus ♀ cylindrus membranaceus subsexdentatus, ovarium ambiens (stamina abortientia?)

Stigmata sessilia.

Drupa monosperma. Albumen aequabile. Embryo?

Ruiz et Pav. prodr. t. 32.

MELANTHACEAE. R. Br.

1483. *MELANTHIUM* Gronov. (1743.)

Cal. corollinus hexasepalus. Sepala unguiculata, ad unguem biappendiculata subcucullata, aut foveis duabus nectariferis.

Stam. unguibus inserta. Antherae extrorsum dehiscentes.

Styli filiformes.

Caps. obovata, tripartibilis, trilocularis, trivalvis, loculis apice saepe cornutis, inter cornu et angulum stylumque dehiscentibus. Semina subtrigona, angulo interiori dissepimentorum affixa.

Bot. mag. 558. Thunb. in act. soc. hist. nat. havn. 4. t. 8. [Leimanthium Willd.]

1484. *KOLBEA* Schlechtend. (1826.)

Cal. corollinus, hexasepalus, campanulatus. Sepala exunguia, exappendiculata.

Stam. basi sepalorum inserta.

Styli nulli. Stigmata tria revoluta, persistentia in angulis fructus.

Caps. prismatica, trilocularis, tripartibilis, trivalvis, inter cornua et columnam mediam superne et interne dehiscentis. Sem. duplici serie angulo interno loculorum affixa.

[*Sisyrinchium Breyn.* cent. t. 36. *Tulipa Breyniana L.* *Melanthium uniflorum Jacqu.* ic. rar. 2. t. 450. *Bot. mag.* 767.]

485. *ANDROCYMBIUM* Willd. (1808.)

Cal. corollinus, hexasepalus. Sepala longissime unguiculata, convoluta-cucullata, nectarifera.

Stam. cucullis sepalorum inserta. Antherae incumbentes, extrorsum dehiscentes.

Styli loculos germinis discretos terminantes.

Caps. tripartibilis, trilocularis, trivalvis: loculi polyspermi. Sem. duplici serie angulo interiori dissepimentorum affixa.

Willd. in *Berl. mag.* 2. t. 2. [*Melanthium capense L.* *M. eucomoides Jacqu.* ic. rar. 2. t. 452. *Bot. mag.* 644.]

486. *OENITHOGLOSSUM* Salisb. (1806.)

Cal. corollinus hexasepalus, patens. Sepala subunguiculata, medio foveola nectarifera.

Stam. basi sepalorum inserta. Antherae extrorsum dehiscentes.

Styli tres in medio germine, basi leviter uniti.

Caps. trilocularis, trivalvis: valvis medio septiferis. Sem. angulato-globosa, grandia.

Salisb. parad. 54. [*Lichtensteinia Willd.* in *Berl. mag.* 2. t. 1. *Cymation Syst. veg.* *Melanthium viride L.* suppl. *Andr. repos.* t. 233.]

487. *WURMBEA* Thunb. (1781.)

Cal. corollinus monosepalus, campanulatus, sexfidus.

Stam. basi laciniarum calycis inserta. Antherae extrorsum dehiscentes.

Styli tres germinis loculos terminantes.

Caps. tripartibilis, trilocularis, tricornis. Sem. subglobosa.

Bot. mag. 694. 1291. *Lam.* ill. t. 270.

1488. *ANGUILLARIA* R. Br. (1810.)

Cor. hexapetala, stellato-patens, regularis.

Stam. basi petalorum inserta. Antherae extrorsum dehiscentes.

Styli tres.

Caps. trilocularis, trivalvis, valvis medio septiferis. Sem. numerosa, subglobosa.

[*Melanthium indicum L.*]

S A R M E N T A C E A E.

1499. FLAGELLARIA L. (1747.)

Cal. corollinus, inferus, persistens, sexpartitus.

Stam. hypogyna. Filamenta subulata. Antherae oblongae.

Styli filiformes. Stigmata lateralialia.

Drupa pisiformis, monosperma. Embryo lenticularis, foveolae baseos albuminis farinacei semiimmersus.

Red. lil. 257. *Lam.* ill. t. 266.

1500. MEDEOLA Gronov. (1743.)

Cal. corollinus, hexasepalus, revolutus.

Stam. basi calycis inserta. Filamenta subulata. Antherae incumbentes.

Styli divaricati.

Bacca trilocularis, loculis subhexaspermis. Sem. trigona.

Bot. Mag. 1316. [*Gyromia Nuttall*, hallucinatione ortum nomen, quod crederet N., *Medeolae* characterem primitus fuisse ad africanas plantas formatum, quas *Tirtnaeus* post huic generi ascripsit. Sed *Medeolam americanam* a *Claytoneo* acceptam *Gronovius* triginta annos antequam *M. asparagoides Linnaeo* diceretur, descripserat.]

1501. MYRSIPHYLLUM Willd. (1808.)

Cal. corollinus, hexasepalus, revolutus.

Filamenta subulata, sepalorum basi inserta.

Styli stricti contigui.

Germen pedicellatum. *Bacca* trilocularis, loculis dispermis.

[*Laurus alexandrina Herm.* hort. lugd. bat. p. 681. ic. *Asparagus africanus Till.* pis. t. 12. f. 1. *Medeola asparagoides L.* mant.]

1502. VERATRUM Plin.

Flores polygami.

Cal. corollinus, hexasepalus, patens.

Stam. receptaculo inserta.

Styli tres breves.

Caps. tres, bivalves, polyspermae. Sem. alata.

Schk. t. 341. *Gärtn.* t. 18.

1503. TRILLIUM Mill. (1739.)

Cal. trisepalus, persistens.

Cor. tripetala, calyce maior, aut longior, aut eum subaequans.

Stam. hypogyna. Filamenta abbreviata. Antherae oblongo-lineares, elongatae, erectae.

Styli revoluti.

Bacca trilocularis, loculis polyspermis.

Salisb. parad. lond. 1. *Willd.* hort. berol. t. 35. *Bot. mag.* 40. 470. 954.

A L I S M E A E.

1504. TRIGLOCHIN Dalech. (1587.) Rupp. (1718.) L. (1737.)

Cal. hexasepalus, duplici ordine: sepala ovata concava, decidua.

Stam. tria exteriora inter sepala ex- et interiora, tria interiora. Filamenta brevissima. Antherae cordato-subrotundae, ambitu circulari dehiscentes.

Styl. recurvi vel nulli. Stigmata penicillata.

Caps. tres, aut sex, evalves, monospermae. Sem. triquetrum, utrinque attenuatum.

Schk. t. 102. *Gärtn.* t. 84. [Tetroncium Willd. in Berl. mag. 2. p. 17. floribus dioecis et capsulis quaternis distinguitur. Styli et persistunt numeroque variant et in aliis speciebus.]

505. HYDROMYSTRIA G. F. W. Meyer. (1818.)

Flores polygami.

Cal. trisepalus, patens.

Cor. tripetala, patens. In femineis dicitur perianthium tubulosum, tripartitum. Squamae tres, petalorum rudimenta.

Stam. Filamenta brevissima. Antherae lanceolatae, elongatae.

Styli apicibus reflexis.

Caps. unilocularis.

506. HYDROGETON Pers. (1805.)

Cal. corollinus trisepalus.

Stam. Filamenta basi dilatata. Antherae terminales.

Styli nulli.

Folliculi tres dispermi. Albumen nullum.

[Ouvirandra Aub. Petit. Thouars nov. gen. madag. n. 3.]

D I O S C O R E A E.

507. DIOSCOREA Plum. (1703.)

Flores dioeci.

Perianthium superum, sexpartitum, persistens.

Stam. basi perianthii inserta. Filamenta subulata. Antherae biloculares.

Styli simplices.

Caps. triangularis, trilocularis; loculis dispermis; valvarum angulis prominentibus dehiscentibus. Sem. compressa, alata.

Salisb. parad. 17. *Gärtn.* t. 17. *Hook.* in Bot. mag. 2825.

[Testudinaria *Salisb.* *Burch.* in Bot. reg. 924. Fructuum maturorum, quos Zeyherus ex Africa australi misit, eandem omnino fabricam ac Dioscoreae, conspicio.]

508. RAJANIA L. (Jan-Raja Plum. 1703.)

Flores dioeci.

Perianthium superum, campanulatum, sexpartitum.

Stam. basi perianthii inserta. Filamenta subulata. Antherae biloculares.

Styli stigmatibus obtusis.

HEXAGYNIA

1518. *DAMASONIUM* Schreb. (1789.)

Spatha monophylla, quinquealata; alis undulatis, duabus in petiolum decurrentibus; ore quinquedentato.

Cal. superus, tripartitus, patens.

Petala tria patentia. Glandulae tres nectariferae.

Filamenta brevissima. Antherae lineares, erectae.

Styli lineares erecti.

Bacca sexvalvis polysperma, valvis placentiferis.

Roxb. coron. 2. t. 186. [Ottelia Pers. Stratiotes alimodes L.]

1519. *IALAMBICEA* Lexarz. (1825.)

Flores monoeci. *Spatha* diphylla.

Cor. ♂ hexa - ♀ tripetala.

Filamenta brevissima. Antherae oblongae, quadrangulares.

Styli sex bipartiti.

Caps. infera, unilocularis, polysperma.

POLYGYNIA

1520. *ALISMA* Diosc.

Cal. trisepalus.

Cor. tripetala.

Stam. hypogyna, gemina petalis opposita. *Filamenta* subulata. Antherae incumbentes.

Styli seni, octoni aut plures brevissimi. *Stigmata* lateralia.

Caps. submonospermae, acuminatae, stellatae. Embryo complicatus.

Scht. t. 102. Engl. bot. 1615. [Damasonium Juss. Actinocarpus R. Br.]

CLASSIS VII. HEPTANDRIA.

CONSPECTUS

CLASSIS SEPTIMAE.

MONOGYNAE.

Floribus perfectis 1521 — 1525.

Floribus imperfectis 1526.

DIGYNA.

1527.

HEPTAGYNA.

1528.

MONOGYNIA.

I. FLORES PERFECTI.

21. TRIENTALIS Rupp. (1718. *Herba trientalis* Val. Cord. 1561.)

Cal. septempartitus.

Cor. patens septempartita, laciniis ovato-lanceolatis.

Stam. Filamenta subulata, basi corollae inserta. Antherae ovaes.

Styl. simplex. Stigma capitatum.

Bacca capsularis, sicca. Placenta centralis libera, subglobosa, excavata. Sem. piura angulata.

Primuleae. Schk. t. 103. Sturm. fl. germ.

2. AESCULUS L. (*Esculus* 1735.)

Cal. campanulatus, quinquedentatus.

Pet. quatuor, quinque, unguiculata, inaequalia.

Stam. disco perigyno inserta, declinata. Filamenta subulata. Antherae ovato-oblongae.

Styl. conicus. Stigma tridentatum.

Caps. coriacea, bi-trilocularis, oligosperma. Cotyledones crassissimae.

Hippocastaneae. Schk. t. 104.

3. ERYTHROSPERMUM Lam. (1791.) Poir. (1811.)

Cal. tetrasepalus, deciduus.

Pet. septem concava.

Stam. perigyna. Filamentum brevissima, subnulla. Antherae oblongae, antherae.

HEXAGYNIA

1518. *DAMASONIUM* Schreb. (1789.)

Spatha monophylla, quinquealata; alis undulatis, duabus in petiolum decurrentibus; ore quinquedentato.

Cal. superus, tripartitus, patens.

Petala tria patentia. Glandulae tres nectariferae.

Filamenta brevissima. Antherae lineares, erectae.

Styli lineares erecti.

Bacca sexvalvis polysperma, valvis placentiferis.

Roxb. corom. 2. t. 186. [Ottelia Pers. Stratiotes alismoides L.]

1519. *IALAMBICEA* Lexarz. (1825.)

Flores monoeci. *Spatha* diphylla.

Cor. ♂ hexa - ♀ tripetala.

Filamenta brevissima. Antherae oblongae, quadrangulares.

Styli sex bipartiti.

Caps. infera, unilocularis, polysperma.

POLYGYNIA

1520. *ALISMA* Diosc.

Cal. trisepalus.

Cor. tripetala.

Stam. hypogyna, gemina petalis opposita. *Filamenta* subulata. Antherae incumbentes.

Styli seni, octoni aut plures brevissimi. *Stigmata* lateralia.

Caps. submonospermae, acuminatae, stellatae. Embryo complicatus.

Schk. t. 102. Engl. bot. 1615. [Damasonium Juss. Actinocarpus R. Br.]

CLASSIS VII.
H E P T A N D R I A.

CONSPECTUS

CLASSIS SEPTIMAE.

M O N O G Y N A E.

l. Floribus perfectis 1521 — 1525.

I. Floribus imperfectis 1526.

D I G Y N A.

1527.

H E P T A G Y N A.

1528.

M O N O G Y N I A.

I. FLORES PERFECTI.

521. TRIENTALIS Rupp. (1718. *Herba trientalis* Val. Cord. 1561.)

Cal. septempartitus.

Cor. patens septempartita, laciniis ovato-lanceolatis.

Stam. Filamenta subulata, basi corollae inserta. *Antherae* ovoides.

Styl. simplex. Stigma capitatum.

Bacca capsularis, sicca. Placenta centralis libera, subglobosa, excavata. Sem. plura angulata.

Primulaeae. Schk. t. 103. *Sturm.* fl. germ.

522. AESCULUS L. (*Esculus* 1735.)

Cal. campanulatus, quinquedentatus.

Pet. quatuor, quinque, unguiculata, inaequalia.

Stam. disco perigyno inserta, declinata. Filamenta subulata. *Antherae* ovato-oblongae.

Styl. conicus. Stigma tridentatum.

Caps. coriacea, bi-trilocularis, oligosperma. Cotyledones crassissimae.

Hippocastaneae. Schk. t. 104.

523. ERYTHROSPERMUM Lam. (1791.) Poir. (1811.)

Cal. tetrasepalus, deciduus.

Pet. septem concava.

Stam. perigyna. Filamenta brevissima, subnulla. *Antherae* oblongae, petala subaequantes.

Styl. simplex. Stigma tri-quinquelobum.

Caps. globosa, unilocularis, pleiosperma.

Flacourtiae Rich. Cand. Lam. ill. t. 274.

1524. PETROCARYA Schreb. (1789.)

Cal. urceolatus, quinquefidus.

Pet. quinque, ovata, calyci inter lacinias inserta.

Stam. Filamenta quatuordecim, capillaria, calycis ori infra petala inserta; quorum septem antherifera, septem in opposita calycis parte sterilia. Antherae subrotundae, introrsum dehiscentes.

Styl. incurvus. Stigma capitatum.

Drupa magna fibrosa: nuce lapidea, biloculari. Sem lanugine obducta.

Chrysobalanaceae R. Br. [Parinari Aubl. guian. t. 204.]

1525. TOVARIA Ruiz et Pav. (1800.)

Cal. heptaphyllus, bracteatus.

Pet. septem, basi villosa.

Stam. pubescentia, disco heptagono inserta.

Stylus persistens. Stigma peltatum.

Bacca pisiformis, pulposa, unilocularis. Sem. compressa.

Melieae. Ruiz et Pav. fl. peruv. 3. t. 306.

II. FLORES IMPERFECTI.

1526. PISONIA Plum. (1703.)

Flores subdioeci.

Cal. campanulatus, persistens, coloratus, limbo plicato, quinqueidentato aut quinquefido.

Stam. plerumque septena, interdum dena, calycis tubo adnata, exserta. Filamenta capillaria. Antherae subrotundae.

Stylus simplex. Stigma bi-multifidum.

Utriculus monospermus, calyce angulato inclusus. Album centrale.

Nyctagineae. Plum. ic. ed. Burm. t. 227. f. 1. Gårds t. 76. [Calpidia Thuars. hist. plant. ins. Afric. t. 8.]

D I G Y N I A.

1527. LIMEUM L. (1759.)

Cal. pentasepalus, duplici serie; sepalis margine membranaceis.

Pet. quinque, ovata, obtusa, subunguiculata, calyce breviora.

Stam. urceolo perigyno inserta. Filamenta basi dilatata, subulata. Antherae ovaes.

Styli staminibus breviores. Stigmata capitata.

Sem. duo hemisphaerica, scabra, hinc excavata. Embryo
periphericus.

Chenopodeae. Gärtn. t. 76.

H E P T A G Y N I A.

528. GILIBERTIA Ruiz et Pav. (1800.)

Cal. superus brevis, septemdentatus.

Pet. septem patentia.

Stam. perigyna. Filamenta subulata. Antherae ovales.

Styli nulli. Stigmata ovalia.

Caps. ovalis, septemlocularis, loculis monospermis.

Araliace. Ruiz et Pav. fl. peruv. 3. t. 312.

CLASSIS VIII. OCTANDRIA.

CONSPECTUS

CLASSIS OCTAVAE.

MONOGYNAE.

- I. Floribus perfectis.
 - A. Corolla monopetala.
 - a. Fructibus superis. 1529 — 1543.
 - b. Fructibus inferis. 1544, 1545.
 - B. Corolla pleiopetala.
 - a. Fructibus superis.
 - α. Corolla tetrapetala. 1546 — 1586.
 - β. Corolla pentapetala. 1587 — 1602.
 - γ. Corolla octopetala. 1603.
 - b. Fructibus inferis. 1604 — 1614.
- II. Floribus imperfectis.
 - Fructibus superis. 1615 — 1632.

DIGYNAE.

1633 — 1638.

TRIGYNAE.

1639 — 1649.

TETRAGYNAE.

1650 — 1663.

MONOGYNIA.

I. Flores perfecti.

A. COROLLA MONOPETALA.

a. Fructus superi.

1529. ERICA Theophr.

Cal. tetrasepalus, persistens.

Cor. monopetala, limbo quadrifido; tubulosa, infundibuliformis, campanulata, ampullacea, urceolata, vel globulosa; ut plurimum calycem excedens, interdum eo minor.

Stam. inclusa vel exserta. Filamenta capillaria, perigyna, receptaculo inserta. Antherae ante anthesin foraminulis duobus lateralibus nexae; basi vel muticae, vel appendiculis aristatis aut cristatis instructae.

Stylus simplex. Stigma capitatum, vel peltatum, vel quadrifidum.

Caps. quadri - octolocularis: dissepimentis ad placentam centralem fixis: valvis totidem. Sem. minuta.

Ericaceae. Schk. t. 107. a. Lam. ill. t. 287. [Calluna Salisb.]

1. *Tubiflorae*, corollis tubulosis, apice dilatatis.
2. *Coniflorae*, corollis cylindricis, apice constrictis.
3. *Ampullaceae*, corollis oblongis, fauce constricta, limbo patente.
4. *Urceolatae*, corollis ovato-subglobosis, basi ventricosis, ore contracto, limbo patulo.
5. *Campanulatae*, corollis sensim in limbum dilatatis.
6. *Calycinae*, calycibus corollas aequantibus vel superantibus.

1530. *SALAXIS* Salisb. (1802.)

Cal. tetrasepalus irregularis.

Cor. campanulata, quadrifida.

Stigma peltatum dilatatum.

Caps. drupacea, trilocularis, trisperma.

1531. *MENZIESIA* Sm. (1789.)

Cal. monophyllus quadridentatus.

Cor. tubulosa aut ovata, quadrifida.

Stam. receptaculo inserta. Antherae bifidae, muticae.

Styl. simplex. Stigma capitatum.

Caps. quadrilocularis, marginibus valvarum inflexis, dissepimenta sistentibus.

Ericaceae. Smith. ic. ined. 1. t. 56. Salisb. parad. lond. t. 44. Engl. bot. 2469. [Erica Dabeoci Engl. bot. 35. E. Bruckenthalii Reichenb. ic. 2. t. 169.]

† 1532. *ELLIOTTIA* Mühlenb. (1813.)

Cal. quadridentatus.

Cor. quadripartita, subtetrapetala.

Stam. hypogyna. Filamenta glandulosa. Antherae sagittatae, biloculares.

Styl. elongatus. Stigma subclavatum.

Caps. ?

Ericaceae?

1533. *DIOSPYROS* Theophr. Dalech. (1587.)

Flores polygami.

Cal. quadripartitus (quandoque tri- aut sexpartitus).

Cor. urceolata, quadrifida, laciniis revolutis.

Stam. basi corollae inserta. Filamenta brevissima, quandoque alterna biantherifera. Antherae basi affixae, subulatae.

Styl. quadrifidus, interdum bipartitus, saepe trifidus.

Bacca globosa, octolocularis, calyce persistente suffulta.

Sem. solitaria, compressa, albuminosa. Embryo obliquus.

Styracaceae. Roxb. corom. 1. t. 47. 50. Ann. mus. 6. t. 62.

Hook. exot. 2. t. 139. Bot. reg. 499. 1139. Gärtn. t. 207.

[Embryopteris Gärtn. t. 29. Roxb. corom. 1. t. 70. Cavanilla Lam.]

1534. *CARGILLIA* R. Br. (1810.)

Flores polygami.

Cal. semiquadrifidus, fructifer cupulaeformis.*Cor.* urceolata, limbo quadrifido.*Stam.* basi corollae inserta.*Germen* quadriloculare, loculis dispermis. *Bacca* globosa.*Styraceae.*1535. *MIMUSOPS* L. (1747.)*Cal.* sex-octopartitus, duplici laciniarum serie.*Cor.* laciniis totidem, duplici ordine; interioribus integris, exterioribus saepe divisis.*Stam.* corollae inserta: sterilia cum fertilibus alternantia.*Antherae* sagittatae.*Bacca* oligosperma.*Sapoteae.* Roxb. corom. 1. t. 14. Lam. ill. t. 300. [Imbricaria Lam. ill. t. 300. Gärtn. t. 206.]1536. *NOLTIA* Schumach. (1827.)

Flores polygami.

Cal. quadrifidus, tomentosus.*Cor.* tubulosa, coriacea, limbo erecto quadrifido.*Stam.* receptaculo inserta. Filamenta inaequalia. *Antherae* subulatae.*Styl.* acutus.*Bacca* coriacea, unilocularis, tetrasperma.1537. *SYMPHOREMA* Roxb. (1815.)*Involucrum* sex-octophyllum, sex-octoflorum: foliolis post anthesin mutatis, membranaceis, venoso-reticulatis.*Cal.* campanulatus, sex-octodentatus.*Cor.* tubulosa, limbo sex-octopartito reflexo.*Stam.* corollae inserta, exserta. Filamenta filiformia, flexuosa. *Antherae* incumbentes.*Styl.* elongatus. Stigma bifidum.*Caryopsis* solitaria, calyce inclusa.1538. *LENNOA* Lexarz. (1824.)*Cal.* profunde octopartitus; laciniis spathulato-linearibus, nervosis.*Cor.* calycem aequans, basi subventricosa, tubulosa; limbo plicato, octodentato.*Stam.* tubo corollae duplici serie inserta, inclusa. Filamenta subulata. *Antherae* bilobae, incumbentes.*Styl.* subulatus. Stigma truncatum.*Caps.* pyriformis, unilocularis, octovalvis, polysperma.

— — [Corallophyllum Kunth. in Humb. nov. gen. 7. t. 660. b.]

1539. *PALLADIA* Lam. (1792.)*Cal.* infundibuliformis, coloratus; limbo quadrifido; laciniis obovatis.

Cor. infundibuliformis, tubo elongato, octoplicato; limbo octofido, laciniis oblongis.

Stam. corollae adnata. Filamenta rigida, persistentia.

Germina duo. Stylus compressus, inter germina positus, ad margines denticulatus. Stigmata duo patula.

Caps. duae, prismaticae, uniloculares, bivalves, valvis post dehiscentiam funis ad instar se contorquentibus.

Placenta fungosa. Sem. numerosissima, angulata.

Gentianeae? Savigny. [Blakwellia Gärtn. t. 117.]

540. *CHLORA* Renealm. (1611.)

Cal. octo- (sex- quadri-) partitus; laciniis lanceolatis acuminatis.

Cor. tubo brevi; limbo patente, octo- (sex- quadri-) fido.

Stam. corollae inserta. Filamenta brevissima. Antherae lineares, erectae.

Styl. filiformis. Stigma bilobum.

Caps. unilocularis, bivalvis; valvis inflexis placentiferis.

Sem. numerosa.

Gentianeae. Reichenb. ic. 3. t. 206 — 209. vol. 5. t. 412.

541. *HEDWIGIA* Swartz. (1788.)

Cal. cyathiformis, quadridentatus.

Cor. tubulosa; limbo quadrifido, erecto, subconnivente.

Stam. basi corollae inserta. Filamenta e basi latiore subulata. Antherae oblongae, acutae, circa stigma conniventes.

Stylus nullus. Stigma obtusum.

Caps. trilocularis, succo gummoso repleta. Nuclei tres ovatae acutae, hinc convexae. Nuclei amari, oleosi.

— — Swartz. fl. ind. occ. t. 13.

542. *MIMOSA* Acost. (1578.) Adans.

Flores polygami.

Cal. suburceolatus, quadri- quinquedentatus; (interdum laciniatus, nonnunquam subinteger.)

Cor. campanulata, persistens; limbo quadri- quinquefido, regulari.

Stam. (numero varia) basi corollae vel gynobasi inserta, longe-exstantia. Filamenta capillaria. Antherae suborbiculatae, dorso affixae, biloculares.

Stylus stigmatate simplici.

Lomentum planum, polyspermum.

Mimoseae. R. Br. Bot. reg. 25. 941.

543. *SCHRANKIA* Willd. (1808.)

Flores polygami.

Cal. urceolatus, minutus, quinquedentatus.

Cor. infundibuliformis; limbo quinquefido regulari.

Stam. octo vel decem, basi corollae vel gynobasi inserta, exstantia. Antherae oblongae, biloculares.

Stylus simplex.

Legumen tetragonum, exsuccum, bivalve, polyspermum; valvis bipartilibus.

Mimoseae. [Mimosa quadrivalvis Banks. rel. *Houst.* t. 25.
M. horridula Vent. sel. t. 28.]

b. *Fructus inferi*.

1544. *VACCINIUM* Virg. Plin.

Cal. quadri-quinquefidus (vel quadri-quinquedentatus, interdum integer).

Cor. urceolata, ore constricto; vel campanulata patens; limbo quadrifido, (quadri-) quinquedentato.

Stam. calycis limbo imposita. Antherae bifidae, apice poro gemino dehiscentes; dorso saepe appendiculis aristatis instructae.

Stylus simplex. Stigma obtusum.

Bacca quadri-quinquelocularis, polysperma.

Ericaceae. Schk. t. 107. a. Bot. reg. 302. 400. *Humb.* nov. gen. 3. t. 248 — 253.

1545. *MICHAUXIA* Herit. (1788.)

Cal. octo- (sexdecim-) fidus, persistens.

Cor. rotata, tubo brevissimo; limbo octopartito, lacinis patentissimis apice revolutis.

Filamenta e basi dilatata, fornicata in apicem subulatum abeuntia. Antherae lineares elongatae, stylo appressae.

Stylus staminibus longior, pubescens. Stigma octopartitum, lacinis revolutis.

Caps. octocostata, octolocularis. Placentae ad angulos loculorum.

Campanuleae. Lam. ill. t. 295. Venten. Cels. t. 81. Gärtn. t. 211.

B. COROLLA PLEIOPETALA.

a. *Fructus superi*.

α. *Corolla tetrapetala*.

SALICARIAE.

1546. *ADENARIA* Bonpl. Kunth. (1823.)

Cal. turbinato-campanulatus, limbo aequaliter quadri- aut quinquefido.

Pet. quatuor aut quinque, inter lobos calycis inserta, aequalia, unguiculata.

Stam. octo vel decem, calyci supra basin inserta, exstantia. Filamenta libera. Antherae biloculares dorso affixae.

Styl. simplex. Stigma bilobum. Germen stipitatum.

Caps. unilocularis, polysperma. Placenta centralis globosa.

Humb. nov. gen. 6. t. 549.

547. *GRISLEA* L. (1737.)

Cal. tubuloso - campanulatus, coloratus, limbo octo - decedentato, dentibus alternis minoribus.

Pet. quatuor aut quinque, calyci inter dentes imposita, aequalia, unguiculata.

Stam. octo aut decem, calyci supra basin inserta, exserta. Filamenta libera. Antherae biloculares, dorso affixae.

Stylus exsertus. Stigma simplex. Germen sessile.

Caps. membranacea, polysperma, unilocularis, non dehiscentis. Placenta centralis.

Bot. reg. 30. *Roxb. corom.* 1. t. 31. [*Woodfordia Salisb. parad.* t. 42.]

548. *LAWSONIA* L. (1747.)

Cal. quadripartitus, persistens.

Pet. quatuor, aequalia, unguiculata, patentia.

Stam. cum petalis alterna, per paria approximata.

Styl. simplex. Stigma capitatum.

Caps. baccata, quadrilocularis, polysperma. Sem. angulata.

Lam. ill. t. 296.

549. *ACRONYCHIA* Forst. (1775.)

Cal. quadrilobus, persistens.

Petala perigyna, lanceolata, erecta, apice inflexo.

Stam. perigyna. Filamenta basi dilatata, persistentia.

Antherae subpeltatae, biloculares, dorso affixae.

Discus perigynus octocrenatus, glandulosus.

Styl. simplex, stigmatibus quadrifido.

Caps. quadrilocularis; loculis mono - dispermis. Sem. strophilolata.

Forst. char. 27. *Labill. nov. Caled.* t. 65. [*Lawsonia Acronychia* L. fil.]

R U T A C E A E.

550. *BORONIA* Sm. (1798.)

Cal. quadripartitus aut quadrifidus.

Pet. quatuor longiora, marcescentia.

Stam. quatuor breviora petalis opposita, interdum alterna abortientia. Filamenta perigyna, incurva; apice dilatato aut incrassato. Antherae subcordatae, dorso affixae.

Styl. vel simplex, stigmatibus capitato, vel e quatuor conflatus.

Caps. quadriloba, tetracocca. Cocci dispermi.

Smith. tracts, t. 4 — 7. *Labill. nov. holl.* t. 124 — 126.

Reichenb. ic. t. 74 — 74. *Mém. mus.* 12. t. 22. f. 26.

551. *CORREA* Sm. (1798.)

Cal. cupulaeformis quadrilobus aut subinteger.

Pet. longiora calyce, basi subcoalita.

Stam. quatuor petalis opposita, breviora. Filamenta lanceolata, glabra. Antherae oblongae.

Styl. quadrisulcus, e quatuor coalitis. Stigma quadrilobum.

Caps. tetracocca.

Andr. rep. t. 18. 436. 653. *Smith. exot. bot.* 2. t. 72. *Labill.* it. 2. t. 17. 19. *Bot. reg.* 3. 26. 515. *Mém. mus.* t. 21. n. 22.

1552. MELICOPE Forst. (1775.)

Cal. quadripartitus persistens.

Pet. unguibus in tubi formam conniventibus, laminis patentibus.

Glandulae quatuor didymae magnae, perigynae.

Stam. petalis breviora. Filamenta subulata. Antherae subcordatae.

Styli quatuor in unum coaliti. Stigma centro concavum.

Cocci quatuor monospermi, superne dehiscentes.

Forst. char. 28. [*Entoganum Banks. Gärtn.* t. 68.]

1553. PORLIERA Ruiz et Pav. (1794.)

Cal. tetrasepalus, aequalis.

Pet. subunguiculata, conniventia.

Filamenta basi squamulosa.

Stylus quadrisulcus, quasi e quatuor coalitis, apice quadrifidus.

Fructus carnosus, quadrilocularis. Sem. solitaria, pendula.

R. et P. prodr. t. 9. *Mém. mus.* 12. t. 16.

1554. CHITONIA Sess. Cand. (1824.)

Cal. quadripartitus, deciduus.

Pet. orbiculata.

Styl. stigmatate peltato-hemisphaerico.

Caps. quadrilocularis, quadrivalvis; valvis carinatis, subulatis. Sem. in singulis loculis bina, ovata, arillo caroso cincta.

1555. CYMINOSMA Gärtn. (1788.)

Cal. tetrasepalus, sepalis orbiculatis.

Pet. lanceolata, patentia.

Stam. disco caroso perigyno imposita: quatuor breviora, petalis opposita. Filamenta complanata, subulata. Antherae ovatae.

Stylus brevis. Stigma quadrisulcum.

Fructus carnosus, crustaceus, quadrilocularis; loculis monospermis.

Gärtn. t. 58. *Mém. mus.* 12. t. 17. [*Iambolifera pedunculata* L. *Vahl. symb.* 3. t. 61. *Iambolifera odorata* et *Gela lanceolata* Lour., quamvis fructu monospermo, tamen *R. Brownio* et *Lindleyo* testibus, huc pertinent.]

1556. XIMENIA Plum. (1703.)

Cal. quadrifidus, minimus, persistens.

Pet. basi conniventia, intus pilosissima, apice revoluta.
Filamenta subulata. *Antherae* lineares, elongatae, adnatae.
Stylus subulatus persistens.
Drupa ovata, monosperma.

Santaleis aff. *Lam.* ill. t. 297.

557. CASTELA Turp. (1806.)

Cal. parvus, quadridentatus.
Pet. cum dentibus calycis alternantia.
Discus perigynus circularis petala cum staminibus imposita gerit.
Stylus simplex.
Drupae quatuor monospermae, gynobasi carnosae insertae.
Ochnaceae. Ann. mus. 7. t. 5.

558. ELVASIA Cand. (1811.)

Cal. tetrasepalus.
Pet. ovalia, cum sepalis alternantia, aequalia, patentia.
Filamenta filiformia, hypogyna, persistentia. *Antherae* ovatae, erectae, rimis duabus dehiscentes.
Stylus simplex. Stigma capitatum.
Fructus quadrilocularis.
Ochnaceae. Ann. mus. 17. t. 20.

559. VITMANNIA Vahl. (1794.)

Cal. brevis, quadripartitus, extus biglandulosus.
Pet. unguiculata, oblonga.
Filamenta basi squamata. *Antherae* lineares, basi bifidae.
Stylus subulatus.
Drupa nuce semilunata, compressa.

Simarubaceae. Vahl. symb. 3. t. 60. [Samadera Gärtn. t. 156. Niota *Lam.* ill. t. 299.]

560. XYLOCARPUS König. (1784.)

Cal. coriaceus, quadrilobus.
Pet. patentissima, subcoriacea.
Urceolus staminifer carnosus, octodentatus. *Filamenta* linearia, emarginata. *Antherae* lineari-oblongae, lateri interiori filamentorum adnatae.
Styl. crassus brevis. Stigma latum, margine sulcato, disco perforato.
Drupa magna, exsucca, globosa, quadrivalvis; nucibus suboctoris monospermis.

Melieae? [Granatum litoreum *Rumph.* amboin. 3. t. 61. Carapa *Aubl.* guian. t. 387.]

561. KÖLREUTERA Laxm. (1772.)

Cal. pentasepalus; sepalis inaequalibus ascendentibus, inferne hiantibus.
Pet. ascendentia, unguiculata; unguibus basi squamatis.
Stam. *Filamenta* subulata, erecta, receptaculo columnari inserta. *Antherae* oblongae, obtusae. Glandulae tres perigynae.

Styl. trigonus ascendens. Stigma trifidum.

Caps. trilocularis, oligosperma.

Sapindeae. Laxmann in nov. comment. acad. petrop. 16. t. 18. Herit. sert. angl. t. 19.

1562. SCHMIDELIA L. (1767.)

Flores polygami.

Cal. quadripartitus.

Pet. inferne barbata.

Filamenta perigyna, subulata. Antherae subrotundae.

Stylus bifidus.

Fructus didymus baccatus, monospermus.

Sapindeae. [Usubis N. L. Burm. ind. t. 32. f. 1. Aporetica Forst. char. 66. Ornitrophe Juss. Lam. ill. t. 309. Labill. nov. Caled. t. 52. Allophylus L. Pal. Beauv. fl. ovar. 2. t. 107. Toxicodendron Gärtn. t. 44. Pometia Forst. char. 55. Gemella Lour.]

1563. AMYRIS L. (1749.)

Cal. quadridentatus persistens.

Pet. hypogyna, unguiculata.

Stam. petalis breviora.

Gynobasis disciformis incrassata. Germen uniloculare.

Stigma sessile.

Drupa nucleo chartaceo, monospermo.

Terebintheae. Humb. nov. gen. 7. t. 610.

1564. BALSAMODENDRON Kunth. (1824.)

Flores diclini.

Cal. quadridentatus persistens.

Pet. lineari-oblonga.

Annulus perigynus staminifer. Stamina cum verruculis annuli alternantia. Filamenta corolla breviora. Antherae oblongae, biloculares.

Styl. brevissimus. Stigma tetragonum.

Bacca mono-disperma.

Terebintheae. [Amyris gileadensis L. Vahl. symb. 1. t. 11. Balsamum P. Alpin. aegypt. p. 60. ic.]

1565. ELAPHRIUM Jacqu. (1763.)

Cal. quadripartitus, persistens.

Pet. basi lata, hypogyna, patentia.

Stam. sub disco orbiculari in fundo calycis inserta. Filamenta corolla breviora. Antherae ovaes, erectae, biloculares.

Styl. brevis. Stigmata duo capitata.

Drupa subbivalvis, nucibus duabus.

Terebintheae. Jacqu. amer. t. 71. Humb. nov. gen. 7. t. 611—613. [Icica Aubl. guian. t. 130—134.]

MELASTOMEAE.

1566. RHEXIA Gronov. (1743.)

Cal. basi ventricosus, ore angustatus, limbo patulo, quinquefidus.

Pet. obovata, ori calycis imposita.

Stam. cum petalis alternantia, iidem calycis ori imposita.

Antherae lineares, declinatae, rostratae, apice poro uno hiantes, dorso adnatae.

Styl. teres. Stigma capitatum.

Caps. globosa, calyce inclusa, quadrilocularis, quadrivalvis. Placentae baccatae, in mediis loculis. Sem. reniformia, scabra.

Lam. ill. t. 283. *Pursh.* amer. bor. t. 10.

567. *MARCEZIA* Cand. (1828.)

Cal. tubulosus, oblongus: limbo quadrifido.

Pet. ovata, acuta.

Stam. ori calycis inserta, aequalia. Antherae basi bituberculatae, oblongae, apice poro solitario hiantes.

Styl. filiformis. Stigma punctiforme.

Caps. quadrilocularis, quadrivalvis. Sem. cochleata.

Cand. Melastom. t. 5. [*Rhexia taxifolia* Bonpl. *Rhex.* t. 57.]

568. *OXYSPORA* Cand. (1828.)

Cal. tubulosus; limbo quadrifido, laciniis ovatis.

Pet. lanceolata, acuta.

Stam. subaequalia. Antherae elongatae, basi bituberculatae, supra basin calcaratae, apice poro solitario hiantes.

Styl. apice incurvus, obtusus.

Caps. quadrilocularis, quadrivalvis. Sem. minima scobiformia, utrinque aristata.

Cand. Melastom. t. 4.

569. *TRICENTRUM* Cand. (1828.)

Cal. urceolatus; limbo quadrifido; laciniis angustis apice setosis.

Pet. ovalia.

Stam. aequalia. Antherae lineares, rostratae, basi tricalcaratae, apice poro solitario hiantes.

Styl. filiformis.

Caps. quadrilocularis. Sem. cochleata.

[? *Rhexia leptophylla* Bonpl. *Rhex.* t. 24.]

570. *ARTHROSTEMMA* Pav. Don. (1824.)

Cal. campanulatus, setosus; limbo quadrifido, laciniis lanceolatis.

Pet. saepe apice aristata.

Stam. aequalia. Antherae oblongae, basi biauriculatae, apice poro solitario hiantes.

Caps. quadrilocularis. Sem. cochleata.

[*Rhexia* Aubl. guian. t. 129. Ruiz et Pav. fl. peruv. 3. t. 318. 319. 321. Bonpl. *Rhex.* t. 5. 14. 27. 29. Bot. reg. 1066. Melastoma Link. ic. nov. t. 24.]

1571. *HETERONOMA* Cand. (1828.)

Cal. tubulosus, dentibus quatuor triangularibus persistentibus.

Pet. ovalia submucronata.

Stam. Antherae alternae longiores; breviores basi bisetosae, longiores appendice lineari apice bifida.

Sem. cochleata, rugulosa, striata.

[*Rhexia Bonpl. Rhex. t. 45.*]

1572. *SPENNERA* Mart. Cand. (1826.)

Cal. globosus, limbi laciniis brevibus acutis.

Pet. lanceolata, acuta.

Antherae ovatae, obtusae, basi nudaе, apice poro solitario hiantes.

Caps. libera subbilocularis. *Sem.* cochleata, echinulata.

[*Melastoma aquatica Aubl. guian. t. 169. Rhexia Bonpl. Rhex. t. 38. 48. 49.*]

1573. *COMOLIA* Cand. (1828.)

Cal. campanulatus, limbi laciniis linearibus.

Pet. obovata.

Antherae aequales, falcatae, apice poro solitario hiantes, basi biauriculatae.

Caps. bilocularis. *Sem.* cochleata, dorso sulcata, tuberculato - scabra.

[*Rhexia berberifolia Bonpl. Rhex. t. 42.*]

1574. *APPENDICULARIA* Cand. (1828.)

Cal. urceolatus, limbo quadrilobo.

Pet. obovata.

Antherae utrinque productae ultra loculos, inferne bisetosae.

Caps. oblonga, trilocularis, trivalvis; columella centrali demum libera. *Sem.* cochleata.

[*Rhexia thymifolia Bonpl. Rhex. t. 50.*]

1575. *MEISNERIA* Cand. (1828.)

Cal. globosus, limbi laciniis lanceolatis acuminatis.

Pet. ovalia.

Antherae ovatae, rostro tubuloso, alternae basi auriculatae.

Caps. bilocularis. *Sem.* minima, ovato - triquetra.

[*Melastoma sessile * syst. 2. p. 303.*]

1576. *BUCQUETIA* Cand. (1828.)

Cal. globosus, limbi laciniis triangularibus.

Pet. obovata.

Stam. aequalia. *Antherae* oblongae, basi exauriculatae, apice poro solitario hiantes.

Germen valvis apice incrassatis truncatum et quatuor tuberculis notatum.

Caps. quadrilocularis. *Sem.* cuneato - angulata.

[*Rhexia glutinosa Bonpl. t. 47.*]

577. *MACAIREA* Cand. (1828.)*Cal.* ovatus, limbi laciniis lineari-lanceolatis.*Pet.* obovata.*Stam.* inaequalia. Filamenta omnia medio articulata, appendiculata; longiora pilis glanduliferis sub appendicula instructa. Antherae lineares, rectae, apice apertae, basi productae.*Caps.* quadrilocularis; libera.

[Cand. Melast. t. 6. [Rhexia Bonpl. Rhex. t. 41.]

578. *CALOPHYSA* Cand. (1828.)*Cal.* fructifer lagenarius, limbi laciniis latis plumoso-ciliatis.*Pet.* obovata.*Filamenta* plana, apice attenuata. Antherae oblongae, exauriculatae.*Caps.* baccata, quadrilocularis.

(Vesicae subrotundae utrinque ad basin petioli.)

579. *MEDINILLA* Gaudich. Cand. (1828.)*Cal.* tubus obovatus, ecostatus; limbo cylindrico, exacte truncato, persistente.*Pet.* subcoriacea, obtusa.*Stam.* aequalia. Antherae elongatae, acutae, apice poro solitario hiantes, basi biauriculatae.*Caps.* baccata, quadrilocularis. Sem. laevia.

(Folia verticillata.)

580. *SAGRAEA* Cand. (1828.)*Cal.* tubulosus; limbo quadridentato.*Pet.* ovalia.*Stam.* aequalia. Antherae basi subexauriculatae.*Caps.* baccata quadrilocularis. Sem. angulata.

[Melastoma rubra Rich. in Bonpl. Melast. t. 39.]

581. *OSSAEA* Cand. (1828.)*Cal.* subglobosus, vel ovato-campanulatus; limbo quadridentato.*Pet.* lanceolata, acuta.*Stam.* aequalia, brevia. Antherae basi subexauriculatae.*Bacca* quadrilocularis, dentibus calycis coronata. Sem. angulata.

[Melastoma Aubl. guian. t. 164. Bonpland. Melast. t. 36—38.]

582. *TETRAZYGIA* Rich. Cand. (1828.)*Cal.* globosus, urceolatus; limbo quadridentato.*Pet.* obovata.*Stam.* aequalia. Antherae lineares, basi obtusae, apice poro solitario hiantes.*Caps.* baccata quadrilocularis. Sem. nitida.

[Melastoma Vahl. ecl. t. 7. ic. amer. t. 26. 28. Bonpl. Melast. t. 10. 13. 34.]

1583. *ACIOTIS* Don. (1823.)

Cal. tubus globosus, carnosus; limbo coarctato persistente, quadridentato.

Pet. apice oblique aristata.

Stam. aequalia, medio articulata. Antherae erectae, basi nuda.

Caps. baccata, quadrilocularis. Sem. oochleata.

1584. *CHARIANTHUS* Don. (1823.)

Cal. ovatus suburceolatus; limbo persistente quadrilobo.

Pet. ovalia.

Stam. aequalia, Filamenta plana breviter. Antherae rimae duplici longitudinali dehiscentes, dorso carnosae.

Caps. baccata quadrilocularis. Sem. ovata.

[*Melastoma* Vahl, ic. amer. t. 16. Bonpl. Melast. t. 44.]

1585. *ANTICHOBUS* L. (1767.)

Cal. tetrasepalus, deciduus.

Pet. obtusa, calycem aequantia.

Stam. Filamenta setacea, erecta. Antherae subrotundae.

Styl. cylindricus. Stigma obtusum.

Caps. prismatica, quadrilocularis, quadrivalvis, polysperma. Sem. angulata.

Tiliaceae. Linn. fl. fasc. plant. hort. ups. t. 2. Lam. ill. t. 293.

1586. *ROXBURGHIA* Jones. Roxb. (1795.)

Cal. tetrasepalus.

Pet. calyce minora.

Antherae geminae, apice foliolo unitae.

Caps. unilocularis, bivalvis, polysperma.

Sarmentaceae? Roxb. corom. 1. t. 32. Bot. mag. t. 1500.

β. *Corolla pentapetala.*

1587. *XANTHOPHYLLUM* Roxb. (1819.)

Cal. pentasepalus.

Cor. irregularis. *Pet.* quatuor superiora subaequalia, quintum inferius carinatum.

Stam. incurva; sena unguibus petalorum, bina receptaculo inserta. Antherae erectae, ovaes.

Styl. stigmatibus bilobo.

Bacca monosperma, stipitata.

Capparidaceae. Roxb. corom. 3. t. 284. [Jackia Blum.]

1588. *BÄCKEA* L. (1757.)

Cal. turbinatus, limbo quinquefido persistente.

Pet. calyci imposita.

Stam. numero varia (5, 8, 10, 15.), calyci inserta, ut plurimum gemina inter petala. Filamenta subulata. Antherae subrotundae, biloculares.

Styl. brevis. *Stigma* capitatum.

Caps. trilocularis, polysperma.

Myrtaceae. Bot. mag. 2694. 2802. *Colla hort. ripul.* t. 6.
Rudg. in Linn. transact. 8. t. 12. 13. [*Leptospermum*
Labill. nov. Caled. t. 61. 62. *Jungia Gärtn.* t. 35. *Imbri-*
caria Sm.]

1589. HONCKENYA Willd. (1793.)

Cal. pentasepalus, coriaceus, intus coloratus.

Pet. oblonga.

Discus perigynus filamentosus.

Stam. Antherae oblongae.

Styl. simplex. *Stigma* denticulatum.

Caps. echinata, quadrilocularis, quadrivalvis, polysperma.

Sem. arillata.

Tiliaceae. Willd. in *Usteri del. opusc.* t. 4.

S A P I N D E A E

1590. MELICOCCA L. (*Melicoccus* P. Brown. 1756.)

Flores polygami.

Cal. quadri- quinquepartitus, persistens.

Pet. quatuor, vel quinque, disco hypogyno inserta, reflexa.

Filamenta subulata. Antherae oblongae, erectae.

Styl. brevissimus. *Stigma* peltatum, obliquum.

Bacca mono- (rarius di- tri- tetra-) sperma.

Juss. mém. du mus. 3. t. 4. 5. *Gärtn.* t. 42. [*Eustathes*
Lour.]

1591. HYPELATE P. Brown. (1756.)

Flores polygami.

Cal. pentasepalus, sepalis deciduis, duobus minoribus.

Pet. ovata, calyce minora.

Annulus perigynus.

Filamenta lanceolata, flexuosa. Antherae ovatae, biloculares.

Styl. brevis. *Stigma* deflexum, trigonum.

Drupa subrotunda, monosperma.

Swartz. fl. ind. occid. t. 14.

1592. CUPANIA Plum. (1703.)

Cal. pentasepalus.

Pet. cucullata, vel planiuscula.

Filamenta subulata, vel longiora, vel brevissima. Antherae incumbentes.

Styl. trifidus. *Stigmata* obtusa.

Caps. coriacea trilocularis, trivalvis, valvis medio septiferis, loculis mono-dispermis. *Sem.* arillata.

Plum. ic. t. 110. [*Molinaea Lam. ill.* t. 305. *Guioa Cav.*
ic. 4. t. 373.]

1593. DIPLOPETALON * (*Dimereza* Labill. 1825.)

Cal. pentasepalus, persistens, bracteatus.

Pet. orbiculata.

Discus glandulosus, turgidus, stamina ambiens, processus petaloideis bifidis, apice callosis, qui petalis incumbunt.

Stam. erecta, basi subdilatata, ciliata. Antherae ovatae erectae.

Caps. coriacea, trivalvis, trisperma.

[*Dimereza* Labill. nov. Caled. t. 51.]

1594. *THOUINIA* Poit. (1804.)

Cal. quadri-quinquepartitus.

Pet. quatuor, vel quinque, intus barbata.

Discus glanduloso-squamosus, stamina ambiens.

Filamenta erecta. Antherae ovatae, erectae, biloculares.

Styl. simplex. Stigma trifidum.

Achenia tria, compressa, superne alata.

Annal. mus. 3. t. 6. vol. 5. t. 26. 27. *Humb.* plant. aequin. 1. t. 56. *Mart.* in nov. act. nat. cur. 12. t. 4.

1595. *EUPHORIA* Commers. Juss. (1789.)

Flores polygami vel monoeci.

Cal. quinquedentatus.

Pet. basi squamam ciliatam foveantia.

Discus perigynus glandulosus.

Stam. erecta, basi connata. Antherae oblongae, erectae, biloculares.

Germen didymum. Stylus brevissimus.

Bacca cortice tuberculato, monosperma. Sem. arillatum.

Bot. reg. 1059. [Li-tchi *Sonnerat.* it. ind. 2. t. 129. *Scytalia* *Gärtn.* t. 42. *Dimocarpus* *Lour.* *Nephelium* *L.* *Marsd.* Sumatr. t. 2.

1596. *MAGONIA* Hilar. (1825.)

Flores polygami.

Cal. quinquepartitus; laciniis inaequalibus, reflexis.

Pet. spathulato-linearia, elongata.

Annulus submembranaceus stamina ambiens, simplex vel duplex.

Stam. declinata. Antherae ellipticae, bifidae.

Styl. curvatus. Stigma trilobum.

Caps. lignosa, trivalvis, polysperma. Sem. plana, undique alata.

Mém. mus. 12. t. 12. 13. A. [*Phaeocarpus* *Mart.* nov. gen. t. 37. 38.]

1597. *EPHIELIS* Schreb. (1789.)

Cal. quinquepartitus.

Pet. unguiculata, patentia. Squamae binae singulis petalis ad basin adnatae.

Filamenta setacea, glandulis inserta. Antherae tetragono-subrotundae.

Discus glandulosus perigynus. *Stylus* nullus. *Stigma* obtusum.

Caps. oblonga, compressa, utrinque sulcata, unilocularis, bivalvis. Sem. duo reniformia.

[*Mataiba Aubl. guian. t. 128.*]

598. *TALISIA* Aubl. (1775.)

Flores polygami.

Cal. quinquepartitus.

Pet. cum laciniis calycis alternantia, unguiculata, disco perigyno imposita.

Squamae pilosae fornicatae *stamina* obtegentes. *Filamenta* brevissima. *Antherae* oblongae, in filamenta decurrentes.

Germen quadriloculare.

Stylus brevis.

Aubl. guian. t. 136.

599. *BLIGHIA* König. (1806.)

Cal. quinquepartitus.

Pet. basi squamis fornicatis instructa, acuminata.

Discus carnosus, crenatus, staminifer.

Filamenta medio incrassata, subulata. *Antherae* didymae, biloculares.

Stylus subnullus. *Stigma* simplex.

Fructus carnosus, sexlobus, trilocularis, trivalvis: loculis monospermis. *Arillus* seminis maximus cordatus, receptaculiformis.

Ann. of bot. 2. t. 16. 17. [Akeesia Tussac. antill. t. 3.]

600. *ACER* Plin.

Flores polygami.

Cal. quinquepartitus.

Pet. patentia, vel nulla.

Filam. subulata, receptaculo inserta.

Styl. filiformis. *Stigmata* duo.

Caryopses duae alatae.

Acera Juss. Schk. t. 351.

601. *TROPAEOLUM* L. (1737.)

Cal. quinquepartitus, coloratus, postice calcaratus.

Pet. calyci inserta, inaequalia.

Stam. *Filamenta* declinata. *Antherae* erectae, oblongae, quadrilobae, biloculares.

Stylus simplex. *Stigma* trifidum.

Fructus tres bacciformes, rugoso-sulcati. *Cotyledones* crassae, demum arcte conferruminatae.

Tropaeoleae Juss. Schk. t. 105. Gärtn. t. 79. Bot. reg. 718.

1602. MAGALLANA Cav. (1797.)

Cal. tripartitus, calcaratus; laciniis duabus divaricatis indivisis, tertia inferiore trifida.

Pet. calyci inserta, inaequalia; duo inferiora sessilia, tria superiora unguibus filiformibus praedita.

Annulus perigynus staminifer.

Filamenta filiformia. *Antherae* ovatae didymae.

Styl. filiformis. *Stigma* bifidum, lacinia altera crassa brevi, altera elongata, subulata divaricata.

Fructus trialatus, abortu monospermus.

Tropaeoleae Juss. *Cav. ic.* 4. t. 374.

γ. Corolla octopetala.

1603. JEFFERSONIA Barton. (1793.)

Cal. coloratus tetra- (tri- penta-) sepalus.

Pet. patentia, oblongo-linearia, obtusa.

Stam. erecta, petalis opposita. *Filamenta* brevissima.

Stigma subsessile, peltatum.

Caps. obovata, substipitata, unilocularis, polysperma, sub apice foramine lunato elastice dehiscens. *Placenta* lateralis. *Sem.* arillo basilari demum lacero instructa.

Podophylleae Cand. *Act. soc. amer.* 3. p. 334. *ic. Bot. mag.* 1513.

b. Fructus inferi.

O N A G R A E Juss.

1604. OENOTHERA L. (1737.) (*Ὠνοθήρα* Theophr.)

Cal. tubulosus, limbo quadripartito deciduo; laciniis binis ternisve saepe cohaerentibus.

Pet. calyci inserta.

Stam. erecta vel declinata. *Filamenta* subulata. *Antherae* incumbentes. *Pollen* viscosum subtriangulare.

Styl. filiformis. *Stigma* quadrifidum (rarius globosum).

Caps. prismatico-cylindrica, vel obovata, alata, quadrilocularis, quadrivalvis. *Placenta* centralis. *Sem.* nuda.

Schk. t. 105. *Hook. exot. fl.* t. 140.

1605. CLARKEA Pursh. (1814.)

Cal. tubulosus, limbo quadripartito; laciniis ternis saepe cohaerentibus.

Pet. calyci inserta; unguibus utrinque unidentatis, limbo trilobo.

Stam. alterna sterilia, unguibus petalorum inserta, clavata; alterna fertilia, laciniis calycis opposita. *Antherae* lineares, biloculares, post anthesin spiratim revolutae.

Styl. filiformis. *Stigma* quadrilobum, subpetaloideum.

Caps. cylindrica, sulcata, quadrilocularis.

Pursh, amer. bor. t. 11. *Bot. reg.* t. 1100.

606. *EPILOBIUM* C. Gesn. (1561.)*Cal.* tubulosus, limbo quadripartito.*Pet.* calyci inserta.*Stam.* ascendentia vel declinata. Filamenta subulata.

Antherae ovales. Pollen viscosum subtriangulare.

Styl. simplex. Stigma quadrifidum.*Caps.* prismatica, quadrilocularis, quadrivalvis, polysperma.

Sem. comosa.

Schk. t. 106. *Gärtn.* t. 31. *Reichenb.* ic. 2. t. 170. 180. 189. 190. 197—200. 341. 342.607. *JUSSIEVA* L. (1747.)*Cal.* tubuloso-prismaticus, limbo quadripartito persistente.*Pet.* patentia, calyci inserta; ut plurimum quatuor, quandoque sex aut quinque.*Stam.* plerumque octona, nonnunquam et dena.*Filamenta* calyci inserta, brevia. Antherae ovales, erectae.*Styl.* cylindricus. Stigma peltatum.*Caps.* prismatica, quadrilocularis, polysperma. Sem. nuda, glaberrima, minuta.*Reichenb.* hort. t. 67. 75. *Bot. mag.* 2530. *Humb.* plant. aequin. 1. t. 3. nov. gen. 6. t. 530—532. [*Cubospermum* Lour.]608. *HAUYA* Sess. Cand. (1828.)*Cal.* cylindricus, elongatus; limbo quadripartito intus colorato.*Pet.* calyci inserta, ovalia.*Stam.* exserta. Antherae versatiles oblongae.*Styl.* filiformis. Stigma capitatum.*Caps.* quadrilocularis, quadrivalvis, polysperma. Sem. apice in alam expansa.609. *GAURA* L. (1751.)*Cal.* tubulosus, (tri-) quadripartitus, caducus.*Pet.* plerumque quatuor (interdum tria).*Stam.* calyci inserta. Filamenta subulata. Antherae incumbentes ovales, rimis binis longitudinalibus dehiscentes.*Stylus* simplex. Stigma quadrifidum.*Nux* tetragona, mono-disperma.*Schk.* t. 115. *Gärtn.* t. 127. *Bot. mag.* 388. 389. *Humb.* nov. gen. 6. t. 529.610. *FUCHSIA* Plum. (1703.)*Cal.* tubulosus coloratus, limbo quadrifido.*Pet.* calycis fauci imposita, conniventia, convoluta (in F. excorticata L. f., patula).*Stam.* vel inclusa, duplici serie tubo calycis adnata, vel exserta, alterna longiora. Antherae ovales erectae.

Pollen viscosum, triangulare.

Stylus filiformis. Stigma capitatum vel quadrifidum.

Bacca quadrilocularis, quadrivalvis, polysperma.

Link. ic. sel. t. 46. Bot. reg. 847. 857. 945. 1048. 1052. 1062.

Humb. nov. gen. 6. t. 534. 535.

1611. MEMECYLON L. (1747.)

Cal. campanulato - turbinatus, subquadri - quinquedentatus aut repandus.

Pet. quatuor aut quinque, ovalia, cum dentibus calycis alternantia.

Filamenta libera filiformia. *Antherae* incurvae, medio affixae, biloculares, hinc rostratae.

Styl. filiformis. *Stigma* simplex.

Bacca limbo calycis coronata, adulta unilocularis monosperma. Sem. pendulum exalbuminosum.

Memecyleae Cand. *Lam. ill. t. 284. Roxb. corom. 1. t. 82. [Cornus sylvestris Burm. zeyl. t. 30. Scutula Lour.]*

1612. MARLEA Roxb. (1819.)

Cal. campanulatus, sex - octodentatus.

Pet. sex, octo, linearia, reflexa.

Filamenta brevissima plana pilosa, corpusculo glanduloso perigyno imposita. *Antherae* lineares, elongatae.

Stylus simplex. *Stigma* quadridentatum.

Drupa nuce biloculari.

Alangiaeae. *Roxb. corom. 3. t. 283. [Stylidium Lour.]*

1613. RHIZOPHORA L. (1747.)

Cal. tubulosus, limbo quadri - (quinque - octo - decem -) lobo.

Pet. tot, quot calycis lobi, iidem inter lobos inserta.

Stam. calyci inserta, octona, (dena, dena sena, vicena).

Filamenta subulata, erecta. *Antherae* basi insertae.

Stylus apice bifidus.

Fructus calyce coronatus, non dehiscent, monospermus.

Embryo germinans, e semine longe exsiliens.

Rhizophoreae R. Br. *Jacqu. amer. t. 89. Gärtn. t. 45. [Mangium Rumph. amboin. 3. t. 68—72. Bruguiera Lam. ill. t. 397.]*

1614. COMBRETUM Löff. (1758.)

Cal. infundibuliformis quadrilobus.

Pet. quatuor, calyci inserta.

Stam. calyci duplici ordine inserta, quatuor petalis opposita. *Antherae* biloculares, longitudinaliter dehiscentes.

Styl. acutus.

Fructus nucamentaceus, monospermus, tetrapterus. Sem. pendulum. Cotyledones plica media reflexae, cruciatio-divaricatae.

Combreteae. *Humb. plant. aequin. 2. t. 152. nov. gen. 6. t. 538. Palis. Beauv. fl. ovar. t. 118. Aubl. guian. t. 137. Reichenb. hort. bot. t. 62.*

II. *Flores imperfecti.**Fructus superi.*

615. **JONESIA** Roxb. (1795.)
Cal. bracteis binis subrotundis suffultus, infundibuliformis, coloratus, tubo clauso carnosq, limbo quadrilobo.
Annullus faucis calycis staminifer.
Stam. longe exserta, basi interdum concreta. *Germen* stipitatum. *Styl.* filiformis.
Legumen acinaciforme, suturis callosis, tetra-octospermum.
Leguminosae. [Saraea N. L. Burm. ind. t. 25. f. 2.]
616. **STADMANNIA** Lam. (1793.)
Cal. exiguus, dentibus quinque brevissimis acutis.
Cor. nulla.
Filamenta subulata, patentia, exserta. *Antherae* subrotundae.
Styl. subnullus. *Stigma* triangulare.
Bacca globosa mono-disperma.
Sapindeae. Lam. ill. t. 312. [Melicocca Juss. mém. mus. 3. t. 7. 8. Schleichera Willd.]
617. **PIERARDIA** Roxb. (1820.)
Flores monoeci.
Cal. quadripartitus subtomentosus.
Filamenta filiformia. *Antherae* biloculares.
Styl. simplex. *Stigmata* tria.
Bacca globosa, trilocularis, trisperma. *Sem.* arillo sapido tunicata. *Embryo* albumine inclusus.
— [Raubeh Marsden Sumatr. t. 6.]
618. **VALENTINIA** Sw. (1788.)
Cal. quinquepartitus, coloratus, patens, laciniis obtusis.
Cor. nulla.
Filamenta subulata. *Antherae* subrotundae.
Styl. teres crassus. *Stigma* capitatum.
Caps. baccata, pulposa, in tres seu quatuor partes revolvendas dehiscens, unilocularis, subtetrasperma.
Sapindeae. Swartz. fl. ind. occident. t. 14.
619. **LLAGUNOA** Ruiz et Pav. (1794.)
Flores polygami.
Cal. quinquefidus, incisura infima productiore.
Discus hypogynus in fundo calycis, subdimidiatus, staminifer.
Stam. pendula inter incisuram inferiorem calycis.
Styl. subulatus. *Stigma* obtusum.
Caps. trigona, trilocularis, trivalvis; loculis submonospermis. *Sem.* ossea, nitida, exalbuminosa. *Embryo* spiritaliter convolutus.
Sapindeae. Ruiz et Pav. prodr. t. 28. Humb. nov. gen. 5. t. 442. [Amirola Pers.]

1620. DODONAEA L. (1737.)

Flores polygami.

Cal. tetra- (tri- penta-) sepalus, subaequalis.*Cor.* nulla. Discus hypogynus nullus.*Filam.* brevissima. Antherae crassae, tetragonae, basi affixae, biloculares.*Styl.* trigonus. Stigma trilobum.*Caps.* membranaceae, triquetrae, trialaetae, triloculares.

Sem. subrotunda, dura. Embryo spiraliter convolutus.

Sapindeae. *P. Brown.* *jam.* t. 18. f. 1. *Humb.* *nov. gen.* 5. t. 442. *Bot. mag.* 2860.

† 1621. ANIBA Aubl. (1775.)

Cal. sexpartitus; laciniis obtusis, concavis.*Filamenta* brevia. Antherae ovaes.Germen glandula cinctum. *Styl.* brevis. Stigma obtusum.*Fructus*?— — *Aubl. guian.* t. 126. [*Cedrota Schreb. Willd.*]

† 1622. BACCAUREA Lour. (1790.)

Flores polygami.

Cal. quinquepartitus, persistens, carnosus.*Cor.* nulla.*Stam.* receptaculo inserta. *Filamenta* brevissima. Antherae subrotundae.

Stigma sessile peltato-concavum.

Bacca subrotunda, tri- (quinque-) locularis, mono-trisperma.

† 1623. GRUBBIA Berg. (1767.)

Involucrum diphyllum, bi-triflorum.*Cor.* tetrapetala, extus lanata.*Filamenta* subulata corolla breviora. Antherae subrotundae.*Stylus* brevis. Stigma simplex.*Nucula* monosperma.*Thymelaeae.* *Berg. act. acad. holm.* 1767. t. 2. fl. esp. t. 2.

1624. DAPHNE Diosc.

Cal. corollinus, infundibuliformis, marcescens, quadrifidus.*Stam.* tubo corollae duplici serie inserta. Antherae subrotundae.*Styl.* brevissimus. Stigma capitatum.*Drupa* monosperma.*Thymelaeae.* *Schk.* t. 107. a. *Sturm. fl. germ.* [Capura L.]

1625. GNIDIA L. (1753.)

Cal. corollinus, infundibuliformis, limbo quadrifido. Squamae quaternae, cum laciniis limbi alternantes, vel octonae, laciniis oppositae.*Stam.* Antherae subsessiles, duplici serie tubo calycis insertae.*Stylus* lateralis. Stigma capitatum vel penicillatum.

Nux monosperma.

Thymelaeae. Bot. reg. 2. 19. 624. 757. [*Nectandra*
Berg.]

526. *PASSERINA* L. (1737.)

Cal. corollinus, infundibuliformis, marcescens; limbo quadrifido.

Filam. setacea, fauci imposita. *Antherae* ovatae, erectae.

Styl. lateralis. *Stigma* capitellatum.

Nux monosperma.

Thymelaeae. Wendl. obs. t. 2. f. 15 — 20. *Gärtn.* t. 215.

[*Stellera* Gmel. *L. Jacqu.* ic. rar. 1. t. 68. *Gärtn.* t. 39.

Lachnaea L. *Lam.* ill. t. 292. *Gärtn.* t. 215.]

627. *LAGETTA* Juss. (1789.)

Flores dioeci.

Cal. corollinus, tubulosus, limbo quadrifido. *Squamulae* octo, lineares setaceae in fundo calycis.

Stam. duplici serie tubo adnata. *Filamenta* brevissima.

Antherae ovatae, biloculares.

Styl. simplex. *Stigmata* duo capitata.

Drupa sicca calyce persistente cincta, monopyrena.

Thymelaeae. Mart. nov. gen. t. 39.

628. *DIRCA* L. (1751.)

Cal. nullus.

Cor. campanulata, limbo obsoleto, margine inaequali.

Stam. *Filamenta* capillaria, tubo medio imposita, exserta, alterna longiora. *Antherae* erectae, subrotundae.

Styl. apice incurvus.

Bacca monosperma.

Thymelaeae. Schk. t. 107. b.

629. *SHEPHERDIA* Nuttall. (1818.)

Flores dioeci,

Cal. quadrifidus, tubulosus, extus lepidotus.

Stam. inclusa, cum totidem glandulis alternantia.

Stylus incurvus.

Bacca monosperma.

Elaeagnaeae. [*Hippophaë canadensis* L., *argentea* Pursh.]

630. *ASPIDISTRA* Ker. (1822.)

Cal. corollinus, subcampanulatus; limbo sex - octofido, patente.

Stam. *Antherae* sessiles, tubo calycis adnatae, biloculares

Styl. clavatus. *Stigma* maximum, peltato - pileatum, convexum.

Germen triloculare, loculis monospermis.

Asphodeleae. Bot. reg. 628. 977. [*Macrogynae* Link. ic. t. 31.]

631. *POPULUS* Plin.

Flores dioeci.

Bractea multifido - laciniata, fimbriata.

Cor. campanulata, limbo integerrimo.

Stam. Filamenta basi corollae inserta. *Antherae* erectae biloculares.

Stigmata sessilia.

Caps. bi-trivalvis. Sem. comosa.

Amentaceae. Schk. t. 330. a. b.

1632. *BUGINVILLEA* Commers. Juss. (1789.)

Bracteae coloratae, ut plurimum ternae, e quarum singularum medio nervo

Cal. corollinus, tubulosus, marcescens; limbo plicato, denticulato.

Stam. inclusa. Filamenta capillaria, libera, basi calycis inserta. *Antherae* oblongae.

Stylus lateralis. *Stigma* clavatum.

Achenium calycis basi inclusum.

Nyctagineae. Lam. ill. t. 294. *Humb. plant. aequin.* t. 49. [Neaea Ruiz et Pav.]

D I G Y N I A.

* 1633. *SPALLANZANIA* Pollin. (1816.)

Cal. exterior decem- aut duodecimfidus; interior quinquefidus persistens.

Pet. quinque.

Stam. subulata.

Caryopsis calyce interno tecta.

Rosaceae? Pollin. plant. veron. fig. 1. *Röm. fl. europ.* t. 1. [Agrimouoides *Column.* ecphr. p. 144. ic. *Agrimonia* Agrimouoides L. *Aremonia* Neck. *Amqnia* Nestl.]

1634. *CODIA* Forst. (1775.)

Flores capitati, involucrati. *Receptaculum* commune vilosum. *Involucri* foliola quatuor, quinque oblonga, sericea, reflexa.

Involucrum parziale cupulare denticulatum.

Cal. quadri-quinquepartitus, intus pilis densis longis stipatus, laciniis oblongis trinerviis.

Pet. quatuor aut quinque linearia, cum laciniis calycis alternantia.

Stam. Filamenta subulata, basi unguibus petalorum adhaerentia. *Antherae* ovatae, biloculares, incumbentes.

Styli duo, stigmatibus acuminatis.

Nucula inter calycem et involucrum, monosperma.

Cunoniaceae R. Br. *Forst. char.* 30. *Labill. nov. Caled.* t. 46.

1635. *WEINMANNIA* L. (*Windmannia* P. Br. 1756.)

Cal. quadripartitus, persistens.

Pet. quatuor aequalia, fundo calycis inserta, cum eius laciniis alternantia.

Stam. inter petala et discum perigynum posita. Filamenta flexuosa, petalis longiora. Antherae subrotundo-ovatae, dorso affixae, biloculares.

Styli persistentes acuti. Discus urceolatus perigynus.

Capsulae binae rostratae, cylindricae, biloculares, oligospermae. Sem. minuta, pilosa.

Cunoniaceae R. Br. Lam. ill. t. 313. Cav. ic. 6. t. 566. 567. Humb. nov. gen. 6. t. 520—525.

636. MÖHRINGIA L. (1745.)

Cal. tetrasepalus, persistens.

Pet. quatuor, cum sepalis alternantia.

Stam. hypogyna, petala aequantia. Filamenta capillaria. Antherae ovatae.

Styli filiformes. Stigmata subclavata.

Caps. unilocularis, quadrivalvis, polysperma. Placenta centralis.

Caryophylleae. Schk. t. 108. Gärtn. t. 129.

637. GALENIA L. (1737.)

Cal. minimus, quadrifidus, concavus, laciniis oblongis.

Cor. nulla.

Filamenta capillaria. Antherae didymae.

Styli reflexi. Stigmata simplicia.

Caps. subglobosa, bilocularis, disperma.

Chenopodeae. Lam. ill. t. 314.

638. ASTILBE Hamilt. Don. (1825.)

Cal. profunde quadri-quinquepartitus, coloratus, imbricatus, laciniis ovalibus, concavis.

Cor. nulla.

Stam. laciniis calycis opposita. Filamenta subulata. Antherae globosae, didymae, rimis longitudinalibus binis dehiscentes.

Styli simplices, stigmatibus truncatis.

Caps. birostris, bilocularis, polysperma.

TRIGYNIA.

639. PAULLINIA Plum. (1703.)

Cal. pentasepalus, imbricatus, persistens.

Petala quatuor, unguiculata, basi squamata. Glandulae duae aut quatuor inter petala et stamina.

Stam. inaequalia. Antherae oblongae, dorso affixae, biloculares.

Styli crassiusculi, abbreviati, conniventes.

Caps. pyriformis, trigona, trilocularis; valvis navicularibus; axi centrali trilato. Sem. basi arillata; arillo bilobo, fungoso.

Sapindeae. Jacq. obs. 3. t. 61. 62. Plum. ic. t. 111. Gärtn. t. 79. [Semarillaria Ruiz et Pav.]

1640. *SERIANIA* Plum. (1703.)*Cal.* pentasepalus, imbricatus, persistens.*Petala* quatuor, unguiculata, basi squamata. Glandulae quatuor inaequales inter petala et stamina.*Stam.* inaequalia. Antherae oblongae, dorso affixae, biloculares.*Styli* crassiusculi, abbreviati, conniventes.*Samarae* tres axi centrali affixae.*Sapindeae.* Juss. in ann. mus. 4. t. 66. Humb. nov. gen. 4. t. 441.1641. *URVILLEA* Kunth. (1821.)*Cal.* pentasepalus, coloratus, sepalis duobus exterioribus multo brevioribus.*Pet.* quatuor unguiculata, basi squamata.*Stam.* imae basi germinis circumposita.*Germen* stipitatum, stipite fundo calycis longitudinaliter adnato. Glandulae duae maculaeformes ad utrumque stipitis latus, duae ad basin petalorum.*Fructus* membranaceus, trialatus, medio subinflatus ibique trilocularis, compositus e tribus samaris.*Sapindeae.* Humb. nov. gen. 5. t. 440. Bot. reg. 1077.1642. *PONAEA* Schreb. (1789.)*Cal.* quinquepartitus, laciniis subrotundis.*Pet.* quatuor lanceolata, patentia.*Discus* glandulosus hypogynus, cui et petala et stamina inserta.*Filamenta* capillaria, patentia. Antherae ovales, dorso adnatae.*Germen* stipitatum, triquetrum. *Styli* breves acuti.*Caps.* trilocularis, trialata; alis bivalvibus; seminibus solitariis.*Sapindeae.* [Toulicia Aubl. guian. t. 140.]1643. *SAPINDUS* Plum. Tournef. (1719.)*Cal.* tetrasepalus, coloratus, sepalis subaequalibus, duobus exterioribus.*Pet.* quatuor unguiculata, basi esquamata, intus glandulosa aut barbata.*Filamenta* villosa.*Fructus* tres globosi, carnosi, connati, monospermi.*Sapindeae.* Roxb. corom. 1. t. 62. Aubl. guian. t. 138-139.1644. *CARDIOSPERMUM* L. (1737.)*Cal.* tetrasepalus, sepalis duobus minoribus.*Pet.* quatuor, basi appendiculata, appendiculis inaequalibus.

Glandulae duae hypogynae inter petala et stamina.

Stam. hypogyna.

Germen stipitatum. Caps. tres inflatae, dorso alatae, evalves, ad axin connatae. Sem. globosa, hilo lato cordato.

Sapindeae. Lam. ill. t. 317. Gärtn. t. 79. Humb. nov. gen. 5. t. 139.

645. DUVAUA Kunth. (1825.)

Flores diclini.

Cal. quadrifidus, regularis, persistens.

Pet. quatuor, patentissima.

Discus urceolatus, sinuatus.

Stam. sub disco inserta. Antherae ovatae, biloculares.

Styli brevissimi. Stigmata capitata.

Drupa globosa, monosperma.

Terebinthaceae. [Amyris polygama Cav. ic. 3. t. 239. Schinus dependens Ort., Sch. dentatus Andr. repos. 620.]

646. PEDICELLIA Lour. (1790.)

Flores polygami.

Cal. quinquepartitus, laciniis ovatis acutis.

Cor. 0.

Discus carnosus crenatus.

Filamenta filiformia, reflexa. Antherae biloculares, erectae.

Germen pedicellatum. Styl. nullus. Stigmata tria acuta, reflexa.

Caps. pedicellata, subrotunda, trivalvis, monosperma. Sem. arillatum.

647. POLYGONUM Colum.

Cal. pentasepalus, persistens.

Cor. nulla.

Stam. octo (saepè sex, interdum quinque, raro quatuor, rarissime tria). Filamenta filiformia, quandoque basi glandulata. Antherae erectae biloculares.

Styli tres, stigmatibus capitatis, (aut stylus bifidus).

Achenium triquetrum, (vel subrotundum) albuminosum.

Embryo lateralis, vel medius, S-formis.

Polygonaceae. Schk. t. 108. Gärtn. t. 119. [Fagopyrum Tourn. Gärtn. l. c.]

648. TRAGOPYRUM Marsch. Bieb. (1819.)

Cal. pentasepalus, sepalis duobus exterioribus reflexis, tribus interioribus fructum obtegentibus.

Cor. nulla.

Stam. octo.

Stylt tres distincti aut basi connati. Stigmata capitata.

Achenium triquetrum, albuminosum. Embryo lateralis.

Polygonaceae. [Polygonum frutescens Gärtn. t. 119. Gmel. sib. 3. t. 12.]

649. COCCOLOBA Jacqu. L. (Coccolobis P. Brown. 1756.)

Cal. quinquepartitus, persistens, demum incrassatus.

Cor. nulla.

Filamenta subulata. *Antherae* didymae.

Styli patentes.

Nux monosperma, calyce baccato inclusa.

Polygonaceae. *P. Brown*, *jam.* t. 14. f. 3. *Gärtn.* t. 45.

Plum. ic. t. 146. f. 1.

T E T R A G Y N I A.

1650. VEREA Andr. (1797.)

Cal. quadripartitus, persistens.

Cor. basi ventricosa, hypocrateriformis, limbo patente quadrifido.

Stam. duplici serie tubo corollae adnata. *Filamenta* brevissima. *Antherae* ovatae, erectae. *Squamae* quatuor lineares hypogynae.

Styli obtusi conniventes.

Caps. quatuor trigonae, polyspermae, intus dehiscentes.

Sedae. *Andr.* *repos.* t. 21. 560. [*Kalanchoë Adans. Cand. succul.* t. 64. 65. 100. *Bryophyllum Salisb. Bot. mag.* t. 1409.]

1651. DIAMORPHA Nuttall. (1818.)

Cal. quadrilobus.

Pet. quatuor.

Caps. quatuor, basi concretionae, cuspidatae, extus dehiscentes, tetraspermae.

Sedae. [*Sedum pusillum Mx. Tillaea cymosa Nutt.*]

652. GONIOCARPUS König. (1805.)

Cal. superus, limbo quadripartito.

Pet. quatuor, cum lacinüs calycis alternantia, summo calycis tubo inserta.

Stam. octo, vel quatuor.

Styli breves, stigmatibus penicillatis: vel stigmata quatuor sessilia.

Drupa octangula octosperma.

Haloragae R. Br. Ann. of bot. 1. t. 12. [*Gonocarpus*

Thunb. fl. jap. t. 15. *Labill. nov. holl.* t. 53.]

1653. HALORAGIS Forst. (1775.)

Cal. superus, limbo quadrifido.

Pet. summo tubo inserta.

Filamenta capillaria, erecta. *Antherae* lineares, incumbentes.

Stigmata sessilia.

Drupa quadrangula tetrasperma.

Haloragae R. Br. Forst. char. 31. *Jacqu. ic. rar.* 1.

t. 69. *Labill. nov. holl.* t. 128. 129. [*Cercodia Murr. comment. gött.* 3. t. 1.]

654. GALVEZIA Ruiz et Pav. (1794.)

Cal. quadripartitus.*Pet.* hypogyna, lanceolata.*Stam.* basi stipitis germen ferentis inserta. Filamenta subulata. Antherae ovatae, dorso affixae.*Stipes* carnosus germina quatuor ferens. *Styli* basi distincti, apice in stigma quadrilobum confluentes.*Drupae* quatuor monospermae.*Rutaceae.* *Adr. Juss.* in mém. mus. 12. t. 25. f. 37.
[Pitavia *Molin.*]655. VEPRIIS Commers. *Adr. Juss.* (1825.)

Flores diglini.

Cal. quadripartitus.*Pet.* quatuor, lanceolata, patentissima.*Stam.* stipiti germen ferenti inserta. Filamenta filiformia.

Antherae ovatae, dorso affixae.

Stipes glandulosus, octolobus germina quatuor ferens, squamulis octo minutissimis instructus. *Styli* quatuor in flore masculo, stigmatibus capitatis: in flore femineo stigma sessile, quadrilobum, convexo-peltatum.*Fructus* carnosus, quadrilocularis, tetraspermus.*Rutaceae.* *Adr. Juss.* mém. mus. 12. t. 26. f. 46.

656. FRANCOA Cav. (1800.)

Cal. quadripartitus persistens.*Pet.* quatuor, unguiculata, acuta.*Stam.* receptaculo inserta. Filamenta basi glandulis totidem obsessa. Antherae biloculares.*Stigmata* sessilia plana.*Caps.* quatuor connatae, polyspermae, bivalves.— — *Cav.* anal. de cienc. nat. 4. t. 36. [Laupanke *Feuill.* obs. 2. t. 31. Panke *Möln.*]

657. POLYSCIAS Forst. (1775.)

Cal. superus, cupularis, obsolete denticulatus.*Pet.* sex, septem, octo, lineari-lanceolata, patentissima.*Filamenta* capillaria, cum petalis alternantia. Antherae biloculares.*Styli* nulli. *Stigmata* tria, quatuor, quinque, patentiuscula.*Bacca* globosa, quadrilocularis. Sem. solitaria, angulata.*Araliaceae.* *Forst.* char. 32.

658. PYTHAGOREA Lour. (1790.)

Cal. inferus campanulatus, suboctosepalus; sepalis linearibus, coloratis.*Petala* totidem lanceolata.*Filamenta* subulata. Antherae didymae.*Styli* subulati, reflexi.*Caps.* ovata, quadrilocularis, polysperma.

— —

1659. MARGARITARIA L. fil. (1781.)

Flores dioeci.

Cal. quadridentatus, inferus.*Pet.* quatuor, subrotunda, calyci inserta.*Filamenta* setacea, receptaculo inserta. *Antherae* subrotundae.*Styli* in ♀ quatuor, vel quinque subulati; in ♂ stylus simplex.*Bacca* globosa, stylis persistentibus coronata, tetra-pentacocca.

— — [Eponymus margaritifera Plukn. t. 176. f. 4.]

1660. ELATINE L. (1737.)

Cal. inferus quadri- (tri-) partitus.*Pet.* quatuor ovalla (interdum tria).*Stam.* octo (sex, tria). *Filamenta* hypogyna subulata.*Antherae* ovales, erectae, biloculares.*Styli* quatuor, stigmatibus capitatis: aut stigmata sessilia tria.*Caps.* quadri- (tri-) locularis, quadri- (tri-) valvis. Sem. linearia, curvata, rugosa.*Caryophylleae.* Schk. t. 109. a. b. *Reichenb.* ic. 5. t. 413.*Cand.* ic. rar. t. 43. [Biolia Bellard.]

1661. PARIS Fuchs. (1542.) Rupp. (1718.)

Cal. inferus tetrasepalus; sepalis lanceolatis acuminatis patentibus.*Pet.* linearia subulata, patentia, persistentia.*Stam.* hypogyna. *Filamenta* subulata. *Antherae* oblongae, mediis filamentis adnatae.*Styli* patentes.*Bacca* quadrilocularis, polysperma.*Sarmentaceae.* Schk. t. 109. a.

1662. ADOXA L. (1737.)

Cal. inferus, bifidus, persistens.*Cor.* supera, plana, quadrifida.*Stam.* *Filamenta* subulata. *Antherae* subrotundae.*Styli* simplices. *Stigmata* capitata.*Fructus* baccatus, cum calyce coalitus, unilocularis, tetraspermus. Sem. membrana tenui obvoluta.*Saxifrageae.* Schk. t. 109. a. *Gärtn.* t. 112.

1663. PETIVERIA Plum. (1703.)

Cal. tetrasepalus.*Cor.* nulla.*Stam.* octo (sex, septem).*Styli* persistentes, demum reflexi, spinescentes.*Semen* unicum nudum, apice spinis armatum.*Chenopodaeae.* Trew. ic. Ehret. t. 67. *Gärtn.* t. 75.

CLASSIS IX. E N N E A N D R I A.

CONSPECTUS

CLASSIS NONAE.

M O N O G Y N A E.

1664 — 1671.

D I - T R I G Y N A E.

1672 — 1676.

H E X A G Y N A E.

1677. 1678.

M O N O G Y N I A.

L A U R I N A E.

1664. LAURUS Plin.

Flores diclini.

Cal. corollinus sexpartitus, deciduus.

Stam. duodecim, quorum tria sterilia. Filamenta utrinque glandula stipitata. Antherae adnatae, biloculares; loculis valva longitudinali a basi ad apicem dehiscentibus.

Styl. simplex. Stigma capitatum.

Bacca monosperma. Sem. exalbuminosum, cotyledonibus magna, iuxta basin peltatis.

Schk. t. 110.

1665. CRYPTOCARYA R. Br. (1810.)

Flores hermaphroditi.

Cal. corollinus, sexpartitus, tubo persistente, limbo deciduo.

Stam. duodecim (saepe pauciora), duplici ordine, interiora tria sterilia. Antherae biloculares; loculis valva longitudinali a basi ad apicem dehiscentibus.

Glandulae sex, cum staminibus interioribus alternantes.

Stylus simplex. Stigma obtusum.

Drupa tubo calycis aucto clauso inclusa, monosperma.

Humb. nov. gen. 7. t. 645.

1666. PERSEA Plum. (1703.)

Flores hermaphroditi.

Cal. corollinus, sexpartitus, tubo persistente, limbo deciduo, vel etiam persistente.

Stam. duodecim (vel novem) duplici ordine disposita, quorum tria interiora sterilia. Filamenta apice plana vel dilatata, basi glandulis stipitatis interstincta. Antherae quadriloculares, loculis distinctis, a basi ad apicem dehiscentibus.

Styl. simplex. Stigma capitatum, trigonum.

Drupa monosperma, calyce aucto, vel sexlobo vel integro suffulta. Semen exalbuminosum. Cotyledones crassissimae, maximae.

[*Laurus Cinnamomum* L. *Hook. exot. fl.* t. 176. *Nees amoen. bonn.* t. 1 — 3. *Gärtn.* t. 224. *Ocotea Aubl. guian.* t. 340. *Porostema Schreb.*]

1667. **TETRANTHERA** Jacqu. (1797.)

Flores dioeci, umbellati. Involucrum tetra-hexaphyllum.

Cal. corollinus, quadri-sexpartitus, vel nullus.

Stam. numero varia (6-15.), nonnulla sterilia. Filamenta apice plana, basi glandulis stipitatis interpositis. Antherae quadriloculares, loculis distinctis, a basi ad apicem dehiscentibus.

Styl. simplex. Stigma lobatum.

Bacca calyce destituta, monosperma.

Jacqu. hort. schöub. 1. t. 113. *Rosb. corom.* 1. t. 147. 148.

[*Litsea* Lam. *Juss.* *Tomex Thunb. W.* *Glaularia* L. *Evoamus Nutt.* *Laurus aggregata* Bot. mag. 2497.]

1668. **CASSYTA** Rumph. L. (1754.)

Flores hermaphroditi.

Cal. corollinus sexfidus persistens; tubo brevissimo; limbi laciniis exterioribus exilibus.

Stam. duodecim, duplici ordine; tria interiora sterilia, reliqua basi biglandulosa. Filamenta clavata. Antherae biloculares, loculis discretis.

Fructus baccatus, calyce tectus, eiusque laciniis coronatus.

Hook. exot. flor. t. 167.

1669. **ANACARDIUM** Fragos. (1572.)

Flores polygami.

Cal. pentasepalus, coloratus, deciduus.

Pet. quinque lanceolata, apice reflexa.

Stam. decem, basi calycis inserta, inferne connata: unum duplo longius et robustius. Antherae dorso adnatae, biloculares, basi bifidae, rimis longitudinalibus, dehiscentes.

Styl. subulatus. Stigma capitatum.

Fructus reniformis, cartilagineus, monospermus, insidens pedunculo pyriformi carnosus.

Terebinthaceae. [*Cassuvium* Lam. *ill.* t. 322. *Acaiuha* - *Gärtn.* t. 40.]

- 1670. *PLEGOREHIZA* Molin. (1782.)
Cal. corollinus, campanulatus, integer.
Stam. brevissima.
Stylus abbreviatus.
Fructus oblongus monospermus.
- 1671. *THALASSIA* Banks. König. (1806.)
Flores dioeci.
Masculorum spatha uniflora, monophylla, bifida; laciniis oblongis obtusis.
Cal. trisepalus, obtusus.
Cor. nulla.
Stam. Antherae sessiles, lineari-lanceolatae, conniventes.
Flores feminei ignoti.
 — — [*Zostera ciliata* et *stipulacea* Forsk.]

D I - T R I G Y N I A

- 672. *MERCURIALIS* Plin.
Flores dioeci.
Cal. tripartitus, patens.
Cor. nulla.
Filamenta capillaria. *Antherae* didymae, globosae. In flore femineo duo filamenta germini adstant.
Styli duo reflexi.
Cocci duo bivalves, monospermi.
Tricoccae. Schk. t. 332.
- 673. *RHEUM* Diosc.
Cal. corollinus sexpartitus, marcescens.
Filamenta basi corollae inserta. *Antherae* ovales, biloculares.
Styli subnulli. *Stigmata* reflexa.
Achenium triquetrum; alatum.
Polygoneae. Schk. t. 110. *Reichenb. hort.* t. 129.
- 1674. *ERIOGONUM* Mx. (1803.)
Flores fasciculati, ex involucro campanulato.
Cal. subcampanulatus, sexfidus; laciniis obtusis, interioribus paulo maioribus.
Filamenta capillaria, calyce longiora. *Antherae* ovatae.
Styli subnulli. *Stigmata* filiformia.
Achenium triquetrum, apterum.
Polygoneae. Mx. fl. bor. amer. t. 24. [*Espinosa* Lag.]
- 1675. *TRIPLARIS* Löff. (1758.)
Flores dioeci, spicati.
Cal. campanulatus, tri-sexfidus.
Stam. basi calycis inserta. *Filamenta* capillaria. *Antherae* cordatae, biloculares. Numerus variat (9-12).

Styli tres acuti. Stam. tria sterilia in flore femineo.

Achenium triquetrum, calyce tectum, cuius laciniae tres elongantur.

Polygonaceae. Jacqu. amer. t. 173. f. 5. Aubl. guian. t. 347.

1676. *PLEEA* Mx. (1803.)

Flores spicati, spathacei.

Cal. corollinus sexpartitus, patentissimus, lacinia linearilanceolata.

Stam. hypogyna. Filamenta subulata. Antherae oblongae, biloculares.

Stigmata sessilia linearia.

Caps. tres connatae, bivalves, polyspermae, intus dehiscentes.

Melanthiaceae. Mx. fl. bor. amer. t. 25.

HEXAGYNIA.

1677. *BUTOMUS* Theophr.

Flores umbellati.

Cal. corollinus, hexasepalus, marcescens.

Stam. perigyna. Filamenta subulata. Antherae ovatae biloculares.

Germina sex, apice in stigmata applanata reflexa abeuntia.

Caps. totidem folliculares, polyspermae. Sem. albuminosa.

Embryo punctiformis.

Hydrocharideae. Schk. t. 111.

1678. *HYDROCHARIS* L. (1737.)

Flores dioeci.

Cal. trisepalus, bracteatus, coloratus.

Petala tria, obovata.

Stam. triplici ordine stylo abortienti inserta. *Stigmata* tria filiformia tribus staminibus interioribus adnata.

Antherae subrotundae, in ambitu dehiscentes.

In flore femineo *Styli* sex bipartiti, cum totidem filamentis abortientibus, quorum tria truncata, tria stylos aequantia.

Caps. infera, sexlocularis, polysperma. Sem. marginata, albuminosa. Embryo lateralis linearis.

Hydrocharideae. Schk. t. 333. Sturm. fl. germ. Ann. mus. 9. t. 30. vol. 17. t. 9. f. 44. 45.

CLASSIS X. DECANDRIA.

CONSPECTUS

CLASSIS DECIMAE.

MONOGYNAE.

- I. Floribus perfectis.
 - A. Corolla monopetala.
 - a. Fructibus superis. 1679 — 1694.
 - b. Fructibus inferis. 1695 — 1698.
 - B. Corolla pleiopetala.
 - a. regulari.
 - α. Fructibus superis. 1699 — 1801.
 - β. Fructibus inferis. 1802 — 1809.
 - b. Corolla irregulari. 1810 — 1859.
- II. Floribus imperfectis.
 - A. Fructibus superis. 1860 — 1876.
 - B. Fructibus inferis. 1877 — 1880.

DIGYNAE.

- I. Floribus perfectis 1881 — 1193.
- II. Floribus imperfectis 1894 — 1896.

TRIGYNAE.

1897 — 1916.

TETRAGYNAE.

1917.

PENTAGYNAE.

1818 — 1941.

DECAGYNAE.

1942 — 1944.

MONOGYNIA.

I. Flores perfecti.

A. COROLLA MONOPETALA.

a. *Fructus superi.*

STYRACEAE.

1679. STYRAX Theophr. Diosc.

Cal. subcampanulatus quinquedentatus.

Cor. basi subcampanulata, profunde quinque- (tri- septem-) fida.

Stam. exserta. Filamenta tubo corollae adnata, basi nonnunquam in annulum coalita. Antherae lineares, biloculares, rimis longitudinalibus interioribus dehiscentes.

Stylus simplex. Stigma sublobatum.

Drupa exsucca, mono- (rarius di- aut tri-) pyrena.

Tournef. inst. t. 369. Gärtn. t. 59. Cav. diag. t. 188. Humb. plant. aequin. 2. t. 101. [Strigilla Cav. diss. t. 201. Foveolaria R. et P. Tremanthus Pers. Trichogamila P. Brown.]

1680. DICLIDANTHERA Mart. (1826.)

Cal. quinquepartitus.

Cor. infundibuliformis, limbo quinquefido.

Stam. Antherae subrotundae, sessiles in fauce corollae, biloculares, e basi sursum et deorsum bivalves.

Styl. filiformis. Stigma subincrassatum.

Bacca globosa, quinquelocularis, pentasperma.

Mart. nov. gen. 2. t. 196. 197.

1681. CYRTA Lour. (1790.)

Cal. quinquedentatus, persistens.

Cor. campanulata, quinquepartita.

Stam. Filamenta brevia, basi dilatata, basi corollae inserta. Antherae oblongae, reflexo-incurvae, dorso adnatae.

Styl. subulatus. Stigma simplex.

Drupa oblonga, utrinque attenuata, incurva, monosperma. Nux oblonga, acuta, sulcata.

E R I C E A E.

1682. ARBUTUS Virgil.

Cal. quinquepartitus.

Cor. urceolata, limbo quinquedentato reflexo.

Stam. basi corollae inserta, inclusa. Filamenta brevia, basi dilatata. Antherae biloculares, apice poris duobus dehiscentes, dorso aristatae.

Germen disco decangulari cinctum.

Stylus simplex. Stigma obtusum.

Bacca quinquelocularis, polysperma.

Engl. bot. 2377. Gärtn. t. 59. Bot. reg. 113. Humb. nov. gen. 3. t. 260.

1683. ARCTOSTAPHYLOS Galen. Adans. (1763.)

Cal. quinquepartitus.

Cor. urceolata; limbo quinquefido, reflexo.

Stam. basi corollae inserta, inclusa. Filamenta brevia. Antherae biloculares, apice poris duobus dehiscentes, dorso aristatae.

Germen disco hypogyno insertum.

Stylus simplex. Stigma obtusum.

Drupa subglobosa, stylo coronata; putamine quinqueloculari, loculis monospermis.

Humb. nov. gen. 3. t. 258. 259. [Arbutus Uva ursi et alpina L. Fl. dan. 33. 73. Sturm. fl. germ. Engl. bot. 714. 2030.]

1684. GAUTIERA Kalm. (1751.)

Cal. bracteatus, campanulatus, quinquefidus.

Cor. ovata, semiquinquefida; limbo parvo, revoluta.

Stam. receptaculo inserta, corolla breviora. Filamenta subulata. Antherae bicornes, corniculis bifidis.

Corpuscula glandulosa quinque aut decem perigyna.

Styl. filiformis. Stigma obtusum.

Caps. calyce baccato tecta, quinquelocularis, quinquevalvis, apice quinquefariam dehiscens; placentis singulorum loculorum polyspermis.

Andr. repos. 116. *Humb. nov. gen.* 3. t. 261. 262. [Kalmius primus dicavit hoc genus amico, medico, canadensi, qui semet Gautier scripsit. *Mém. étrang. de l'ac. de Paris*, 2. p. 378. Linnaeus *Gaultheriam*, alii *Gualtheriam* scripserunt.]

1685. EPIGAEA L. (1751.)

Cal. bracteis tribus suffultus, quinquepartitus, laciniis lanceolatis.

Cor. tubulosa; limbo hypocrateriformi quinquefido.

Stam. basi corollae inserta, tubum aequantia. Filamenta filiformia. Antherae oblongo-lineares, rimis geminis longitudinalibus dehiscentes.

Stylus filiformis, Stigma sublobatum.

Caps. quinquelocularis, quinquevalvis, polysperma; valvis inflexis dissepimenta formantibus; columna centrali quinquepartita.

Andr. repos. 102. *Bot. reg.* 201.

1686. ENCYANTHUS * (Enkianthus Lour. 1790.)

Cal. bracteis coloratis stipatus, quinquefidus, persistens.

Cor. campanulata, quinquefida. Foveolae nectariferae quinque in basi corollae.

Stam. basi corollae inserta. Filamenta inferne pilosa dilatata. Antherae bicornes, apice poro gemino dehiscentes.

Styl. filiformis, persistens.

Capsula.

Bot. mag. 1649. *Bot. reg.* 884. 885. [*Melidora Salisb. horticult. transact.* 2. p. 156.]

1687. ANDROMEDA L. (1737.)

Cal. quinquepartitus, coloratus, persistens.

Cor. campanulata, ampullacea vel urceolata, limbo quinqueudentato, reflexo.

Stam. basi corollae inserta, inclusa. Filamenta subulata. Antherae bicornes, apice poro gemino dehiscentes, (raro dorso aristatae).

Styl. cylindricus, persistens. Stigma obtusum.

Caps. quinquelocularis, quinquevalvis, polysperma; valvae

septis contrariae, medio dehiscentes; placenta centralis, quinqueloba.

Engl. bot. 713. *Gärtn.* t. 63. 178. Bot. mag. 667. 905. 970. 1095. 1286. 1566. 1579. *Humb.* nov. gen. 3. t. 263. [Nominis rationem potius exposuit Linnæus in flora lapp. n. 163.]

1688. *LYONIA* Nuttall. (1818.)

Cal. quinquedentatus.

Cor. globoso-urceolata, pubescens.

Stam. inclusa. Antherae bicornes, muticae.

Styl. cylindricus.

Caps. pentagona, quinquelocularis, quinquevalvis; valvae medio septiferae, marginibus ope valvarum accessoriarum clausis; placenta centralis quinqueloba. Sem. imbricata, subulata.

[*Andromeda paniculata* L., frondosa *Pursh.*, ferruginea *Ait.*]

1689. *RHODODENDRON* L. (1758.)

Cal. quinquefidus, persistens.

Cor. infundibuliformis, limbo quinquepartito patente.

Stam. hypogyna, declinata. Filamenta filiformia. Antherae ovals.

Styl. filiformis. Stigma capitatum.

Caps. quinquelocularis; valvis inflexis dissepimenta formantibus; columna centralis quinqueloba. Sem. exilia, linearia, membrana obvoluta.

Schk. t. 117. *Gärtn.* t. 63. *Andr.* repos. 4. 36. Bot. mag. 37. 194. 195.

1690. *KALMIA* L. (1751.)

Cal. quinquefidus, persistens.

Cor. hypocrateri-infundibuliformis; limbo patente quinquepartito. Foveolae quinque sub limbo antheras recipientes, extus totidem tuberculis prominulae.

Stam. basi corollae inserta, eamque longitudine subaequantia. Filamenta filiformia. Antherae ovals.

Styl. filiformis, elongatus, persistens. Stigma obtusum.

Caps. quinquelocularis, quinquevalvis; loculis polyspermis. Placenta centralis, quinqueangularis.

Schk. t. 116. Bot. mag. 138. 175. 177.

1691. *PTEROSPORA* Nuttall. (1818.)

Cal. quinquepartitus.

Cor. ovata, limbo quinquedentato, reflexo.

Stam. hypogyna, inclusa. Filamenta subulata, basi dilatata. Antherae biloculares, basi appendiculatae, transverse dehiscentes.

Styl. columnaris, brevis. Stigma subquinquelobum.

Caps. quinquelocularis, quinquevalvis. Placenta centralis quinqueloba. Sem. numerosa, minuta, apice alata.

1692. *INOCARPUS* Forst. (1775.)*Cal.* bipartitus.*Cor.* infundibuliformis, limbo quinquepartito.*Stam.* tubo adnata, duplici serie. *Antherae* erectae ovales biloculares.*Styl.* nullus. *Stigma* sessile, subrotundum.*Drupa* monosperma; nuce fibrosa, venoso-reticulata.*Sapoteae*. Forst. char. t. 33. *Rexb.* corom. 3. t. 263.1693. *POTALIA* Aubl. (1775.)*Cal.* coriaceus, quinquelobus, coloratus, persistens.*Cor.* tubulosa, limbo decemfido, laciniis oblique imbricatis.*Stam.* inclusa, basi membrana annulari nexa. Filamenta breviter subulata. *Antherae* lineares, biloculares, rimis longitudinalibus dehiscentes.*Styl.* filiformis. *Stigma* capitatum.*Bacca* bilocularis, polysperma.*Potalieae* Mart. *Aubl. guian.* t. 151. Mart. nov. gen. t. 170. [*Nicandra* Schreb.]1694. *CODON* Royen. L. (1767.)*Cal.* decempartitus, laciniis alternis minoribus.*Cor.* campanulata, basi torulosa; limbo decempartito, aequali.*Stam.* basi squamis conniventibus et receptaculum obtinentibus instructa. *Antherae* crassae.*Styl.* simplex. *Stigmata* duo, setacea, divergentia.*Fructus* baccatus, bilocularis, polyspermus. Sem. papillosa.— — *Meerburgh* ic. t. 73. *Gärtn.* t. 95.b. *Fructus inferi.*1695. *CERATOSTEMA* Juss. (1789.)*Cal.* turbinatus, quinquefidus; laciniis magnis persistentibus.*Cor.* coriacea, tubulosa; limbo erecto, quinquefido.*Stam.* perigyna. Filamenta breviter. *Antherae* longissimae, erectae, apice bifidae.*Styl.* simplex.*Caps.* tomentosa, quinquelocularis.*Ericaeae.*1696. *THIBAUDIA* Pav. Kunth. (1818.)*Cal.* quinquefidus.*Cor.* tubulosa, basi ventricosa, limbo suberecto, quinquefido.*Stam.* calyci inserta. Filamenta breviter. *Antherae* longissimae, erectae, apice bifidae.*Styl.* simplex. *Stigma* obtusum.*Bacca* calyce vestita, quinquelocularis; loculis polyspermis.

Sem. angulata.

Ericaeae. *Hamb.* nov. gen. 3. t. 254 - 256.

1697. *LUSSACIA* * (Gay - *Lussacia* Kunth. 1818.)*Cal.* germini adnatus, limbo quinquefido.*Cor.* tubulosa, basi ventricosa; limbo quinquefido.*Stam.* calyci inserta, inclusa. Filamenta brevia membranacea. Antherae dorso affixae, apice in tubulos duos fissae.*Stylus* simplex. Stigma capitatum.*Drupa* subglobosa, calyce vestita, decemlocularis, decasperma. Sem. lenticularia.*Ericaceae.* Humb. nov. gen. 3. t. 257.1698. *POLYPHRAGMON* Desfont. (1820.)*Cal.* quinquedentatus persistens.*Cor.* infundibuliformis; limbo decemfido.*Stam.* in medio tubo. Antherae sessiles, lineares.*Styl.* crassus. Stigmata hex aut plura, acuta, recurva.*Bacca* multilocularis, polysperma. Placenta centralis.*Rubiaceae.* Mém. du mus. 6. t. 2.

B. COROLLA PLEIOPETALA,

a. *regularis.*α. *Fructus superi.*

SALICARIAE.

1699. *ACISANTHERA* P. Brown. (1756.)*Cal.* basi ventricosus, limbo quinquefido persistente.*Pet.* quinque obovata, fauci calycis inserta.*Stam.* ibi inserta, exstantia. Antherae sagittatae, versatiles.*Stylus* brevis. Stigma acutum.*Caps.* bilocularis, polysperma, calyce tecta, limbique laciniis coronata. Sem. subrotunda.*P. Brown.* jam. t. 22. f. 1.

MELASTOMEAE.

1700. *MELASTOMA* J. Burm. (1737.)*Cal.* ovatus, saepe squamatus, limbo quinquefido, deciduo.*Pet.* quinque, calycis apici inserta, cum laciniis calycis alternantia.*Stamina* ibidem inserta. Filamenta aequalia. Antherae elongatae, arcuatae, apice poro dehiscentes, basi bicalcaratae aut bisetosae.*Styl.* simplex.*Caps.* baccata quinquelocularis. Sem. cochleata.*Bot. reg.* 672. *Gärtn.* t. 126. *Labill.* nov. Caledon. t. 64.

1701. *PLEUROMA* Don. (1823.)

Cal. ovatus, bracteis binis suffultus, limbo quinquelobo deciduo.

Pet. obovata, quinque.

Stam. calycis margini inserta. Filamenta glabra. Antherae elongatae, arcuatae, basi biauriculatae.

Styl. simplex.

Caps. baccata, quinquelocularis. Sem. cochleata.

[? *Melastoma heteromalla* Bot. reg. 644. *Rhexia vaminea* ib. 664.]

1702. *LASIANDRA* Cand. (1828.)

Cal. ovatus, setosus aut sericeus, bracteis binis munitus; limbo quinquefido deciduo, laciniis angustis elongatis.

Petala quinque obovata.

Stam. calycis margini inserta. Filamenta pilosa. Antherae arcuatae, basi stipitiformi biauriculata aut papilloso-barbata.

Styl. simplex.

Caps. sicca, quinquelocularis. Sem. pauca, angulata.

Cand. *Melast.* t. 7. [*Rhexia Bonpl.* *Rhex.* t. 36. 51. Bot. reg. 323.]

1703. *CHAETOGASTRA* Cand. (1828.)

Cal. turbinatus, ebracteatus; limbo quinquefido persistente.

Pet. quinque obovata.

Stam. calycis margini inserta, similia. Filamenta glabra. Antherae rectiusculae, apice poro dehiscentes, basi bituberculatae.

Stylus simplex.

Caps. libera, quinquelocularis. Sem. cochleata.

Cand. *Melast.* t. 8. Bot. mag. 2836. [*Meriania Vent.* sel. t. 34. *Rhexia Vahl.* ic. amer. t. 15. *Bonpl.* *Rhex.* t. 1. 6. 7. 8. 10. 13. 15. 21. 31. 43.]

1704. *OSBEKIA* L. (1757.)

Cal. ovatus, stellato-pubescent; limbo quinque (quadri-) fido. Appendices inter lacinias exteriores variae formae.

Pet. quinque (quatuor).

Stam. aequalia. Filamenta glabra. Antherae breviter rostratae, basi biauriculatae.

Styl. simplex.

Caps. libera, quinque- (quadri-) locularis, polysperma. Sem. cochleata.

Osbek. it. t. 2. Bot. reg. 542. 565. 574. Bot. mag. 2838. *Hook.* fl. exot. t. 31. 37. [*Rhexia Bonpl.* *Rhex.* t. 45.]

1705. *TIBUCHINA* Aubl. (1775.)

Cal. turbinatus, squamis imbricatis vestitus, basi involucri duplici cinctus, utroque e bracteis binis connatis; limbo quinquefido, laciniis lanceolatis.

Pet. quinque, ovalia.

Stam. Filamenta glabra. Antherae basi in aurículas binas obtusas productae.

Styl. simplex.

Caps. libera, quinquelocularis. Sem. cochleata.

Aubl. guian. t. 177.

1706. *TRISTEMMA* Juss. (1789.)

Cal. tubulosus, triplici setarum annulo cinctus; limbo quinquefido.

Pet. quinque unguiculata.

Stam. calyci inserta. Filamenta glabra. Antherae subarctatae, basi subauriculatae.

Styl. simplex.

Bacca calyce vestita, quinquelocularis, polysperma.

Ventén. sel. t. 35.

1707. *MICROLICIA* Don. (1823.)

Cal. subglobosus; limbo quinquefido persistente, laciniis subulatis.

Pet. obovata.

Stam. calycis margini inserta. Filamenta glabra. Antherae ovatae, breviter rostratae, apice poro dehiscen-tes, inferne appendicula stipitiformi obtuse calcaratae, alternatim longiores.

Styl. simplex.

Caps. libera, sicca, trilocularis, trivalvis. Sem. cochleata.

[*Melastoma Aubl. guian.* t. 155. *Rhexia Bonpl. Rhex.* t. 59.]

1708. *TREMBLEYA* Cand. (1828.)

Cal. urceolatus; limbo quinquefido, laciniis subaristatis.

Pet. quinque ovalia.

Stam. calycis margini inserta, inaequalia. Antherae alternae ovatae, breviter rostratae, basi ligula obcordata vel spatulata; alternae basi subnudaе.

Styl. simplex.

Caps. glabra, quinquelocularis. Sem. cochleata?

1709. *DIPLOSTEGIUM* Don. (1823.)

Calycis limbus quinquefidus persistens, in calyptra duplici cucullata hispida inclusus.

Pet. quinque.

Antherae subaequales, basi biauriculatae.

Caps. baccata, quinquelocularis.

1710. *LEANDRA* Radd. (1820.)

Cal. bracteatus, urceolatus; limbo duplicato, laciniis interioribus membranaceis, exterioribus rigidioribus subulatis.

Pet. lineari-lanceolata.

Stam. Antherae lineares, basi obsolete auriculatae.

Styl. basi hirsutus.

Bacca trilocularis. Sem. ovata, angulata.

1711. *Tschudia* Cand. (1828.)

Cal. globosus, dentibus quinque minimis subulatis persistentibus coronatus.

Stam. Filamenta planiuscula. Antherae lineari-oblongae, apice poro dehiscentes, basi exauriculatae.

Styl. filiformis. Stigma capitatum, subbilobum.

Caps. libera, globosa, membranacea, quinquelocularis.

Sem. minima incurva, appendicula celluloso-lacunosa.

Cand. Melastom. t. 9.

1712. *Clidemia* Don. (1823.)

Cal. ovatus, nudus; limbo quinquefido, persistente, laciniis angustis acutis.

Stam. aequalia. Antherae basi constrictae, subbiauriculatae, apice poro dehiscentes.

Germen setis circinatis coronatum.

Styl. filiformis.

Caps. baccata, quinquelocularis. Sem. ovata, subangulata.

[*Melastoma* Aubl. guian. t. 165 — 167. *Boiss.* *Melast.* t. 3. 5-52. 59.]

1713. *Myriaspora* Cand. (1828.)

Cal. ovato-globosus, hispidissimus; limbi laciniis apice coalitis, cuculli formam referentibus, dein irregularibus, lanceolatis.

Stam. Filamenta brevia. Antherae crassae, obtusae, basi gibbae.

Styl. filiformis.

Caps. baccata, quinquelocularis. Sem. numerosa, angulata.

1714. *Tococa* Aubl. (1775.)

Cal. oblongus, limbo quinquefido persistente.

Pet. obovata.

Stam. aequalia. Filamenta glabra. Antherae lineares, acutae, basi subsagittatae, rectae.

Styl. brevis. Stigma peltatum.

Bacca trilocularis. Sem. angulata.

Aubl. guian. t. 174.

1715. *Maieta* Aubl. (1775.)

Cal. oblongus, pentagonus, coloratus, bracteis imbricatis suffultus; limbo quinquefido, persistente.

Pet. subrotunda, unguiculata.

Stam. Filamenta brevia. Antherae lanceolatae, apice in ligulam productae, basi bicalcaratae.

Stylus brevis. Stigma capitatum.

Bacca calyce vestita, quinquelocularis. Sem. minutissima.

Aubl. guian. t. 176.

1716. *Huberia* Cand. (1828.)

Cal. infundibuliformis, limbo quinquefido.

Pet. obovata.

Stam. Antherae apice productae, acuminatae, dorso setigerae.

Styl. filiformis.

Caps. baccata, quinquelocularis.

Cand. Melast. t. 10. [Maleta Vent. sel. t. 32.]

1717. *CALYCOGONIUM* Cand. (1828.)

Cal. subglobosus angulatus, stellato - punctatus; limbi laciniis linearibus elongatis.

Pet. obovata.

Stam. Antherae oblongae apice poro dehiscentes, basi exauriculatae.

Bacca angulata, quadri-quinquelocularis, laciniis calycis coronata. Sem. angulata.

[Melastoma calycopteris Rich. in Bonpl. Melast. t. 42.]

1718. *HETEROTRICHUM* Cand. (1828.)

Cal. globosus, limbo quinque - octofido, laciniis elongatis acuminatis.

Pet. quinque, octo, ovalia.

Stam. aequalia. Filamenta glabra. Antherae oblongae, apice poro dehiscentes, basi exauriculatae.

Styl. filiformis.

Caps. baccata, quinque - octolocularis.

[Melastoma Plum. ic. t. 141.]

1719. *CONOSTEGIA* Don. (1823.)

Cal. oblongus; limbo circumscisso, integro.

Pet. oblonga.

Stam. Antherae basi biauriculatae.

Styl. brevis. Stigma peltatum.

Caps. baccata, quinquelocularis. Sem. angulata.

[Melastoma Bonpl. Melastom. t. 31. 41. 46. 51. 54. 55. 57. 60.]

1720. *DIPLOCHITON* * (Diplochita Cand. 1828.)

Cal. junior bracteis duabus amplis obtectus; limbo quinquedentato persistente.

Pet. oblonga.

Stam. Antherae apice poro dehiscentes, basi biauriculatae.

Styl. filiformis. Stigma capitato - peltatum.

Caps. sicca quinquelocularis. Sem. angulata.

[Fothergilla Aubl. guian. t. 175. Melastoma Bonpl. Melast. t. 18. 32. 33. Chitonita Don.]

1721. *MICONIA* Ruiz et Pav. (1794.)

Cal. subglobosus vel ovatus; limbo quinquedentato persistente, intus membrana lata aucto.

Pet. obovata.

Stam. Antherae oblongo-lineares, basi breviter auriculatae.

Styl. filiformis. Stigma obtusum.

Caps. baccata, quinquelocularis. Sem. triquetra.

[Melastoma Aubl. guian. t. 154. 156. 158. 159. 170. Bonpl. Melast. t. 1. 2. 6. 7. 11. 15. 16. 17. 21. — 27. 29. 35. 40. 43. 47. 49. 53. 56. 62. Bot. reg. 363.]

1722. CREMANIUM Don. (1823.)

Cal. campanulatus; limbo quinquedentato persistente.

Pet. obovata.

Stam. Antherae breviores, apice biporosae, basi attenuatae exauriculatae.

Styl. filiformis. Stigma peltatum.

Caps. baccata, tri-quinquelocularis. Sem. angulata.

[*Melastoma Bonpl.* Melast. t. 22. 19. 48.]

1723. MERIANIA Sw. (1800.)

Cal. campanulatus, laciniis membranaceis dorso dente elongato subulato notatis.

Pet. obovata.

Stam. Antherae oblongae, utrinque obtusae, apice poro gemino hiantes.

Styl. clavatus.

Caps. libera, quinquelocularis. Sem. angulata. Placentae lunatae.

Swartz. fl. ind. occid. t. 15.

1724. LAVOISERIA Cand. (1828.)

Cal. turbinate-oblongus, laciniis limbi quinque aut decem acutis.

Pet. obovata.

Stam. Filamenta planiuscula. Antherae oblongae utrinque obtusae, apice appendicula ovata forata, processu basilari arcuato, ad articulationem cum filamento, bifobo.

Caps. libera, quinque-decemlocularis. Sem. angulata.

Cand. Melast. t. 2.

1725. DAVIA Cand. (1828.)

Cal. campanulatus, limbo integro membranaceo.

Pet. quinque.

Stam. Antherae lineares, elongatae, apice acuminata-rostratae, basi appendicula calcarata setiformi.

Styl. filiformis.

Caps. quinquelocularis.

Cand. Melast. t. 3.

1726. GRAFFENRIEDIA Cand. (1828.)

Calycis tubus oblongus, limbus campanulatus, obtusissime quinquedentatus.

Pet. obovata.

Antherae lineares, acutae, apice poro unico instructae, appendicula basilari setacea vel obtusa.

Styl. filiformis.

Caps. ovata, quinquevalvia, apice dehiscens. Sem. angulata.

[*Rhexia Bonpl.* Rhex. t. 25. 34.]

1727. CENTRONIA Don. (1823.)

Cal. oblongus, undique retrorsum setosus; limbo coarctato integro.

Pet. quinque.

Antherae longissimae rostratae, basi processu longo subulato instructae.

Styl. in collo ovarii tubuloso elongato inclusus. Stigma obtusum.

Caps. quinquelocularis.

1728. *RHYNCHANThERA* Cand. (1828.)

Cal. ovato-globosus; laciniis subulatis persistentibus.

Pet. quinque.

Stam. inaequalia; alternorum antherae abortientes aut minores, alternorum longissime rostratae, appendicula stipitiformi basi auriculata.

Styl. simplex.

Caps. libera, quinquelocularis. Sem. angulata.

[*Melastoma Aubl. guiau*, t. 160. *Rhexia Bonpl. Rhex.* t. 11.]

1729. *CAMBESSEDESIA* Cand. (1828.)

Cal. ovata-globosus; laciniis linearibus persistentibus.

Pet. obovata.

Stam. aequalia. *Antherae* falcatae subrostratae, basi gibbae; appendicula exili auriculata.

Styl. filiformis.

Caps. libera, trilocularis. Sem. ovata, angulata.

[*Rhexia Bonpl. Rhex.* t. 56. 58. 60.]

1730. *TRIBLEMMMA* R. Br. (1822.)

Cal. campanulatus, lobis obtusis, saepe in limbum integrum confluentibus.

Pet. obovata.

Stam. subinaequalia. *Antherae* ovatae, obtusae; apice porosae, basi attenuatae, exappendiculatae.

Caps. trigona, trivalvis; valvis uncinatis, sub apice circumscissis. Sem. cuneato-triquetra scabra.

[*Rhexia Bonpl. Rhex.* t. 53 — 55. *Bertolonia Radd. Cand.*]

1731. *CHAENOPLEURA* Rich. Cand. (1828.)

Cal. germi adnatus; limbo obtuse quinque-dentato.

Pet. subrotunda.

Stam. petalis vix longiora. *Antherae* longitudinaliter birimosae; appendicula biauriculata.

Styl. clavatus. Stigma orbiculatum, subumbilicatum.

Bacca? tri-quadrilocularis.

[*Melastoma Bonpl. Melast.* t. 30.]

1732. *ASTRONIA* Blum. (1827.)

Cal. germi adnatus; limbo persistente quinque-sex-dentato.

Pet. quinque aut sex.

Stam. decem, duodecim. Filamenta membranaeco-compressa. *Antherae* longitudinaliter dehiscentes, dorso (subrostratae) carnosae.

Styl. stigmatē peltato.

Bacca exsucca, umbilicata, tri-quadrilocularis, polysperma. Sem. paleacea, arillo marginata membranaceo.

T E R E B I N T H E A E.

1733. *BOSWELLIA* Colebr. (1814.)

Cal. quinquedentatus.

Pet. quinque.

Urceolus glandulosus crenulatus perigynus.

Styl. cylindricus. Stigma trilobum.

Caps. trigona, trilocularis, trivalvis. Sem. solitaria, marginato - alata.

Asiat. research. 9. p. 377. Roxb. corom. 3. t. 207. Linn. transact. 15. t. 5.

1734. *PEGIA* Colebr. (1827.)

Cal. quinquelobus, minutus, persistens.

Pet. subrotunda.

Annulus carnosus perigynus.

Stam. externa basi annuli perigyni inserta; alterna breviora. Filamenta subulata. Antherae subrotundae, biloculares.

Stylus conicus. Stigma simplex.

Bacca globosa, monosperma.

1735. *GABUGA* Roxb. (1819.)

Cal. quinquedentatus.

Pet. quinque, calyci inserta.

Glandulae quinque perigynae.

Stam. calyci inserta.

Stylus brevis. Stigma quinquelobum.

Drupa subdisperma.

Roxb. corom. 3. t. 208.

1736. *BUCHANANIA* * (1801.)

Cal. quinquedentatus,

Pet. sub disco inserta, revoluta.

Discus urceolatus perigynus; intus dentibus subulatis quinque munitus.

Stam. petalis breviora.

Stylus brevis. subulatus.

Drupa monosperma, putamine bivalvi.

Roxb. corom. 3. t. 262. [Cambessedea Kunth.]

1737. *BURSERIA* Jacqu. (1763.)

Flores polygami.

Cal. tri-quinquelobus.

Pet. tria, quinque, patentia.

Discus annularis crenatus.

Filamenta subulata.

Stylus brevis, stigmatē trifido.

Drupa putamine trivalvi, submonosperma. Sem. pellicula pulposa cinctum.

Jacqu. amer. t. 65. Linn. transact. 15. t. 4. f. 1.

1738. *MARIGNIA* Commers. Kunth. (1823.)

Cal. quinquefidus, persistens,

Pet. quinque, calyce duplo longiora,

Discus integer.

Stam. calycem aequantia.

Stigma subsessile subquinelobum.

Drupa stigmatē coronata, mono- (penta-) pyrena. Sem. pendula, exalbuminosa.

[*Dammara Gärtn. t. 103.*]

1739. *MAURIA* Kunth. (1825.)

Cal. urceolatus, quinquedentatus, persistens.

Pet. patentia.

Discus orbicularis hypogynus.

Stam. sub disco inserta. Antherae biloculares,

Styl. crassus, brevissimus. *Stigma* trilobum.

Fructus ellipticus, compressus, carnosus. Semen reniforme, exalbuminosum.

Humb. nov. gen. 7. t. 605. 606. [Mangifera axillaris Desrousse.]

1740. *RHINOCARPUS* Berter. Kunth. (1825.)

Flores polygami.

Cal. quinquepartitus, deciduus.

Pet. calyci inserta, patentissima.

Stam. calyci inserta, inaequalia; duo vel quatuor perfecta, reliqua abortientia. Filamenta nonnulla inferne connata, petalis hinc adnata. Antherae biloculares.

Styl. sublateralis. *Stigma* obtusum.

Fructus compressus monospermus, pedicello crasso arcuato suffultus.

Humb. nov. gen. 7. t. 601.

1741. *CYRTOCARPA* Kunth. (1825.)

Flores polygami.

Cal. quinquepartitus, lacinii obtusis.

Pet. ovalia, patentissima, calycem duplo superantia.

Discus perigynus orbicularis, crenatus.

Stam. sub disco inserta, corolla dimidio breviora. Antherae biloculares.

Styl. simplex. *Stigma* subquadrifidum.

Drupa monosperma, calyce, corolla filamentisque persistentibus suffulta.

Humb. nov. gen. 7. t. 609.

A U R A N T I A.

1742. *FERONIA* Corr. (1800.)

Cal. planus, quinquepartitus.

- Pet.* oblonga.
Stam. Filamenta basi dilatata, villosa. Antherae oblongae.
Stigma quinquelobum.
Pomum quinqueloculare, loculis polyspermis.
Roxb. corom. 2. t. 141.
743. *GLYCOSMIS* Corr. (1805.)
Cal. quinquefidus.
Pet. quinque.
Stam. Filamenta complanata subulata. Antherae ovales.
Stylus brevis cylindricus. *Stigma* simplex.
Germen quinqueloculare. *Fructus* carnosus uni-bilocularis. *Sem.* pendula.
[Limonia Roxb. corom. 1. t. 84. 85. *Sclerostylis Blum.]*
744. *LIMONIA* P. Brown. (1756.)
Cal. quinquefidus.
Pet. quinque.
Stam. Filamenta subulata. Antherae lineares, erectae.
Styl. cylindricus. *Stigma* capitatum.
Pomum quinqueloculare, pentaspermum.
Roxb. corom. 1. t. 86. *Bot. mag.* 2416.
745. *COOKIA* Sonnerat. (1782.)
Cal. quinquepartitus.
Pet. quinque, navicularia, villosa.
Stam. Filamenta linearia. Antherae subrotundae.
Pomum abortu biloculare, loculis monospermis.
Sonner. it. ind. 2. t. 131. *Jacqu. schöubr.* 1. t. 101. [*Quinaria et Aulacia Lour.*]
746. *MURRAYA* L. (1771.)
Cal. quinquepartitus.
Pet. quinque unguiculata, erecto-patentia.
Stam. Filamenta planiuscula, erecta, conniventia. Antherae biloculares subrotundae.
Styl. crassus. *Stigma* capitatum.
Bacca submonosperma.
Murr. comm. gotting. 9. t. 1. *Hook. fl. exot.* 2. t. 134.
[Marsana Sonner. it. ind. 2. t. 139. *Claucena N. L. Burm.*
ind. t. 29. *Chalcas Lour.* *Bergera L.*]
747. *MICROMELUM* Blum. (1825.)
Cal. urceolatus, integer, persistens.
Pet. quinque, linearia, patentia.
Stam. alterna breviora. Filamenta subulata. Antherae subrotundae, didymae.
Styl. crassus, stigmate obtuso.
Bacca exsucca, dissepimentis quinque chartaceis intortis lamellata, mono-disperma.

E R I C E A E.

748. *LEDUM* Lobel. (1570.)
Cal. quinquedentatus, persistens.

Pet. quinque patentia.

Stam. receptaculo inserta. Filamenta capillaria, recta.

Styl. cylindricus. Stigma quinquelobum.

Caps. quinquelocularis, quinquevalvis; valvis a basi sursum dehiscentibus. Placenta quinqueangularis, centralis.

Sem. linearia, membrana tenera reticulata obvoluta.

Schk. t. 117.

1749. *LEIOPHYLLUM* Pers. (1805.)

Cal. quinquepartitus; laciniis lanceolatis.

Pet. quinque ovata, obtusa, patentia.

Stam. receptaculo inserta. Filamenta capillaria, corollae longiora. Antherae ovatae, didymae.

Styl. cylindricus. Stigma truncatum.

Caps. trilocularis, trivalvis. Sem. ovata, minuta.

[*Ledum buxifolium* Berg. in act. petropol. 1777. t. 3. f. 2. *Ammyrsine* Pursh. Bot. reg. 531. *Fischera* Sw. in act. soc. mosqu. 5.]

1750. *CLETHRA* L. (1737.)

Cal. quinquepartitus, persistens.

Pet. quinque, spathulata, concaviuscula, calyce longiora.

Stam. receptaculo inserta. Filamenta capillaria. Antherae obovatae, apice dehiscentes.

Styl. filiformis, persistens. Stigma trifidum.

Caps. calyce vestita, trilocularis; loculis polyspermis; placentis liberis in quovis loculo. Sem. ovalia, solida.

Schk. t. 118. *Gärtn.* t. 63. *Bot. mag.* 1057. *Humb.* nov. gen. 3. t. 264. [*Cuellaria* R. et P.]

1751. *CLIFTONIA* Banks. (1805.)

Cal. minimus, quinquelobus.

Pet. quinque spathulata, subunguiculata, concaviuscula.

Stam. Filamenta crassiuscula, corolla breviora, receptaculo inserta, alterna longiora. Antherae subrotundae.

Stigma sessile, subpeltatum, lobatum.

Caps. quadrilocularis, quadrialata. Sem. solitaria.

Bot. mag. 1625. *Gärtn.* t. 225. [*Mylocaryum* Willd. *Pursh.* fl. amer. sept. t. 14.]

1752. *PYROLA* Brunf. (1532.)

Cal. quinquefidus, vel quinquepartitus.

Pet. subrotunda vel ovalia, patentia vel conniventia.

Stam. hypogyna, conniventia vel ascendentia. Antherae apice bicornes, tubulis dehiscentes.

Styl. filiformis, erectus vel declinatus, staminibus longior.

Stigma quinquefidum vel quinquelobum.

Caps. pentagona, quinquelocularis, quinquevalvis, angulis dehiscentibus. Placenta fungosa in quovis loculo. Sem. saepius membranula laxa obvoluta.

Schk. t. 119. *Sturm.* fl. germ. *Gärtn.* t. 63.

753. *CHIMAPHILA* Pursh. (1814.)*Cal.* quinquesfidus.*Pet.* rotundata, patentia.*Stam.* hypogyna. Filamenta medio incrassata, discophora. Antherae bicornes.*Styl.* brevissimus. Stigma quinquelobum, tuberculatum.*Caps.* quinquelocularis, quinquevalvis, apice dehiscens.*Rad.* Pyrol. t. 5. f. 2. [*Pyrola umbellata* et *maculata* L. Bot. mag. 897.]754. *MONOTROPA* Gronov. (1743.)*Cal.* pentasepalus connivens, vel eius loco bractee persistentes.*Pet.* quinque unguiculata, basi carnosae, saccata.*Stam.* hypogyna, erecta, cum petalis alternantia. Filamenta filiformia. Antherae depressae, subreniformes, basi bigibbae, transverse dehiscences.*Styl.* cylindricus. Stigma peltatum depressum.*Caps.* quinqueangularis, quinquelocularis. Placenta centralis. Sem. numerosa, minuta, membrana tenerrima laxa obvoluta.*Schk.* t. 116. *Sturm.* fl. germ. *Gärtn.* t. 185. *Reichenb.* ic. 5. t. 481. 482. *Hook.* fl. exot. t. 85. [*Hypopitys Dillen.* Nutt. *Monotropis Schweinitz* apud *Elliott.*]755. *GOMPHIA* Schreb. (1789.)*Cal.* quinquepartitus.*Pet.* cum laciniis calycis alternantia, latiora.*Stam.* Filamenta brevissima aut nulla. Antherae pyramidatae, apice biporosae.*Styl.* quingesulcatus. Stigma acutum.*Fructus* subdrupacei, monospermi, immersi receptaculo post anthesin turgido carnosio.*Ochneae, Cand.* in ann. mus. 17. t. 17—29. *Hilar.* in mém. mus. 10. t. 17. [*Jabotapita Plum.* ic. 153. *Correia Velloz,* in *Röm.* script. t. 6. f. 11.]756. *QUASSIA* L. (1763.)*Cal.* quinquepartitus, quinquepartitus.*Pet.* quinque.*Stam.* perigyna. Filamenta e squamis ciliatis orta. Antherae ovales, basi emarginatae.*Styl.* sulcatus. Stigma capitatum, quinquelobum.*Fructus* quinque drupacei, bivalves, monospermi.*Simarubeae.* *Adr. Juss.* in mém. mus. 12. t. 27. [*Simaruba Aubl.* guian. t. 331. 332. *Gärtn.* t. 70. *Adr. Juss.* l. c. *Aruba Aubl.* guian. t. 115. *Simaba ib.* t. 153. *Hilar.* in mém. mus. 10. t. 18. plant. mémor. brasil. 1. t. 17. 18. *Humb.* nov. gen. 6. t. 514. a. b. *Adr. Juss.* mém. mus. 12. t. 27.]

R U T A C E A E.

1757. RUTA Colum. Plin.

Cal. quadri-quinquepartitus.

Pet. quatuor, quinque, unguiculata.

Stam. disco hypogyno inserta. Filamenta filiformia. Antherae ovales.

Styli quatuor, quinque superne in unum confluentes.

Stigma quadri-quinquelobum.

Discus clavatus germina quatuor aut quinque ferens, poris lateralibus nectariferis instructus.

Caps. quatuor, quinque, plerumque in unicam coacta, apice et introrsum dehiscentes. Sem. semilunata, Embryo in centro albuminis.

Schk. t. 115. *Gärtn.* t. 111. *Jacqu.* ic. rar. 1. t. 78. *Bot. mag.* 2018. 2254. *Bot. reg.* 367. [*Aplophyllum* *Adr. Juss.*]

1758. DICTAMNUS Brunf. (1532.)

Cal. quinquepartitus, deciduus.

Pet. quinque lanceolata, basi attenuata, inferiora declinata.

Stam. hypogyna, declinata. Filamenta longitudine inaequalia, filiformia, superne glandulosa. Antherae subrotundae.

Styl. sulcatus. Stigma obtusum.

Folliculi quinque coriacei, dispermi. Sem. nitida. Cocci elastici, bivalves.

Schk. t. 114. *Adr. Juss.* in mém. mus. 12. t. 18. [*Fraxinella* *Gärtn.* t. 69.]

1759. ERIOSTEMON Sm. (1798.)

Cal. quinquepartitus, persistens.

Pet. marcescentia.

Stam. hypogyna, subaequalia, erecta. Filamenta plapiuscula, pilosa. Antherae biloculares, apice appendilatae.

Styli quinque in unum coacti. Stigma quinquelobum.

Discus elevatus germina quinque ferens.

Capsulae quinque, mono-dispermae. Sem. ovalia strophiolata.

Adr. Juss. in mém. mus. 12. t. 21. n. 25.

1760. PHEBALIUM Vent. (1803.)

Cal. brevis, persistens, quinquedentatus aut subinteger.

Pet. patula.

Stam. hypogyna, alterna breviora. Filamenta subulata, glabra. Antherae subrotundae, quadriloculares, callo minuta glanduloso coronatae.

Styli quinque in unum coacti. Stigma quinquelobum.

Discus elevatus, germina quinque ferens.

Cocci duo aut tres aut quinque monospermi.

Vent. Malmaia. t. 102. *Adr. Juss.* in act. soc. hist. nat. par. 2. t. 10 — 12. [*Eriostemon* *Labill.* nov. holl. t. 141.]

1761. *Crowea* Sm. (1798.)*Cal.* quinquepartitus.*Pet.* quinque, ovalia, patentia.*Stam.* sub disco germinifero inserta. Filamenta planiuscula, pilis intertextis nexa, apice in appendiculam villosam abeuntia. Antherae biloculares, lateri interiori filamentorum adnatae.*Styl.* subnullus. Stigma carnosum, sublobatum.*Discus* carnosus, germina quinque ferens.*Caps.* quinque monospermae.*Andr.* repos. t. 79. *Adr. Juss.* in mém. mus. 12. t. 21. n. 24.1762. *Choisia* Kunth. (1823.)*Cal.* pentasepalus, deciduus.*Pet.* quinque ovalia, brevissima, unguiculata.*Stam.* hypogyna. Filamenta subulata, alterna breviora. Antherae basi affixae, cordatae, biloculares, rima longitudinali dehiscentes.*Styl.* brevissimus, sulcatus. Stigma capitatum; quinquelobum.*Discus* nullus.*Caps.* quinquelocularis, quinquerostrata.*Humb.* nov. gen. 6. t. 513. [*Juliania Lexarz.*]1763. *Zygophyllum* L. (1737.)*Cal.* pentasepalus.*Pet.* unguiculata.*Stam.* hypogyna. Filamenta subulata, basi squama fornicata aucta. Antherae oblongae, incumbentes.*Styl.* simplex. Stigma minutum.*Discus* hypogynus.*Caps.* quinquelocularis, quinquevalvis, prismatica, loculis pleiospermis.*Bot. mag.* 372. *Delil.* flor. aeg. t. 27. f. 3. *Forsk.* ic. t. 11. f. 12. *Gärtn.* t. 112. [*Röperia Adr. Juss.* solo numero partium desciscit.]1764. *Larrea* Cav. (1800.)*Cal.* pentasepalus.*Pet.* unguiculata.*Stam.* hypogyna. Filamenta subulata, basi squama fornicata aucta. Antherae cordatae.*Styl.* pentagonus, acutus.*Fructus* pentacoccus.*Anal. de cienc. natur.* 2. t. 18. 19. *Cav.* ic. 6. t. 559. 560. *Mém. du mus.* 12. t. 15. n. 5.1765. *Guaicum* Monard. Clus. (1601.)*Cal.* pentasepalus.*Pet.* unguiculata.*Stam.* hypogyna. Filamenta subulata, basi squamata. Antherae oblongae.*Styl.* brevis. Stigma acutum.

Caps. duae aut quinque, compressae, coriaceae, intus coalitae, monospermae.

Clus. exot. p. 314. *Plum.* gen. t. 17. *Gärtn.* t. 113. *Adr.* *Juss.* in mém. mus. 12. t. 16. n. 7.

1766. **FAGONIA** Tournef. (1700.)

Cal. pentasepalus, deciduus.

Pet. unguiculata.

Stam. hypogyna. Filamenta subulata, basi esquamata.

Antherae subrotundae.

Styl. angulatus, persistens. Stigma acutum.

Fructus pentacoccus. Cocci bivalves, monospermi.

Tourn. instit. t. 141. *Gärtn.* t. 113. *Delil.* fl. aeg. t. 27. 28.

1767. **TRIBULUS** Theophr. Plin.

Cal. pentasepalus.

Pet. patentia.

Stam. alterna petalorum basi inserta, alterna hypogyna; haec singula extus glandula stipata. Filamenta filiformia.

Antherae ovales, rimis duabus longitudinalibus dehiscentia.

Styl. crassus, conicus. Stigma quinquecostatum.

Cocci quinque, initio connati, extus (plerumque) tuberculati aut spinosi, intus transverse divisi in loculos duos aut quatuor, monospermos.

Schk. t. 115. *Jacqu.* ic. rar. 3. t. 462. *Adr. Juss.* in mém. mus. 12. t. 14. n. 1. [*Ehrenbergia* Mart. nov. gen. 2. t. 163.]

LEGUMINOSAE.

1768. **CASTANOSPERMUM** Cunningh. Hook. (1830.)

Cal. coloratus, tubulosus, limbo quinqueidentato, subbilabiato.

Pet. quinque, conniventia, subaequalia, unguiculata, subpapilionacea.

Stam. libera, exserta. Antherae dorso affixae, biloculares.

Styl. stigmatate simplici.

Legumen stipitatum, grande, oblongum, bivalve, tetraspermum; valvis coriaceis intus spongioso-pulposis.

Hook. bot. miscell. 1. t. 51. 52.

1769. **GYMNOCLADUS** Lam. (1783.)

Flores dioeci.

Cal. tubulosus, limbo quinquefido.

Pet. oblonga, calyci inserta, cum eius laciniis alternantia.

Stam. inclusa. Filamenta tubo calycis adnata, alterna breviora. Antherae ovales, biloculares.

Styl. simplex. Stigma capitatum.

Legumen magnum, coriaceum, pleiospermum, intus pulposum.

Michaux fl. bor. amer. t. 51. *Reichenb.* Mag. t. 40.

1770. **ENTADA** Rheed. Adans. (1763.)

Flores polygami.

Cal. campanulatus, quinquedentatus.

Pet. minuscula, oblonga.

Stam. decem aut plura, elongata. Antherae subrotundae.

Legumen maximum, compressum, articulatum; valvis demum in membranas duas separatis; costis persistentibus.

Cand. legum. t. 61. 62. [*Mimosa Plum.* ic. t. 12. *Gigalobium P. Brown* jam.]

1. **DESMANTHUS** Willd. (1800.)

Cal. quinquedentatus.

Pet. quinque distincta aut coalita, spatulata. Flores inferiores apetalii neutri, filamentis abortientibus dilatato-membranaceis.

Stam. decem, rarius quinque.

Legumen continuum, pleiospermum.

Kunth. legum. t. 35. *Cand. legum.* t. 67. [*Mimosa Roxb. corom.* 2. t. 119. 174. *Humb. plant. aequin.* 1. t. 16. *Neptunia Lour.*]

2. **GAGNEBINIA** Neck. (1790.)

Cal. quinquedentatus.

Pet. quinque, oblongo-linearia, distincta.

Stam. decem distincta. Antherae oblongae, emarginatae.

Styl. elongatus, filiformis, deciduus.

Legumen complanatum, exsuccum, non dehiscens; margine ex utraque sutura alatum, intus transverse multiloculare, loculis monospermis.

Cand. legum. t. 64.

3. **PROSOPIS** L. (1767.)

Flores polygami.

Cal. quinquedentatus.

Pet. quinque, libera.

Stam. decem, basi ima vix coalita. Antherae oblongae, biloculares.

Legumen continuum lineare, subtorulosum, pulpa farctum, transverse articulatum, valvis non dehiscens.

Roxb. corom. 1. t. 63. *Kunth. legum.* t. 33. 34. [*Lagonychium M. B.*]

4. **ADENANTHERA** L. (1737.)

Cal. quinquedentatus.

Pet. lanceolata.

Stam. Filamenta erecta, subulata. Antherae subrotundae, incumbentes, glandula substipitata coronatae.

Styl. subulatus. Stigma simplex.

Legumen elongatum, membranaceum, transverse subarticulatum, seminibus remotis.

Lam. ill. t. 334. [*Corallaria et Clypearia Rumph. amboin.* 3. t. 109. 111.]

5. **MORINGA** Acost. Burm. (1737.)

Cal. subpentasepalus, deciduus.

Pet. calyci inserta, oblonga, subaequalia, superascendens.

Stam. inaequalia, discreta, interdum quinque abortient

Styl. filiformis, acutus, elongatus.

Legumen trivalve, elongatum, polyspermum. Sem. prurumque alata.

Burm. zeyl. t. 75. Lam. ill. t. 337. Gärtn. t. 147. [Ano Lour. Hyperanthera Vahl.]

1776. *GUILANDINA* L. (1737.)

Cal. pentasepalus, urceolatus.

Pet. calyci inserta, quinque, sessilia, subaequalia.

Filamenta basi villosa, decumbentia, inaequalia. *Antherae* oblongae, dorso affixae.

Styl. brevis, filiformis.

Legumen ovale, compressum, subventricosum, oligospermum. Sem. ossea.

Lam. ill. t. 336. Gärtn. t. 149.

1777. *SWEETIA* * (1825.)

Cal. campanulatus, quadri-quinquefidus.

Pet. quinque obovata, patentissima, calyci inserta.

Stam. imo calyci inserta. *Antherae* globosae.

Legumen lineare, polyspermum.

[*Acosmium Schott.*]

1778. *EXOSTYLES* Schott. (1827.)

Cal. campanulatus, laciniis tribus tandem revolutis.

Pet. quinque subaequalia, imbricato-conniventia, caly medio inserta.

Stam. subaequalia. *Filamenta* basi dilatata. *Antherae* sagittatae, cuspidatae.

Styl. elongatus, cylindricus. Stigma simplex.

Legumen crassum, polyspermum.

1779. *MELANOXYLON* Schott. (1827.)

Cal. pentasepalus, sepalis basi concretis.

Pet. quinque subaequalia.

Stam. *Filamenta* hirsuta. *Antherae* longitudinaliter dehiscentes.

Legumen compressum, falcatum. Sem. arillo in alio producto, transverse leguminum cavitates implente.

1780. *BARYXYLON* Lour. (1790.)

Cal. pentasepalus, deciduus, sepalis reflexis.

Pet. hypogyna, subrotunda, unguiculata, plicata, patentia

Stam. hypogyna. *Antherae* quadiloculares.

Styl. filiformis. Stigma concavum.

Legumen crassum, teretiusculum, polyspermum. Sem. subrotundo-angulata.

1781. *CADIA* Forsk. (1775.)

Cal. campanulatus, quinquepartitus, basi intus glandulosus.

- Pet.* quinque, calyci inserta.
Stam. Filamenta basi gibboso - geniculata.
Stigma sessile acutum.
Legumen breviter stipitatum, lineare, polyspermum.
 [Panciatica *Piccivoli* hort. Pauciat. 1783.]

782. ZUCCAGNIA Cav. (1799.)

- Cal.* quinquepartitus, coloratus; laciniis oblongis obtusis
Pet. quinque, subaequalia, calyci inserta.
Stam. ascendentiā. Filamenta subulata, inferne pilosa.
 Antherae ovatae, biloculares.
Styl. filiformis, incurvus. Stigma infundibuliforme.
Legumen ovatum, pilosissimum, monospermum. Sem.
 apici suturae affixum.

Cav. ic. 5. t. 403.

783. HAEMATOKYLON L. (1737.)

- Cal.* urceolatus, coloratus, limbo quinquelobo.
Pet. quinque, aequalia, calyci inserta.
Stam. calyci inserta. Filamenta inferne pilosa. Antherae
 ovaes.
Styl. filiformis, incurvus. Stigma infundibuliforme.
Legumen lanceolatum, utrinque acuminatum, dispermium;
 valvis medio ruptis, suturis non dehiscentibus.

Lam. ill. t. 340.

784. MELANOSTICTA Cand. (1826.)

- Cal.* quinquepartitus, glandulosus; limbo deciduo.
Pet. aequalia, ovalia.
Stam. aequalia. Filamenta subulata, inferne dilatata,
 sparse plumosa. Antherae ovaes, biloculares.
Styl. filiformis, acutus.
Legumen oblongum, compressum, subtetraspermum.

Cand. legum. t. 69.

785. OMPHALOBIUM Gärtn. (1788.)

- Cal.* quinquepartitus (quandoque quadrifidus).
Pet. unguiculata, hypogyna, erecto - patentia.
Stam. aequalia. Filamenta glabra, subulata, basi sub-
 connata.
Styl. simplex, persistens. Germen pedicellatum.
Caps. leguminosa, monosperma. Sem. arillo magno di-
 midiato.

Gärtn. t. 46. *Jacqu.* fil. ecl. t. 126. *Cand.* in act. soc. hist.
 nat. paris. 2. t. 16. [Connarus *Cav.* diss. 7. t. 221 - 223.]

786. CYNOMETRA L. (1741.)

- Cal.* quadripartitus, limbo reflexo, deciduo.
Pet. oblonga, aequalia.
Stam. decem, vel plura. Antherae apice bifidae.
Fructus semilunaris, carnosus, non dehiscentis, monosper-
 mus.

Lam. ill. t. 331. *Roxb.* corom. 3. t. 286.

MALPIGHIACEAE.

1787. *BUNCHOSIA* Juss. (1811.)

Cal. quinquepartitus, extus ad basin glandulis octo aut decem instructus.

Pet. quinque orbiculata, unguiculata, patentissima.

Stam. basi nexa.

Styl. simplex, stigmatibus peltato; aut apice bi-trifidus.

Bacca bilocularis, disperma.

[*Malpighia Cav. diss. 8. t. 238. 239.*

1788. *HIPTAGE* Gärtn. (1788. 1791.)

Cal. quinquepartitus, basi quinque glandulis stipatus.

Pet. obovata, fimbriata, unguiculata.

Stam. basi annulo cohaerentia, unico longiori. Filamenta subulata. Antherae ovoides.

Styl. simplex, acutus.

Fructus subglobosus, quadrialatus, alis inaequalibus.

Gärtn. t. 116. [*Banisteria tetraphylla* s. *Madagascariensis* Sonnerat. it. ind. 2. t. 135. Molina Cav. diss. 9. t. 263. Gärtner Roxb. corom. 1. t. 18.]

1789. *VARGASIA* Bert. * (1825.)

Cal. corollinus, pentasepalus, deciduus, eglandulosus.

Pet. spatulata, sepalis subaequalia.

Stam. hypogyna. Filamenta basi libera. Antherae ovoides.

Styl. conicus. Stigmata tria revoluta.

Samarae binae aut tres apice alatae.

1790. *ZYMUM* Noron. Juss. (1811.)

Cal. quinquefidus.

Pet. quinque, unguiculata, inflexa.

Stam. alterna breviora.

Germen poris tribus glandulosis instructum.

Styl. incurvus.

Fructus tres monospermi, appendicibus sex planis radiatis coronati, tribus aliis centralibus.

[*Tristellateia Thuars.*]

1791. *DIONAEA* Ellis. (1770.)

Cal. pentasepalus, persistens.

Pet. quinque, obtusa, concava.

Stam. decem aut viginti. Filamenta subulata. Antherae subrotundae, lateraliter dehiscentes.

Styl. brevissimus, filiformis. Stigma peltatum.

Caps. sublocularis, quinquevalvis, polysperma.

Droseraceae. Ellis in nov. act. ups. 1. t. 8., in monographia anglica 1770, latina et germanica, curante Schreber 1771. Vent. Malmais. t. 29.

792. *HYDROPITYON* Gärtn. fil. (1805.)*Cal.* pentasepalus, sepalis rotundis concavis.*Pet.* quinque subrotunda, concava.*Stam.* disco perigyno carnosio villosoque inserta. Filamenta brevia. Antherae cordatae, biloculares.*Styl.* brevis conicus. Stigma subpeltatum.*Caps.* monosperma.*Caryophylleae?* Gärtn. t. 183. [*Hottonia* Burm. zeyl. t. 55.]793. *LEPTOCHLAENA* Thuars. (1806.)*Involucrum* carnosum, urceolatum, uniflorum.*Cal.* trisepalus, villosus.*Pet.* quinque unguiculata, inferne in tubum conniventia, laminis patentibus.*Stam.* urceolo inserta, exstantia. Filamenta capillaria.

Antherae dorso adnatae, biloculares.

Styl. crassus. Stigma trilobum.*Caps.* monosperma.*Chlaenaceae.* Thuars. hist. afr. austr. t. 11.794. *HEISTERIA* Jacqu. (1763.)*Cal.* campanulatus, quinquefidus, post anthesin ampliatus.*Pet.* quinque, concava, patentia.*Stam.* alterna breviora. Filamenta plana. Antherae subrotundae.*Styl.* brevis. Stigma trifidum.*Drupa* monosperma; calyce maximo cincta.*Santaleae.* Jacqu. amer. t. 81.795. *BALANITES* Delil. (1802.)*Cal.* quinquepartitus.*Pet.* linearia, patentissima.*Stam.* Filamenta subulata, sub disco perigyno orta. Antherae subrotundae.*Styl.* brevis. Stigma truncatum.*Drupa* monosperma, nucleo pentagono.— — Delil. flor. aeg. t. 28. [*Ximenia aegyptiaca* L.]796. *RHAFTOSTYLUM* Bonpl. (1815.)*Cal.* quinquefidus.*Pet.* ovata, hypogyna.*Stam.* hypogyna. Filamenta basi dilatata, secum et cum petalis cohaerentia. Antherae ellipticae, biloculares.*Stigma* sessile, trilobum.*Germen* triloculare.*Humb.* plant. aequin. 2. t. 125.797. *CHILMORIA* Hamilt. (1822.)

Flores polygami.

Cal. pentasepalus.*Pet.* quinque, cum squamis totidem alternantia.*Stam.* incerti numeri (5 - 15.)

Stigma peltatum sessile.

Bacca lignosa, unilocularis, polysperma.

1798. *CERATOPETALUM* Sm. (1793.)

Cal. quinquepartitus, coloratus, persistens; lacinii lanceolatis, acutis.

Pet. calyci inserta, inaequaliter pinnatifida.

Stam. calyci inserta. Antherae subrotundae, biloculares, basi calcaratae.

Styl. brevis, acutus.

Caps. bilocularis.

Cunoniaceae R. Br. *Sm. nov. Holl.* t. 3.

1799. *VIVIANIA* Cav. (1804.)

Cal. pentasepalus, persistens.

Pet. quinque obovata, unguiculata.

Stam. hypogyna: alterna squamis inserta quinque hypogynis, alterna interstitiis squamarum.

Styl. simplex, stigmatibus tribus crassiusculis.

Caps. trilocularis, trivalvis. Sem. solitaria:

— — *Anal. de cienc. nat.* 7. 3. 49.

1800. *PHYSOPODIUM* Desv. (1826.)

Cal. bracteatus, turbinatus, quinquedentatus, intus pilosistrigosus. Pedicellus tumidus, articulatus.

Pet. ovali-oblonga.

Stam. alterna longiora. Antherae oblongae, biloculares

Stylus capillaris.

Fructus?

1801. *MACRAEA* Lindl. (1828.)

Cal. campanulatus, quinquedentatus.

Pet. quinque, arida, persistentia.

Stam. hypogyna. Antherae biloculares.

Styl. stigmatibus tribus.

Caps. trilocularis.

β. *Fructus inferi.*

1802. *PETALOMA* Sw. (1788.)

Cal. bracteatus, urceolatus, quinquedentatus.

Pet. summo calyci inserta, aestivatione contorta, oblonga acuta.

Stam. calyci inserta, subinaequalia. Filamenta subulata

Antherae obovatae, apice poro gemino dehiscentes.

Styl. filiformis. Stigma capitatum.

Bacca mono-tetrasperma.

Myrteae - Melastomeae. Swartz fl. ind. occ. t. 14

[*Mouriri Aubl. guian.* t. 180.]

803. *POIVREA* Commers. Thuars. (1806.)

Cal. infundibuliformis, quinquelobus, deciduus.

Pet. quinque, calyci inserta.

Stam. calyci inserta, exstantia. Filamenta filiformia.

Antherae ovaes, biloculares, longitudinaliter dehiscentes.

Styl. filiformis.

Fructus nucamentaceus, pentapterus aut quinquangularis, monospermus. Sem. pendulum. Cotyledones spiratim convolutae.

[*Cristaria Sonnerat.* it. ind. 2. t. 140. *Combretum Roxb. corom.* 1. t. 59. *Bot. reg.* 429. *Gonocarpus Gul. Hamilton* ind. occid. p. 39.]

804. *QUISQUALIS* Rumph. (1750.)

Cal. longissime tubulosus, filiformis; limbo quinquedentato.

Pet. calyci inserta, oblonga obtusa.

Stam. calycis fauci inserta, exstantia, alterna longiora.

Antherae ovatae, acutae, erectae.

Styl. filiformis, staminibus longior. Stigma capitatum.

Drupa sicca, quinquangularis, monosperma. Cotyledones carnosae, maximae.

Combreteae. *Rumph. amboin.* 5. t. 38. *N. L. Burm.* ind. t. 28. 35. *Bot. mag.* 2033. *Bot. reg.* 492. *Pal. Beauv.* fl. ovar. t. 34. [*Lumnitzera Willd.* in nov. act. scrutat. berol. 4. p. 186.]

805. *SCHOUSBOA* Willd. (1799.)

Cal. tubuloso-campanulatus; limbo quinquedentato, deciduo.

Pet. ovato-lanceolata, cum dentibus calycis alternantia.

Stam. imo calyci inserta, longe exstantia, aequalia. Antherae ovaes, erectae.

Styl. filiformis, acutus.

Drupa quinquangularis, mono-disperma. Cotyledones crassae, carnosae.

Combreteae. [*Cacoucia Aubl. guian.* t. 179. *Combretum Venten. sel.* t. 58.]

806. *LAGUNCULARIA* Gärtn. fil. (1805.)

Cal. tubuloso-campanulatus, persistens; limbo quinquelobo.

Pet. calyci inserta, minima, caduca.

Stam. duplici serie calyci inserta, alterna longiora. Filamenta subulata. Antherae ovaes, erectae.

Styl. subulatus. Stigma capitatum.

Nux coriacea, obovato-oblonga, sulcata, calyce coronata, monosperma. Cotyledones foliaceae, spiratim convolutae.

Combreteae. *Gärtn.* t. 217. [*Gonocarpus Jacq. minor.* t. 53.]

807. *GUIERA* Juss. (1789.)

Flores capitati, involucrati.

Cal. tubulosus, gracilis; limbo campanulato quinquedentato.

Pet. oblongo-linearis, parva.

Stam. longe exserta. Antherae globosae.

Styl. filiformis.

Cap. prismatica, unilocularis, subpentasperma.

Combretaceae. Lam. ill. t. 360.

1808. CERATOSTACHYS Blum. (1826.)

Flores dense capitati, processibus filiformibus spongiosis mixti.

Calycis limbus integer.

Pet. calyci inserta, octo vel decem, parva.

Stam. numero varia, inaequalia.

Styl. brevis. Stigma bifidum.

Drupa baccata, monosperma, calycis limbo coronata.

Combretaceae?

1809. CHAMAELAUCIUM Desf. (1819.)

Flos ante anthesin bracteis binis concavis deciduis obvolutus.

Cal. quinquefidus; laciniis vel integris vel palmato-partitis ciliatis.

Pet. subrotunda, denticulata.

Stam. calycis ori inserta, ligulis alternis interpositis.

Filamenta brevia, vel aequalia, vel alterna longiuscula.

Antherae subrotundo-reniformes.

Styl. filiformis, apice incurvus, pilosus. Stigma capitatum.

Fructus monospermus. Sem. globosum.

Myrteae. Desf. in mém. mus. 5. t. 3. B. t. 4. 19. [Verticordia Cand.]

b. Corolla irregularis.

1810. RHODORA L. (1764.)

Cal. quinquedentatus, persistens.

Cor. calyci adnata, ringens, subdipetala; petalo superiore profunde bi-tripartito, inferiore latiore bilobo.

Stam. calyci inserta, declinata. Filamenta subulata, inaequalia. Antherae didymae, poro gemino dehiscentia.

Styl. filiformis. Stigma capitatum.

Cap. quinquelocularis, quinquevalvis, loculis polyspermis.

Placenta centralis, quinquangularis. Sem. angusta, membrana tenera obvoluta.

Ericaceae. Bot. mag. 474. Herit. stirp. 1. t. 68. Gärtn. t. 209.

LEGUMINOSAE.

1811. LACARA * (1822.)

Cal. campanulatus, quinquedentatus, persistens.

Pet. quinque, imo calyci inserta, unguiculata; summum immumque latiora, concava.

Stam. fundo calycis inserta, exstantia. Antherae transverse oblongae, versatiles, biloculares, longitudinaliter dehiscentes.

Styl. cylindricus obtusus, persistens.

Legumen compressum, oligospermum, pilosum.

[*Caulotretus Schott.* Cur. post. p. 406.]

1812. *HYMENAEA* L. (1737.)

Cal. turbinatus, bracteatus, subcoriaceus, limbo quadri-quinquepartito, deciduo.

Pet. oblonga, sessilia, inaequalia.

Filamenta subulata, glabra. Antherae dorso affixae, biloculares.

Stylus simplex. Stigma capitatum.

Legumen lignosum, pulpa exsucca farctum, monospermum.

Lam. ill. t. 330. *Gärtn.* t. 145. *Humb.* nov. gen. 6. t. 566. 567.

1813. *BAUHINIA* Plum. (1703.)

Cal. campanulato-infundibuliformis, laciniis longissimis, liberis, vel ligulae forma cohaerentibus; saepe latere fissis, aut basi hians; nonnunquam subbilabiatus, labio superiori bi-inferiori tridentato.

Pet. calyci inserta, unguiculata, suprema maiora, a ceteris distantia.

Stam. calyci inserta, declinata, vel omnia libera, fertilia; vel basi connata, abortientibus quinque, septem aut novem, unico longissimo fertili superstiti. Antherae elongatae, subincumbentes.

Styl. longissime exsertus. Stigma capitatum.

Germen stipitatum.

Legumen elongatum, polyspermum.

a. *Caulotretus Rich.* (*Bauhinia Kunth.* legum. t. 46.)
Stamina omnia fertilia. *Cal.* bilabiatus. *Germen* sessile.
Burm. zeyl. t. 18 *.

β. *Symphyopoda Cand.* *Stam.* tria fertilia. *Cal.* tubulosus, irregulariter dentatus. *Cand.* legum. t. 70. *Vahl.* symb. 3. t. 62. *Roxb.* corom. 3. t. 285. [*Phanera Lour.*]

γ. *Pauletia Cav.* *Stam.* omnia fertilia, vel alterna abortientia. *Cal.* lacinae longissimae. *Plum.* ic. t. 44. f. 4. *Cav.* ic. 5. t. 410.

δ. *Casperea Kunth.* *Stam.* unicum perfectum, reliquis abortientibus, basi coalitis. *Linn.* hort. Clifford. t. 15. *Cav.* ic. t. 404—407.

1814. *CASSIA* Jo. Actuar. Dodon. (1557.)

Cal. pentasepalus, obtusus vel acutus.

Pet. quinque, inferiora maiora.

Stam. basi calycis inserta, inaequalia: inferiora declinata, longiora, antheris maioribus; summa saepe abortientia. Antherae apice vel poris vel rimis binis dehiscentes.

Styl. filiformis, arcuatus.

Legumen membranaceum, coriaceum (rarissime lignosum); compressum, vel teretiusculum; quandoque membranis transversis septatum; ubi hoc contingit, pulpa etiam farctum est. Sem. compressa, in pulposis leguminibus septis incumbunt, in aliis valvarum parietibus parallela, seu verticalia.

1. *Fistula* Cand. (*Bactrylobium Willd.*) Cal. obtusissimus. Antherae birimosae. Legumen lignosum, non dehiscent, septatum, pulpa farctum. Sem. horizontalia, aeu septis parallela. *Tourn. instit. t. 392. Gärtn. t. 147. Colladon. Cass. t. 1.*

2. *Chamaefistula* Cand. Cal. obtusissimus. Antherae biporosae. Legumen membranaceum, teres, septatum; loculis pulpa parca obductis. Sem. horizontalia. *Jacqu. ic. rar. 1. t. 73. Kunth. legum. t. 38. 39. 40. Senna Tourn. inst. t. 390. Gärtn. t. 146. Jacqu. ecl. t. 87.*

3. *Herpetica* Cand. Cal. obtusiusculus. Antherae biporosae. Legumen membranaceum, dorso alatum. septatum, loculis vix pulpa obductis. Sem. horizontalia. *Jacqu. obs. 2. t. 45. f. 2.*

4. *Chamaesenna* Cand. Cal. obtusus. Antherae biporosae. Legumen compressum, valvis dehiscent, imperfecte septatum, loculis epulposis. Sem. valvis parallela. *Plum. ic. t. 76. Jacqu. ic. rar. 1. t. 71. 170. vol. 3. t. 459. Colla hort. ripul. t. 24. Colladon Cass. t. 12.*

5. *Absus* Cand. Cal. acutiusculus. Stamina subaequalia, omnia perfecta. Antherae apice birimosae. Legumen plano-compressum, eseptatum. Sem. verticalia, seu valvis parallela. *Jacqu. ecl. 1. t. 53.*

6. *Chamaecrista* Breyn. (*Grimaldia Schrank.*) Cal. acuminatus, sepalis stipulisque nervosis. Pedunculi bracteati. Antherae biporosae. Legumen compressum eseptatum. Sem. verticalia. *Colladon Cass. t. 9. 16—20. Linn. hort. Cliff. t. 36.*

1815. CAESALPINIA Plum. (1703.)

Cal. turbinatus; limbo quinquepartito, deciduo; lacinis inferiore maiore, concava.

Cor. subpapilionacea, petalis unguiculatis, summo difformi.

Stam. omnia perfecta, ascendentia. Filamenta basi villosa. Antherae biloculares, dorso affixae, longitudinaliter dehiscentes.

Stylus filiformis, apice clavatus. Stigma truncatum, fimbriatum.

Legumen compressum, lignosum vel spongiosum. Sem. subcompressa.

Roxb. corom. 1. t. 16. Lam. ill. t. 335. Gärtn. t. 144. Kunth. legum. t. 45. [Poinciana Tournef. L. leguminibus subseptatis differt. Gärtn. t. 150.]

1816. COULTERIA Kunth. (1823.)

Cal. turbinatus, limbo quinquepartito, obtuso, deciduo; lobo inferiori maiori pectinato-fimbriato.

Cor. subpapilionacea, petalis unguiculatis, summo difformi.
Stam. ascendentia, in lobo inferiori calycis latentia. Filamenta basi pilosa. Antherae biloculares, dorso affixae, longitudinaliter dehiscentes.

Styl. ascendens, basi articulatus.

Legumen compressum, subspongiosum, septatum. Sem. albuminosa.

Humb. nov. gen. 6. t. 568. 569. [Tara *Molin.* *Adenocalyx Berter.*]

817. *BOWDICHIA* Kunth. (1823.)

Cal. campanulatus, quinqueidentatus, dentibus duobus superioribus approximatis.

Cor. subpapilionacea; petala lateralia reliquis longiora.

Stigma? suburceolatum.

Legumen stipitatum, compressum, membranaceum, ad suturam marginato-alatum, non dehiscens. Sem. compressa.

818. *REICHARDIA* Roth. (1821.)

Cal. campanulatus, margine crenulato.

Cor. subpapilionacea, hexa-decapetala; petalis margini calycis inserta.

Stam. ascendentia, inferne villis nexa. Antherae biloculares, dorso affixae.

Stylus filiformis. *Stigma* barbatum.

Legumen coriaceum, monospermum, samaroideum in alam oblongam terminatum.

819. *MEZONEVRON* Desf. (1818.)

Cal. quinquepartitus; lobo inferiori fornicato ceteros in aestivatione involvente.

Pet. quinque unguiculata, summum minus.

Stam. declinata, inferae villosa.

Styl. incurvus.

Legumen foliaceum, planum, non dehiscens, polyspermum; sutura seminifera superne in alam expansa; hinc legumen quasi medio nervo instructum.

Desfont. in mém. mus. 4. t. 10. 11.

820. *HOFFMANNSEGGIA* Cav. (1798.)

Cal. quinquepartitus, laciniis aequalibus acutiusculis.

Pet. unguiculata; superius difforme, concavum.

Stam. (quandoque octo) plerumque decem, filiformia, basi hirsuta aut glanduloso-pilosa. Antherae ovaes.

Styl. filiformis. *Stigma* capitatum.

Legumen compressum, oligo-vel polyspermum.

Cav. ic. 4. t. 392. 393. [*Pomaria Cav.* ic. 5, t. 402.]

821. *CUBAEA* Scop. Schreb. (1789.)

Cal. turbinatus, quinquepartitus; lobis concavis subrotundis, infimo declinato.

Pet. calycis ori inserta, unguiculata; duo inferiora declinata.

Stam. inferne villosa; tria superiora breviora, erecta; inferiora longiora, declinata.

Styl. filiformis, acutus.

Legumen stipitatum, coriaceum, monospermum.

[Tachigali Aubl. guian. t. 143.]

1822. *PARKINSONIA* Plum. (1703.)

Cal. quinquepartitus, coloratus; laciniis acutis, reflexis, deciduis.

Pet. unguiculata, patentissima; quintum reniforme, longe unguiculatum.

Stam. inferne villosa, declinata. Antherae oblongae incumbentes.

Styl. filiformis, ascendens. Stigma obtusum.

Legumen longissimum, teres, inter semina compressum, unde moniliforme.

Linn. hort. Cliff. t. 13.

† 1823. *AFZELIA* Sm. (1798.)

Cal. tubulosus, limbo quadrifido, deciduo.

Pet. quatuor unguiculata, superiori maiori.

Stam. duo abortientia.

Styl. subulatus.

Legumen lignosum, septatum. Sem. arillata.

[Paucovia Willd.]

† 1824. *ALOËXYLON* Lour. (1790.)

Cal. tetrasepalus, deciduus: sepalum inferius falcatum, elongatum.

Pet. inaequalia.

Styl. filiformis.

Legumen lignosum, falcatum, monospermum. Sem. oblongum incurvum, arillatum.

† 1825. *METROCYNIA* Thuars. (1810.)

Cal. campanulatus, laciniis elongatis, coloratis, reflexis.

Pet. quinque, erecta.

Filamenta hirta. Antherae subrotundae.

Styl. stamina aequans.

Legumen reniforme, verrucosum vel plicatum, monospermum.

1826. *MYROSPERMUM* Jacqu. (1763.)

Cal. campanulatus, quinquedentatus, persistens.

Pet. unguiculata; supremum subrotundum, reliqua angusta.

Stam. glabra, persistentia. Antherae tuberculo vel rostro terminatae, basi bifidae.

Stylus brevis, persistens, lateralis.

Legumen stipitatum, inflatum, membranaceum, apice oblique obtuseque alatum, non dehiscens, monospermum.

Humb. nov. gen. 6. t. 570. 571. [Myroxylon et Tolivifera L.]

COROLLA PAPILIONACEA.

827. ANAGYRIS Diosc.

Cal. bilabiatus, quinquedentatus.

Cor. Alae vexillo longiores, carina dipetala elongata breviores.

Stam. ascendentia.

Styl. simplex. Stigma villosum.

Legumen stipitatum, teretiusculum, polyspermum, subseptatum.

Tournef. inst. t. 415. *Lam.* ill. t. 328. [*Macrotropis Cand.*]

828. CERCIS Theophr.

Cal. urceolatus, inferne gibbus, obtuse quinquedentatus.

Cor. Alae insignes, divaricatae. Carina dipetala.

Stam. declinata; quatuor longiora, rectiuscula. Antherae incumbentes.

Styl. stigmatate obtuso.

Legumen oblongum, membranaceum, polyspermum; sutura seminifera subalata.

Schk. t. 112. *Duham.* arb. ed. nov. t. 7. *Gärtn.* t. 144.

829. ANTHONOTHA Pal. Beauv. (1805.)

Cal. bilobus, coriaceus.

Cor. Carina emarginata, longe unguiculata. Vexillum maximum.

Stam. inaequalia, tria longissima, septem calyce minora.

Styl. filiformis. Stigma subbifidum.

Legumen compresso-planum, vix dehiscens, polyspermum, valvis rugoso-reticulatis.

Pal. Beauv. fl. ovar. 1. t. 42.

830. SOPHORA L. (1737.)

Cal. subcampanulatus, quinquedentatus.

Cor. Carinae petala subconnata.

Legumen moniliforme, apterum, polyspermum.

Plum. ic. t. 101. *Pall.* astrag. t. 87. *S. alopecuroides* L. desciscit staminibus diadelphis.

831. PODALIRIA Lam. (1792.)

Cal. campanulatus, basi truncatus, subbilabiatus; labio superiori emarginato-bidentato, inferiori tripartito.

Cor. Vexillum maximum, obcordatum.

Stam. ascendentia, persistentia; tria superiora libera, septem inferiora basi connata.

Stylus simplex, glaber.

Legumen ventricosum polyspermum.

Venten. Cels. t. 99. Bot. reg. 869. [*Hypocalyptus Thunb.*]

832. BAPTISIA Vent. (1808.)

Cal. campanulatus, basi subattenuatus, quadri-quinquedentatus, persistens.

Pet. unguiculata, subaequalia. Vexillum lateribus reflexum.

Stam. libera, decidua.

Legumen ventricosum, subcoriaceum, pedicellatum, stylo coronatum, pleiospermum.

[*Podaliria Vent. Cels. t. 56. Bot. mag. 1099. 1177.*]

1833. *VIRGILIA* Lam. (1792.)

Cal. campanulatus, quinquedentatus.

Pet. subaequalia. Vexillum planum.

Stigma imberbe.

Legumen compressum, polyspermum.

Bot. mag. 1590. [Robinia Herit. surp. 1. t. 75.]

1834. *AMMODENDRON* Fisch. Cand. (1822.)

Cal. quadrifidus, persistens, laciniis aequalibus.

Cor. Vexillum alis brevius, complicatum. Alae carina obtusa multo latiores.

Stamina libera.

Stylus filiformis. *Stigma* parce transverse barbatum.

Legumen lanceolatum, membranaceum, planum, monospermum.

[*Sophora bifolia Pall. astrag. t. 91.*]

1835. *CYCLOPIA* Venten. (1808.)

Cal. campanulatus, basi truncatus; limbo inaequaliter quinquefido, dente imo longiori.

Cor. Vexillum maximum, rugis longitudinalibus. Alae transverse plicatae. Carina inflexo-mucronata.

Stam. decidua.

Stigma hinc barbatum.

Legumen pedicellatum, coriaceum, compressum, pleiospermum.

[*Ibbetsonia Bot. mag. 1259.*]

1836. *THERMIA* Nuttall. (1818.)

Cal. campanulatus, bilabiatus, labio utroque bifido.

Cor. Vexillum lateribus reflexum. Carina obtusa, alis multo amplior.

Stam. persistentia.

Legumen compressum, polyspermum.

[*Sophora Pall. astrag. t. 89. 90. Thermopsis R. Br.*

Baptisia nepalensis Hook. fl. exot. t. 131.]

1837. *EDWARDSIA* Salisb. (1805.)

Cal. ollaeformis, supra fissus, margine reliquo obsolete quinquedentatus.

Cor. Vexillum connivens. Carina obtusa, paullo longior alis.

Stam. disco annulari perigyno imposita.

Styl. filiformis, exsertus.

Legumen moniliforme, quadrialatum, polyspermum.

Bot. reg. 738. [Sophora Ait. Bot. mag. 167.]

1838. *ORMOSIA* Jacks. (1811.)

Cal. bilabiatus, labio superiori bilobo, inferiori tripartito.

Cor. Vexillum subrotundum, alas carinamque subaequans.

Stam. Filamenta basi dilatata.

Styl. incurvus. Stigmata duo approximata obtusa, altero unilaterali.

Legumen compressum, lignosum, subdispermum.

Jacks. in *Linn. transact.* 10. t. 25 — 27. [*Robinia coccinea*
Aubl. *Sophora monosperma Sw.*]

839. *BAPHIA* Afzel. (1825.)

Cal. calyptraeformis, longitudinaliter sub carina ruptus, sub vexillo erectus, demum deciduus.

Cor. Vexillum patens, subrotundum. Alae lineares, vexillum aequantes. Carina acuta.

Legumen falciforme.

Loddig. bot. cabin. 4. t. 367.

840. *CHORIZEMA* Labill. (1798.)

Cal. bilabiatus, lab. sup. bi-, inf. trifido.

Cor. Carina ventricosa, alis brevior.

Styl. brevis uncinatus.

Legumen ventricosum, polyspermum, subsessile.

Labill. it. 1. t. 21. *Bot. mag.* 1032. *Bot. reg.* 986.

841. *PODOLOBIUM* R. Br. (1811.)

Cal. turbinatus, subbilabiatus, quinquefidus.

Cor. Vexillum amplum, planum. Carina compressa, alarum longitudine.

Styl. rectus, persistens.

Legumen pedicellatum, oblongum, compressum, oligospermum.

Bot. mag. 1477.

842. *OXYLOBIUM* Andr. (1809.)

Cal. quinquepartitus, subbilabiatus, demum reflexus.

Cor. Vexillum planum. Carina compressa, alas aequans.

Styl. ascendens. Stigma simplex.

Legumen subsessile, ovatum, acutum, ventricosum, polyspermum.

Andr. repos. 492. *Bot. mag.* 1544. 2442. *Bot. reg.* 392.
[*Callistachys elliptica Vent.*]

843. *CALLISTACHYS* Vent. (1803.)

Cal. bilabiatus, labio sup. bifido, inf. tripartito.

Cor. Vexillum amplum, erectum, emarginatum. Carina dipetala cum alis demissa.

Stam. disco inserta.

Legumen pedicellatum, lignosum, apice dehiscens, iunius septatum, polyspermum.

Vent. *Malm.* 115. *Bot. reg.* 216. *Bot. mag.* 1925.

844. *GOMPHOLOBIUM* Sm. (1798.)

Cal. quinquepartitus, subaequalis; laciniis late lanceolatis.

Cor. Vexillum amplum, expansum, emarginatum. Carinae petala antice concreta.

Styl. incurvus, deciduus.

Legumen globulosum, obtusissimum, pleiospermum.

Reichenb. hort. 1. t. 76. 97. *Smith. exot. bot.* t. 5. 58. *Labill. nov. holl.* t. 133. 134. *Salisb. parad. lond.* t. 6. (G. grandiflorum *Sm.* carina dipetala.)

1845. *BURTONIA* R. Br. (1811.)

Cal. quinquepartitus.

Pet. longitudine subaequalia, decidua. Carinae petala dorso concreta.

Styl. subulatus, basi dilatata. Stigma obtusum.

Legumen subrotundum, subventricosum.

[*Gompholobium* scabrum et minus *Sm.*]

1846. *JACKSONIA* R. Br. (1811.)

Cal. quinquepartitus subaequalis.

Cor. petala decidua, patentia. Vexillum planum, emarginatum. Carina dipetala.

Styl. subulatus, persistens.

Legumen ovatum, monospermum, valvis intus pubescentibus.

[*Gompholobium* spinosum *Labill. nov. holl.* 1. t. 136.]

1847. *VIMINARIA* Sm. (1805.)

Cal. campanulatus, angulatus, aequaliter quinquefidus.

Pet. longitudine subaequalia.

Styl. capillaris.

Legumen coriaceum, evalve, intus farctum, monospermum.

Smith. exot. bot. t. 27. *Bot. mag.* 1190. [*Daviesia* denudata *Vent. sel.* t. 6.]

1848. *SPHAEROLOBUM* Sm. (1805.)

Cal. subbilabiatus, quinquefidus.

Cor. Vexillum reflexum. Carina dipetala.

Stam. duo superiora distantia.

Styl. hinc membrana auctus.

Legumen pedicellatum, sphaericum, submonospermum.

Labill. nov. holl. t. 138.

1849. *GASTROLOBUM* R. Br. (1811.)

Cal. subbilabiatus, quinquefidus.

Cor. Vexillum erecto-reflexum. Carinae petala antice concreta.

Styl. subulatus, ascendens.

Legumen ventricosum. Sem. strophiolata.

Bot. reg. 411.

1850. *EUCHILOS* R. Br. (1811.)

Cal. bracteatus, quinquepartitus, bilabiatus; lab. sup. lobi maximi.

Cor. Vexillum amplum, subreniforme, emarginatum. Alae carinam aequantes.

Styl. subulatus, ascendens.

Legumen compressum, pedicellatum. Semina strophiolata. Stropholi lobi postici integri.

Bot. reg. 403.

851. *PULTENAEA* Sm. (1793.)*Cal.* bracteatus, bilabiatus, lab. sup. bi-inf. trifido.*Cor.* Vexillum planum aut subconnivens. Carina prominula saturius colorata.*Styl.* subulatus, saepe persistens.*Legumen* compressum, dispermum. Sem. strophiolata.

Stropholi lobii postici incisi.

Smith. nov. holl. t. 12. *Andr.* repos. t. 98. *Bot. mag.* 475.1394. 1588. 2086. 2091. *Labill.* nov. holl. t. 130. 131.*Reichenb.* hort. 2. t. 192 — 196.852. *SCLEROTHAMNUS* R. Br. (1811.)*Cal.* bracteatus, bilabiatus; lab. sub. bi-inf. trifido.*Cor.* Carina alarum longitudine.*Stylus* ascendens.*Legumen* pedicellatum, dispermum, ventricosum.853. *DAVIESIA* Sm. (1798.)*Cal.* ebracteatus, campanulato-turbinatus; quinquedentatus.*Cor.* Vexillum obcordatum, erecto-reflexum. Carina alis brevior.*Styl.* strictus persistens.*Legumen* compressum, monospermum, ad suturam inferiorem dilatatum, trapezoideum, elastice dissiliens. Sem. strophiolata. Strophiolus postice integer.*Bot. reg.* 728. 1005. *Labill.* nov. holl. t. 137.854. *EUTAXIA* R. Br. (1811.)*Cal.* ebracteatus, subcampanulatus, bilabiatus; lab. super. latiori obtuso, emarginato, inf. tridentato.*Cor.* Vexillum obovatum, erectum. Alae distantes.*Styl.* uncinatus. Stigma capitatum.*Legumen* ventricosum, submonospermum. Sem. strophiolata.[*Dillwynia* *Labill.* nov. holl. t. 140.]855. *DILLWYNIA* Sm. (1805.)*Cal.* bilabiatus, quinquefidus.*Cor.* Vexillum latum, divaricato-bilobum.*Pet.* cum staminibus calyci medio inserta.*Styl.* uncinatus.*Legumen* ventricosum, dispermum. Sem. strophiolata.*Sm.* exot. bot. t. 25. 26. *Labill.* nov. holl. t. 139.856. *AOTUS* Sm. (1805.)*Cal.* ebracteatus, quinquefidus.*Pet.* subaequalia.*Stam.* decidua.*Stylus* erectus, filiformis, superne parumper incrassatus.*Legumen* dispermum. Strophiolus seminum nullus.*Bot. mag.* 949. *Labill.* nov. holl. t. 132.

1857. *BRACHYSEMA* R. Br. (1811.)*Cal.* ventricosus, basi truncatus, limbo quinquefido.*Cor.* Vexillum minimum. Alae carinam aequantes.*Stam.* basi incrassata.*Stylus* filiformis.*Germen* stipitatum, stipite basi vaginato.*Legumen* ventricosum, polyspermum.

Bot. reg. 118. 642.

1858. *MIRBELIA* Sm. (1805.)*Cal.* bilabiatus, quinquefidus.*Cor.* Vexillum dilatatum, erectum. Alae porrectae, truncatae, carina longiores.*Legumen* turgidum, biloculare, polyspermum, mucronatum. Sutura inflexa.

Vent. Malmais. t. 119. Reichenb. hort. 2. t. 191.

1859. *ADESMIA* Cand. (1825.)*Cal.* quinquefidus, laciniis acutis aequalibus.*Cor.* Vexillum initio connivens. Carina truncata.*Lomentum* compressum, sutura altera rectiuscula, altera sinuato-lobata.

Cand. legum. t. 48 — 50. [Hedysarum muricatum Jacq. ic. rar. 3. t. 568. Patagonium Schrank.]

II. *Flores imperfecti.*A. *Fructus superi.*

LEGUMINOSAE.

1860. *HARDWICKIA* Roxb. (1819.)*Cal.* corollinus pentasepalus.*Cor.* nulla.*Stam.* alterna longiora. Antherae ovatae apiculatae.*Styl.* ascendens. Stigma capitatum.*Legumen* membranaceum, lanceolatum, monospermum.

Roxb. corom. 3. t. 209.

1861. *COPAIFERA* L. (1749.)*Cal.* quadripartitus, patens.*Cor.* nulla.*Stam.* subaequalia. Antherae biloculares, dorso affixae.*Legumen* stipitatum, coriaceum, monospermum. Sem. arillo obvolutum.

Desf. in mém. mus. 7. t. 13. 14. Humb. nov. gen. 7. t. 659.

Hayne plant. off. 10. t. 16 — 23. [Weigeltia Reichenb.]

1862. *CRUDIA* Schreb. (1789.)*Cal.* bracteatus, turbinatus, quadripartitus, laciniis inaequalibus.*Cor.* nulla.*Stam.* calycis collo inserta. Antherae dorso adnatae.

Stylus capillaris, flexuosus. Stigma capitatum.

Legumen coriaceum, crassum, monospermum.

[*Apalatoa* *Aubl.* guian. t. 147. Ann. of bot. 1. t. 7. f. 5. *Tou-chiroa* *Aubl.* ib. t. 148. *Parivoa* ib. t. 304. *Vouarana* ib. t. 374. *Chalcas* *Schreb.*]

863. *DETARIUM* Juss. (1789.)

Cal. quadrifidus.

Cor. nulla.

Stam. alterna breviora.

Drupa crassa, farinacea, monosperma.

R U T A C E A E.

864. *DIPLOCHLAENA* R. Br. (1814.)

Involucrum multiflorum duplex, exterius quinquepartitum, interius decaphyllum.

Cal. quinquepaleaceus.

Stam. alterna longiora, persistentia.

Styli quinque in unum coaliti. Stigma quinquelobum.

Folliculi quinque monospermi.

Desfont. in mém. mus. 3. t. 19. 20.

T E R E B I N T H E A E.

865. *HETERODENDRON* Desfont. (1818.)

Cal. urceolatus, margine repando.

Cor. nulla.

Filamenta brevissima. *Antherae* tetragonae, biloculares.

Styl. nullus.

Germīna quatuor subglobosa.

Desfont. in mém. mus. 4. t. 3.

866. *STYLOBASIUM* Desfont. (1819.)

Flores polygami.

Cal. urceolatus, quinquelobus, coloratus.

Cor. nulla.

Stam. hypogyna. *Filamenta* capillaria, persistentia. *Antherae* crassae, oblongae, biloculares.

Styl. filiformis. Stigma capitatum.

Drupa monosperma.

Desfont. in mém. mus. 5. t. 2.

867. *RHYNCHOTHECA* Ruiz. et Pav. (1794.)

Cal. pentasepalus persistens.

Cor. nulla.

Stam. hypogyna, persistentia.

Styl. brevissimus. Stigmata quinque.

Fructus pentacoccus.

Geranieae. *Humb.* nov. gen. 5. t. 464. 465.

868. *GEISSOIS* Labill. (1825.)

Cal. tetrasepalus.

Cor. nulla.

Stam. hypogyna. Filamenta capillaria. Antherae ovatae.

Styl. bifidus.

Fructus linearis, siliquaeformis, bilocularis, bivalvis. Sem. alata.

— — *Labill.* nov. Caledon. t. 50.

1869. *DARWINIA* Rudg. (1813.)

Cal. corollinus tubulosus, limbo quinquefido.

Cor. nulla.

Stam. Filamenta brevissima duplici serie fauci calycis inserta. Antherae oblongae.

Styl. longissime exsertus.

Fructus ?

— — *Linn.* transact. 11. t. 22.

1870. *ADENOSTEMON* Pers. (1805.)

Cal. corollinus coriaceus duplex; externus quadripartitus, internus trisepalus.

Stam. decem aut plura, triplici serie, sensim minora. Glandulae binae ad basin singulorum staminum. Antherae oblongae laterales.

Styl. subulatus. Stigmata tria.

Drupa monosperma.

— — [*Gomortega R.* et *P. Keulia Molin.*]

1871. *SAMYDA* P. Brown. (1756.) Jacqu. (1777.)

Cal. campanulatus, coloratus, persistens, quinquefidus, laciniis inaequalibus.

Stam. tubo calycis adnata. Filamenta brevina, lanceolata, cuspidata, basi subconnata. Antherae oblongae, erectae, dorso supra basin affixae, biloculares.

Styl. simplex. Stigma capitatum.

Caps. coriacea, apice trifariam dehiscens, unilocularis.

Sem. numerosa, baccata, valvis affixa.

Samydeae. *Venten.* sel. t. 43. *Plum.* ic. t. 146. f. 3.

[*Guidonia Plum.* gen. t. 24.]

1872. *CASEARIA* Jacqu. (1763.)

Cal. quinquepartitus, coloratus, persistens.

Stam. perigyna, (numero varia,) basi subconnata, interiectis squamis totidem hirsutis. Antherae biloculares.

Styl. simplex. Stigma capitatum, aut trifidum.

Caps. coriacea trivalvis, valvis medio seminiferis. Sem. baccata.

Samydeae. *Jacqu.* act. helv. 8. p. 58. ic. *Vent.* sel. t. 44.

45. 46. *Humb.* nov. gen. 5. t. 479. 480. [*Samyda P.*

Brown. jam. in textu p. 217. *Guidonia* in ic. t. 23. f. 3.

Iroucana Aubl. guian. t. 127. *Pitumba ib.* t. 385. *An-*

vinga Lam. ill. t. 355. *Gärtn.* t. 224. *Melistaureum Ford.*

char. t. 72. *Athenaea Schreb.* *Chaetocrater R.* et *P. Cr-*

teria Pers.]

873. AQUILARIA Lam. (1785.)*Cal.* coriaceus, turbinatus, quinquelobus, persistens.*Urceolus* perigynus, quinquepartitus, laciniis bifidis.*Stam.* Filamenta intra urceolum, sub eius fissuris inserta, brevissima. Antherae lineares, dorso affixae, versatiles.*Styl.* nullus. Stigma simplex.*Caps.* pyriformis, bilocularis, bivalvis; valvis medio septigeris. Sem. solitaria, extus spongiosa.*Aquilarinae* R. Brown. *Cav. diss.* t. 224.**874. OPHISPERMUM Lour. (1790.)***Cal.* campanulatus, sexpartitus.*Urceolus* perigynus, decemlobus.*Stam.* receptaculo inserta, brevissima. Antherae oblongae, erectae.*Styl.* staminibus longior. Stigma bifidum.*Caps.* compressa, bivalvis, monosperma. Sem. ovatum, ala laterali elongata, subtereti, sinuosa.**875. AUGEA Thunb. (1794.)***Cal.* quinquepartitus, persistens; laciniis ovatis, erectis, concavis.*Urceolus* perigynus, brevis, decemdentatus.*Stam.* Filamenta brevissima, dentibus urceoli inserta. Antherae erectae, subulatae, sulcatae.*Styl.* filiformis, brevissimus. Stigma simplex.*Caps.* baccata, decemlocularis, decemvalvis. Sem. numerosa, membrana obvoluta.**876. DAIS Royen. (1740.)**

Flores capitati. Involucrum quadrifidum.

Cal. corollinus, basi ventricosus, tubulosus, persistens, limbo patente quinquefido.*Stam.* duplici serie tubo calycis inserta. Filamenta brevissima. Antherae ovoides, erectae.*Styl.* filiformis.*Bacca* monosperma.*Thymelaeae.* Bot. mag. 147. *N. L. Burm. fl. ind.* t. 32. f. 2. *Gärtn.* t. 39.**B. Fructus inferi.****877. BUCIDA L. (1759.)***Cal.* campanulato-urceolatus, persistens; limbo quinque-dentato.*Stam.* limbo imposita, longe exstantia. Antherae didymae.*Styl.* subulatus.

Drupa monosperma; putamine angulato. Cotyledones spiratim convolutae.

Combretaceae. *Vahl*, eclog. 1. t. 8. *Gärtn.* t. 217. [*Buccaras* *P. Brown*, jam. t. 23. f. 1. Nomen e fortuita metamorphosi, qua apex spicae florigeræ abijt in processum falcato-cornutum.]

1878. AGATHISANTHES Blum. (1826.)

Flores dioeci.

Cal. quinquefidus.

Stam. disco inserta. Filamenta brevissima. Antherae didymae.

Styl. brevis bifidus, discum planum perforans.

Drupa baccata, umbilicata, monosperma.

Combretaceae.

1879. TERMINALIA L. (1767.)

Flores polygami.

Cal. campanulato-urceolatus; limbo quinquefido, basi villosa.

Stam. calyci inserta, exstantia. Antherae dorso affixae, biloculares.

Styl. filiformis acutus, basi annulo quinquelobo cinctus.

Drupa monosperma; vel carnosae, putamine angulato; vel exsucca, alata. Cotyledones spiratim convolutae.

Combretaceae. *Jacqu.* ic. rar. 1. t. 197. *Lam.* ill. t. 848. *Roxb.* corom. 2. t. 97. 98. vol. 3. t. 224. *Mart.* nov. gen. t. 27. [*Tanibuca* *Aubl.* guian. t. 178. *Catappa* *Gärtn.* t. 127. *Badamia* et *Myrobalanus* *Gärtn.* t. 97. *Chuncoa* *Pav.* Jun. *Guibernatia* *Ruiz* et *Pav.* prodr. t. 36.]

Drupae et putaminis formae variae hae sunt:

1. *Drupa* carnosae, glaberrimae, putamine angulato. *T. Belirica* *Roxb.*, *Badamia* *Cand.*
2. *Drupa* glabra, putamine apice alato. *T. glabrata* *Ford.*
3. *Drupa* angulata, longitudinaliter sulcata. *T. Chebala* *Retz.*, *clitvina* *Roxb.*, *moluccana* *Lam.*, *procera* *Roxb.*, *Pamea* *Cand.*, *madagascarensis* *, *rhomboidea* *.
4. *Drupa* lacunoso-foraminulata. *T. latifolia* *Sw.*
5. *Drupa* rugoso-sulcata, putamine bialato. *T. Tanibuca* *Sw.*, *Catappa* *L.*, *Vernix* *Lam.*, *mauritanica* *Lam.*, *angustifolia* *Jacqu.*
6. *Drupa* exsucca, alata. *T. brasiliensis* *, *fagifolia* *Mart.*, *monoptera*, alata, cuneata et crenulata *Roth.* [*Pentaptera* *Roxb.* Ceterarum fructus non observati.]

1880. GETONIA Roxb. (1795.)

Cal. campanulatus, quinquepartitus; laciniis lanceolatis trinerviis.

Stam. calyci duplici serie inserta, laciniis calycis breviora.

Filamenta alterna latiora. Antherae globosae, didymae.

Styl. filiformis, pubescens.

Nux calyce coronata, pentagona, monosperma.

— — *Roxb.* corom. 1. t. 87. *Gärtn.* t. 217. [*Calycoptera* *Lam.* ill. t. 357.]

DIGYNIA.

I. Flores perfecti.

881. ROYENIA L. (1737.)

Cal. urceolatus, quinquefidus, persistens.

Cor. urceolata, quinquefida, laciniis ovatis revolutis.

Stam. corollae adnatae. Filamenta brevissima. Antherae erectae, didymae.

Styli stamina superantes.

Bacca quadrilocularis. Sem. quatuor, aut abortu duo, plano-convexa.

Styraceae. Lam. ill. t. 370. Jacqu. coll. suppl. t. 13.

f. 1. Gärtn. t. 94.

882. TURARIA Molin. (1782.)

Cal. tubulosus quinquedentatus.

Cor. infundibuliformis, quinquefida.

Stam. Filamenta brevissima, filiformia. Antherae didymae.

Styli setacei.

Caps. bilocularis, disperma.

Styraceae?

883. PACHYNEMA R. Br. Cand. (1818.)

Cal. pentasepalus, persistens, sepalis subrotundis, concavis.

Pet. quinque ovalia, cum sepalis alternantia.

Stam. perigyna. Filamenta inferne crassissima, apice attenuata. Antherae subrotundae, dorso adnatae. (Stam. aliquot abortientia.)

Germina duo in stylos filiformes desinentia.

Dilleniaceae. Lessert. ic. sel. t. 73.

884. ADRASTEA Cand. (1818.)

Cal. pentasepalus, persistens; sepalis cuspidatis.

Pet. ovalia, calyce breviora.

Stam. perigyna. Filamenta plana, apice subemarginata.

Antherae biloculares, loculis lateralibus discretis, longitudinaliter dehiscentibus.

Styli conici.

Fructus membranacei, monospermi?

Dilleniaceae.

1885. RECCHIA Scss. Cand. (1818.)

Cal. pentasepalus, patens; sepalis ovatis aequalibus.

Pet. oblonga, apice denticulata, basi attenuata.

Stam. perigyna.

Styli filiformes, breves. Stigmata capitata, dilatata.

Fructus?

Dilleniaceae.

886. GYPSOPHILA L. (1751.)

Cal. campanulatus, angulatus, quinquefidus, (interdum basi squamatus).

Pet. quinque spathulata, obovata, integerrima vel emarginata.

Stam. hypogyna, petalis breviora vel longiora. Antherae biloculares, erectae.

Styli capillares. Superficies stigmatica lateralis.

Caps. unilocularis, polysperma, apice quinquefariam dehiscens. Placenta centralis libera.

Caryophylleae. Schk. t. 120. *Kit.* plant. hungar. 1. t. 41. *Desfont.* fl. atlant. 1. t. 97. *Delil.* fl. aeg. t. 29. [Banffy Baumg. Reichenb. ic. 4. t. 400. abortu staminum alternorum et minore seminum numero differt.]

1887. *SAPONARIA* Trag. (1551.)

Cal. tubuloso - cylindricus, quinquedentatus, basi nudus.

Pet. unguiculata, integra, quandoque emarginata, rarissime bifida. Laminae basi interdum coronatae.

Stam. hypogyna. Filamenta filiformia, basi annulo cohaerentia. Antherae biloculares.

Styli filiformes. Superficies stigmatica lateralis.

Caps. oblonga, apice quinquefariam dehiscens, unilocularis, polysperma. Placenta centralis, libera.

Caryophylleae. Schk. t. 121. *Cav.* ic. 2. t. 134. *Marsch.* *Bieb.* pl. ross. cent. 2. t. 66. *Bot. mag.* 2855.

1888. *DIANTHUS* L. (1737.)

Cal. tubulosus, quinquedentatus, basi squamis imbricatis suffultus.

Pet. unguiculata, plerumque dentata aut multifida, raro integerrima.

Stam. annulo perigyno inserta. Filamenta capillaria. Antherae biloculares.

Styli filiformes. Superficies stigmatica lateralis.

Caps. oblonga, apice quadri-quinquefariam dehiscens, unilocularis, polysperma. Placenta centralis, libera.

Caryophylleae. Schk. t. 121.

1889. *CUNONIA* L. (1764.)

Cal. quinquepartitus, parvus.

Pet. ovato-oblonga, patentia.

Stam. hypogyna, exstantia. Filamenta filiformia. Antherae subrotundae.

Styli stigmatibus obtusis.

Caps. ovata acuminata, stylis coronata, a basi in duos folliculos corticatos dehiscens. Sem. plura triangularia.

Cunoniaceae. Lam. ill. t. 371. *Bot. reg.* 828. *Gärtn.* t. 225. [Oosterdykia Burm. afr. t. 96.]

1890. *HYDRANGEA* Gronov. (1743.)

Cal. germini adnatus, quinquedentatus, persistens.

Pet. subrotunda, calyce maiora.

Stam. alterna longiora. Filamenta filiformia. Antherae didymae.

Styli divergentes, persistentes.

Caps. infera, bilocularis, stylis birostrata, foramine inter stylas dehiscens. Placenta spongiosa. Sem. plura, oblonga.

Saxifrageae. Schk. t. 119. Gärtn. t. 30. Lam. ill. t. 370. Bot. mag. 437. 438. 975.

891. **SAXIFRAGA** Dioscor.

Cal. semisuperus, persistens, quinquefidus.

Pet. unguiculata vel sessilia; aequalia vel rarissime inaequalia.

Stam. calyci inserta, perigyna vel subepigyna. Filamenta filiformia. Antherae biloculares.

Styli simplices, subdivergentes. Stigmata obtusa, glabra aut barbata.

Caps. semiinfera, bilocularis, calyce vestita, polysperma.

Saxifrageae. Schk. t. 119. Lapeyrouse monographia Saxifrag. 1801. fol. Sternberg revisio Saxifrag. 1810. fol. [Bergenia Mönch., Megasea Haw., Geryonia Schrank., Robertsonia Haw., Gymnopera Don., Diptera Borkh. Link., Leiogyne et Micranthes Don.]

892. **TIARELLA** L. (1751.)

Cal. quinquepartitus, persistens, coloratus.

Pet. unguiculata, integrâ, calyci inserta.

Stam. calyci inserta, subpersistencia, exstantia. Filamenta capillaria. Antherae subrotundae.

Styli brevissimi.

Caps. unilocularis, bivalvis, valva altera longiore. Sem. plurima, ovata, nitida.

Saxifrageae. Bot. mag. 1589.

893. **MITELLA** Tournef. (1719.)

Cal. basi ventricosus, quinquefidus, persistens.

Pet. calyci inserta, angusta, pinnatifida.

Stam. calycis tubo affixa, inclusa. Antherae ovales biloculares.

Styli nulli. Stigmata obtusa.

Caps. unilocularis, bivalvis. Sem. suturis affixa.

Saxifrageae. Schk. t. 120. Lam. ill. t. 373. Gärtn. t. 44. Bot. reg. 166. [Tellima R. Br. Bot. reg. 1178. placētis parietalibus differt.]

II. *Flores imperfecti.*

894. **CHRYSOSPLENIUM** Tournef. (1719.)

Cal. corollinus superus, quadri-quinquelobus, patens.

Annulus perigynus subcarnosus undulatus.

Stam. sub annulo inserta, brevia. Antherae biloculares.

Styli brevissimi, divaricati. Stigmata obtusa.

Caps. bipartibilis, polysperma. Sem. fundo affixa.

Saxifrageae. Schk. t. 108. Engl. bot. 490.

1895. *TRIANTHEMA* Sauvag. (1751.)

Cal. corollinus, persistens, quinquepartitus; laciniis sub apice mucronatis.

Stam. imo calyci inserta, numero varia, ut plurimum decem, in duabus vel tribus speciebus quinque, in unica plura. Antherae reniformes.

Styli plerumque duo, (saepe unicus) persistentes.

Caps. bimucronata, bilocularis, supra basin circumscissa.

Sem. in singulis loculis bina, unum super altero.

Portulacaceae. *Cand. succul.* t. 109. *Lam. ill.* t. 375. *Gärtn.* t. 128. [*Zaleya* N. L. *Burm. ind.* t. 31. *Rocama* et *Papularia* Forsk.]

1896. *SCLERANTHUS* L. (1737.)

Cal. urceolatus, quinquefidus, persistens.

Stam. fauci calycis inserta. Antherae biloculares.

Achenium calycis basi indurata tectum.

Paronychiaee. *Schk.* t. 120. *Gärtn.* t. 126.

T R I G Y N I A.

1897. *HOLOLACHNA* Ehrenb. (1827.)

Cal. campanulatus, quinquefidus, bracteatus, persistens.

Cor. pentapetala, receptaculo affixa.

Stam. glandulae hypogynae inserta.

Styli breves subulati (duo, tres, quatuor).

Caps. bi-quadrangulata, bi-quadrilocularis. *Sem.* pauca, pilosa, basi affixa.

Tamariscineae. [*Tamarix songarica* Pall. in nov. act. petrop. 10. t. 10. f. 4.]

1898. *BEMBIX* Lour. (1790.)

Cal. inferus, tripartitus.

Pet. quinque, ovata, concava.

Stam. quinque, alterna longiora, petalorum basi inserta.

Filamenta filiformia. Antherae biloculares, erectae.

Styli oblongo-turbinati. Stigmata verticaliter compressa, emarginata.

Bacca ovata trilocularis.

1899. *DEUTZIA* Thunb. (1781.)

Cal. inferus, campanulatus, quinquefidus; laciniis ovatis, obtusis, erectis.

Petala receptaculo inserta, oblonga, obtusa, (quinque aut sex).

Stam. hypogyna. Filamenta linearia, sub apice emarginata bi- (tri-) fida. Antherae globosae, didymae.

Styli filiformes. Stigmata clavata.

Caps. trigona, stylis coronata, trilocularis, trivalvis, basi dehiscens; loculis pleiospermis.

— — *Thunb. diss. nov. gen.* t. 1.

900. *ERYTHROXYLON* P. Brown. (1756.)*Cal.* campanulatus, quinquepartitus.*Pet.* quinque hypogyna, basi squamata.*Stam.* urceolo perigyno inserta. Filamenta capillaria. Antherae ovatae, biloculares, erectae, longitudinaliter dehiscentes.*Styli* filiformes. Stigmata capitata. Interdum in unum conflunt.*Drupa* monosperma. Sem. angulatum.*Erythroxyleae* Kunth. *P. Brown.* *jam.* t. 38. f. 2. *Cav.* diss. t. 225 — 233. *Vahl.* *symb.* 3. t. 63. *Roxb.* *corom.* 1. t. 88. *Humb.* *nov. gen.* 5. t. 453.901. *MALPIGHIA* Plum. (1703.)*Cal.* hemisphaericus, quinquefidus, extus glandulis decem stipatus.*Pet.* quinque orbiculata, unguiculata, patentia.*Stam.* hypogyna. Filamenta basi cohaerentia. Antherae subrotundae.*Styli* simplices. Stigmata capitata, vel uncinata.*Drupa* tripyrena.*Malpighiae.* *Cav.* diss. t. 234 — 236.902. *BYRSONIMA* Rich. (1811.)*Cal.* hemisphaericus, quinquefidus, extus glandulis decem stipatus.*Pet.* quinque orbiculata, unguiculata, patentia.*Stam.* hypogyna. Filamenta basi cohaerentia. Antherae subrotundae.*Styli* simplices. Stigmata acuta, subulata.*Drupa* trilocularis, loculis monospermis.*Malpighiae.* *Humb.* *nov. gen.* 5. t. 446 — 449. [*Malpighia Aubl. guian.* t. 181 — 184. *Cav.* diss. t. 237. 240. 241.]903. *GALPHIMIA* Cav. (1799.)*Cal.* quinquepartitus, persistens, eglandulosus.*Pet.* quinque subrotunda, unguiculata.*Stam.* hypogyna. Filamenta subulata, basi subcohaerentia. Antherae ovatae, erectae, basi emarginatae.*Styli* tres simplices.*Caps.* tricocca, coccis monospermis, dorso longitudinaliter dehiscentibus.*Malpighiae.* *Cav.* in anal. de cienc. nat. 1. t. 2. ic. 5. t. 489. vol. 6. t. 563. *Humb.* *nov. gen.* 5. t. 452. [Huc pertinere *Thryallis* *Lin.* videtur. Ipse e *Marcgrav.* p. 79. ed. 1648. sumsisse videtur. Nec in *Linnaei* collectione, nec usquam alibi conspecta est eiusmodi planta. Nuper vero *Lindleyus* *Bot. reg.* 1162. invenisse eam sibi visus est. Congruit tamen icon cum *Galphiinia*, ipsaque trigyna est. De *Caucantho* *Forsk.* multo magis dubitandum: drupam enim ferre incolae asseverarunt.

1904. *BANISTERIA* Houst. L. (1737.)

Cal. quinquepartitus, extus glandulis decem stipatus, persistens.

Pet. subrotunda, unguiculata.

Stam. hypogyna. Filamenta subulata, basi connata.

Antherae ovals, erectae.

Styli simplices. Stigmata dilatata, subfoliacea.

Samarae tres, -axi centrali affixae, unialatae.

Malpighiaceae. Cav. diss. t. 246 — 257. [Heteropteris Kunth. in Humb. nov. gen. 5. t. 450.]

1905. *TRIOPTERIS* Plukn. (1700.)

Cal. quinquepartitus, extus glandulis decem stipatus.

Pet. subrotunda, unguiculata.

Stam. hypogyna, alterna longiora. Filamenta subulata, basi connata. Antherae ovals.

Styli simplices.

Samarae tri-quadrilatae, alis aequalibus vel inaequalibus.

Malpighiaceae. Cav. diss. t. 259. Humb. nov. gen. 5. t. 451. [Tetrapteris Cav. diss. t. 260 — 262.]

1906. *HIRABA* Jacqu. (1763.)

Cal. quinquepartitus, extus saepe glandulis stipatus.

Pet. subrotunda, unguiculata.

Stam. hypogyna. Filamenta basi subcohaerentia.

Styli simplices.

Samarae tres, dorso cristatae, ala membranacea cinctae.

Malpighiaceae. Roxb. corom. 2. t. 160. [Flabellaria Cav. diss. t. 264. Mascagnia Berter. in Coll. hort. ripul. p. 86.]

1907. *SCHINUS* L. (1737.)

Flores dioeci.

Cal. quinquepartitus, patens.

Pet. quinque ovalia, unguiculata.

Stam. filiformia. Antherae subrotundae.

Styl. nullus. Stigmata tria vel quatuor.

Drupa globosa, monosperma.

Terebintheae. Gärtn. t. 140. [Molle Monard. in Clus. exot. p. 322. Clus. cur. post. p. 94.]

1908. *STELLARIA* L. (1745.)

Cal. quinquepartitus.

Pet. quinque, bipartita (quandoque biloba).

Stam. hypogyna, basi libera. Filamenta filiformia. Antherae biloculares, dorso affixae.

Styli acuti. Stigmata lateralialia.

Caps. unilocularis, apice quinque-sexfariam dehiscens.

Placenta basilaris.

Caryophylleae. Schk. t. 122. [Schiedea Chamiss. Schlechtend.]

909. *ARENARIA* Chabr. (1666.) Rupp. (1718.)*Cal.* pentasepalus.*Pet.* quinque, integra.*Stam.* hypogyna, numero varia, (plerumque decem, interdum alterna abortientia, aut quinque).*Styli* simplices. Stigmata lateralialia.*Caps.* apice tri- quinque-sexfariam dehiscens, aut tri-sexvalvis. Placenta basilaris vel centralis. Sem. numerosa, vel pauca. (A. bavarica *L.* semina terna; A. polygonoides *Wulff.* semina quaterna, terna vel bina gerit; vel, abortiente staminum dimidia parte et styliorum triente, Gouffea dicitur arenarioides *Cand. Reichenb.* ic. 4. t. 399. *Brachystemma calycinum Don.* simili modo exponendum est ac *Alsine Linnaei.*)*Caryophylleae.* Schk. t. 122, Sturm. fl. germ. [*Alsine L.*, *Spergularia Pers.* *Lepigonum Fries.*]910. *SILENE*. [Nomen plane fictitium, cuius prima mentio fit in *Lobel. hist.* p. 242. (1576), ascriptis auctoritatibus Theophrasti, qui nusquam id nomen habet, et Aldroandi (*Aldrovandi*), qui tunc temporis saltem nihil ediderat.]*Cal.* cylindricus, clavatus vel inflatus; striatus vel sulcatus; quinquedentatus; persistens.*Pet.* quinque bipartita, bifida, biloba; vel integra; ad laminarum basin appendiculis erectis ornata, vel nuda.*Stam.* hypogyna. Filamenta alterna basi petalorum inserta, alterna basi annulo coniuncta. Antherae ovales, biloculares.*Styli* simplices. Stigmata lateralialia.*Caps.* basi trilocularis, polysperma; apice subsexfariam dehiscens. Placenta columnaris, centralis.*Caryophylleae.* Schk. t. 122. *Gärtn.* t. 130. [*Cucubali L.* pleraeque species. *Viscago Dillen. Hornem.*]911. *CUCUBALUS* Dalech. (1587.)*Cal.* ventricosus, quinquedentatus, persistens; dentibus revolutis.*Pet.* bifida, ad ungues appendiculata.*Stam.* alterna basi petalorum inserta, alterna perigyna.*Styli* filiformes, persistentes.*Caps.* baccata, uniloculares, polysperma. Placenta centralis, libera.*Caryophylleae.* Eng. bot. 1577. *Gärtn.* t. 77.912. *CHEBLERIA* Haller. (1740.)*Cal.* pentasepalus, striatus.*Pet.* quinque minima, emarginata, seu squamae nectariferae et staminiferae.*Stamina* alterna petalis inserta.*Styli* filiformes.

Caps. trivalvis, oligosperma.

Caryophylleae. Hall. stirp. helv. t. 21. Engl. bot. t. 1212.

1913. *GABIDELIA* Tournef. (1719.)

Cal. pentasepalus, coloratus, deciduus.

Pet. seu nectaria quinque petaloidea, bilabiata; labio altero bifido elongato, altero breviori integro.

Stam. hypogyna, decem aut plura.

Styli brevissimi.

Caps. tres folliculares, pleiospermae, ita nexae, ut capsulam trilocularem mentiantur.

Ranunculeae. Bot. mag. 1266. Gärtn. t. 118.

1914. *BRUNNICHIA* Banks. Gärtn. (1788.)

Cal. ampullaceus, coriaceus, persistens, quinquefidus; laciniis revolutis, obtusiusculis. Pedunculus sub calyce arcuatus, hinc alatus, inde sulcatus.

Pet. nulla.

Stam. decem vel octo.

Caps. pyramidata, trigona, monosperma. Sem. pyramidatum, sexlobatum.

— — Gärtn. t. 45.

1915. *QUERIA* Löffl. (1752.)

Cal. quinquepartitus.

Pet. nulla.

Stam. decem inaequalia, interdum quinque sterilia, interdum tria sola perfecta.

Styli gracillimi.

Caps. unilocularis, trivalvis, monosperma.

Paronychieae. Lam. ill. t. 52.

1916. *DIAPHORA* Lour. (1790.)

Flores spicati, androgyni.

Cal. uniflorus, trivalvis; valva tertia aristata.

Cor. bivalvis, calyce longior, mutica.

Filamenta subnulla. Antherae filiformes, inaequales, apice caudato discolore. Paleae numerosae adstantes, seu stamina dixeris sterilia.

Styl. nullus. Stigmata tria filiformia.

Caryopsis trigona.

Gramen.

T E T R A G Y N I A.

1917. *MICROPETALON* Pers. (1805.)

Cal. pentasepalus.

Pet. quinque minutissima integra, aut nulla.

Stigmata quatuor sessilia subligulata.

Caps. ovata, quadrivalvis.

Caryophylleae. [Spergulastrum Mx.]

P E N T A G Y N I A.

918. CERASTIUM Dillen. (1719.)

Cal. pentasepalus aut quinquepartitus, persistens.

Pet. bifida, calyce minora, vel eum aequantia aut superantia.

Stam. alterna hypogyna, alterna basi petalorum inserta.

Styli capillares. Stigmata lateralialia.

Caps. globosa, vel cylindrica, unilocularis, polysperma; apice dentibus decem erectis vel revolutis dehiscens.

Placenta centralis, libera.

Caryophylleae. Schk. t. 125. Reichenb. ic. t. 181. 230. 233. 244. 245. [Larbrea Hilar.]

919. LYCHNIS Theophr.

Cal. tubulosus, quinquedentatus; vel cylindrico-clavatus, costatus; vel campanulatus costatus, aut coriaceus, vel inflatus ecostatus.

Pet. quinque unguiculata, ad laminarum basin appendiculata, plerumque bifida, nonnunquam laciniata, interdum integra.

Stam. hypogyna, basi annulo unita. Antherae oblongae, biloculares, dorso affixae.

Styli filiformes. Stigmata lateralialia.

Caps. unilocularis, apice dentibus decem dehiscens. Placenta centralis libera.

Caryophylleae. Schk. t. 124. Gärtn. t. 130. [Agrostemma L. Hedona Lour.]

920. SPERGULA Dodon. (1583.)

Cal. quinquepartitus, persistens.

Pet. quinque, integra.

Stam. hypogyna. Antherae biloculares.

Styli filiformes, superne undique stigmatosi.

Caps. unilocularis, semiquinquevalvis. Placenta centralis, libera. Sem. quandoque marginata.

Caryophylleae. Schk. t. 125. Fl. dan. 1930.

921. BERGIA L. (1771.)

Cal. quinquepartitus, persistens.

Pet. quinque lanceolata, patula, calycem aequantia.

Stam. hypogyna.

Styli brevissimi, approximati.

Caps. quinquelocularis, quinquevalvis, valvis patentissimis.

Caryophylleae. Roxb. corom. 1. t. 142. [Elatine Delil. fl. aeg. t. 26.]

922. OXALIS Diosc. (de Rumice) L. (1737.)

Cal. quinquepartitus.

Pet. quinque hypogyna, aequalia, unguibus quandoque cohaerentia.

Stam. hypogyna, alterna breviora; annulo basilari saepe unita. Antherae biloculares, erectae.

Styli erecti, staminibus vel omnibus vel alternis longiores brevioresve. Stigmata capitata, aut emarginata.

Caps. oblonga, pentagona, quinquelocularis, decemvalvis. Semina pauca, angulo loculorum centrali affixa, iuniora arillo carnoso praedita, cuius vi proiciuntur.

Oxalideae. Schk. t. 123. *Jacquin.* monogr. 1792. [*Biophytum* Cand.]

1923. *LEDOCARPON* Desf. (1818.)

Cal. pentasepalus persistens, involucre cinctus foliolis bipartitis linearibus.

Pet. quinque, ovali-oblonga.

Stam. hypogyna, persistentia, alterna breviora. Antherae oblongae, biloculares.

Styli incrassati.

Caps. quinquevalvis; valvis introflexis totidem loculos formantibus. Placenta centralis. Sem. numerosa.

Oxalideae. Desf. in mém. mus. 4. t. 13. [*Balbisia verticillata* Cav. in anal. de cienc. nat. 7. t. 46.]

1924. *AVERRHOA* L. (1747.)

Cal. quinquepartitus vel pentasepalus.

Pet. patentia, calyce longiora.

Stam. alterna longiora, basi in annulum iuncta. (In *A. Carambola* stamina alterna abortiunt.) Antherae subrotundae.

Styli simplices. Stigmata capitata.

Bacca magna, quinesulcata, quinquelocularis; loculis paucifloris. Sem. angulo centrali adnata.

Oxalideae R. Br. *Corr.* in aun. mus. 8. t. 2. *Cav.* diss. t. 219. 220.

1925. *DROSOPHYLLUM* Link. (1806.)

Cal. pentasepalus.

Pet. quinque, unguibus approximatis.

Stam. hypogyna, alterna longiora. Antherae oblongae, incumbentes.

Styli filiformes. Stigmata capitata.

Caps. unilocularis, quinquevalvis. Placenta centralis.

Droseraeae. Hilar. in mém. mus. 2. t. 4. f. 13. [*Drosera lusitanica* L. *Spergula rorida* Griseb. Sp. *droseroides* Brot.]

1926. *GRIELUM* L. (1764.)

Cal. quinquefidus, in fructum abiens.

Pet. calyci inserta.

Stam. ibidem inserta, extra annulum glandulosum, persistentia; spinescentia.

Styli breves, persistentes. Stigmata capitata.

Caps. calyce indurato vestita, staminibus spinescentibus stylisque coronata, decemlocularis, loculis monospermis.

Rosaceae. Sweet. *Geran.* 2. t. 171. *Gärtn.* t. 36.

927. BIEBERSTEINIA Stephan. (1811.)

Cal. pentasepalus, irregularis, persistens.*Pet.* unguiculata.*Stam.* hypogyna. Filamenta subulata. Antherae incumbentes, versatiles.*Styli* simplices.*Caryopses* quinque?*Rosaceae?*

928. SURIANA Plum. (1703.)

Cal. pentasepalus, persistens.*Pet.* calyci inserta, obovata, patentia.*Stam.* nonnulla abortientia.*Styli* laterales. Stigmata obtusa.*Nuculae* quinque.*Rosaceae?* Plum. ic. t. 249. f. 1.

929. QUILLATA Molin. (1782.)

Flores polygami.

Cal. quinquefidus persistens.*Pet.* quinque decidua. Discus quinquelobus calycis basin tegens.*Stam.* apici disci inserta.*Styli* subconnati, subulati.*Caps.* quinque, trigonae, coriaceae. Sem. biseriata, imbricata, apice alata.*Rosaceae.* Kunth. *Cand.* [Smegmadermos *R.* et *P.* Smegmaria Pers. *W.* Aliam habet Molina (saggio sulla storia nat. del Chili, ed. 2. p. 162.) *Quillajam*, quae ad oram maritimam crescat, quam monoecam dicit, cuique tribuit calycem tetrasepalum, corollam nullam, stamina duodecim, stylos quatuor, capsulam quadrilocularem (p. 298.) Hanc male descriptam esse existimat Candolleus (prodr. 2. p. 547.)]

930. POUPARTIA Commers. Juss. (1789.)

Flores diclini.

Cal. quinquefidus, minimus.*Pet.* quinque patentissima.*Stam.* disco crenato hypogyno inserta.*Styli* approximati.*Drupa*, nucleo bi-quinqueloculari.*Terebintheae.*

931. CNESTIS Juss. (1789.)

Cal. quinquepartitus, extus tomentosus.*Pet.* hypogyna, lanceolata, reflexa.*Stam.* hypogyna. Antherae ovatae, biloculares.*Styli* divergentes. Stigmata capitata.*Caps.* quinque leguminiformes, stylo coronatae, extus villosae-prurientes, dorso debiscentes, mono-dispermae.*Terebintheae.* Lam. ill. t. 387. [Byrsocarpus Schumacher. pl. guin. p. 226.]

1932. *TAPIRIA* Juss. (1789.)

Cal. pentasepalus, sepalis subrotundis deciduis.

Pet. quinque, disco annulari hypogyno annexa.

Stam. ibidem inserta, alterna petalis, sepalis alterna opposita. Antherae biloculares.

Stylus nullus. Stigmata quinque.

Caps. quinesulcata, quinquevalvis, pentasperma. Sem. valvis affixa.

Terebintheae. [Tapirira Aubl. guian. t. 188. Joncquetia Schreb.]

1933. *ROUREA* Aubl. (1775.)

Cal. quinquepartitus, extus tomentosus.

Pet. quinque patentissima.

Stam. hypogyna, basi calycis inserta, in anulum connata, alterna breviora. Antherae dorso affixae, biloculares.

Styli elongati. Stigmata dilatata.

Fructus oblongus, coriaceus, follicularis, monospermus.

Terebintheae? Aubl. guian. t. 187. [Robergia Schreb.]

1934. *SPONDIAS* L. (1737.)

Flores diclini.

Cal. parvus, quinquefidus, coloratus, deciduus.

Pet. quinque, sub disco inserta, patentia.

Stam. sub disco crenulato inserta. Antherae biloculares.

Styli brevissimi. Stigmata obtusa.

Drupa putamine fibroso vel echinato, quineloculari, pentaspermo.

Terebintheae. Sonnerat it. ind. 2. t. 123. Gärtn. t. 103-104.

1935. *KIGGELARIA* L. (1737.)

Flores dioeci.

Cal. quinquepartitus.

Pet. quinque, unguibus glandulifera.

Stam. Filamenta brevissima. Antherae oblongae, apice biporosae.

Styli simplices. Stigmata obtusa.

Caps. baccata, unilocularis, valvis quatuor dehiscens?

Sem. angulata, valvis adnata?

Flacourtiaceae Cand. Linn. hort. Cliff. t. 29. Jacqu. ic. rar. 3. t. 628. Gärtn. t. 44.

1936. *SEDUM* Colum. Plin.

Cal. quinquepartitus, persistens.

Pet. quinque lanceolata.

Squamae hypogynae nectariferae.

Stam. basi petalorum inserta. Antherae biloculares.

Stigmata germinum apices.

Caps. quinque patentes, sutura interna dehiscentes. *Sem.* minuta, plurima, ad angulum internum fixa.

Sedae. Schk. t. 123. Gärtn. t. 65. [Anacampeeros
Tourn. Hau. Rhodiola L.]

937. COTYLEDON Nicand.

Cal. quinquefidus aut quinquepartitus.

Cor. tubulosa, campanulata; limbo quinquefido.

Stam. tubo adnata, alterna plerumque breviora. *Antherae* subrotundae.

Styli subulati.

Caps. quinque patentes, sutura interna dehiscentes. *Sem.* minuta, plurima, ad angulum internum fixa.

Sedae. Engl. bot. 325. 1522.

Orostachys Fisch. et *Echeveria Cand.* corollis profundius partitis sectionem constituunt, non genus. [Umbilicus et *Pistorinia Cand.* solo habitu differunt.]

938. PENTHORUM Gronov. (1743.)

Cal. quinquefidus.

Pet. quinque, linearia.

Stam. basi petalorum inserta, subulata.

Stigmata obtusa.

Caps. quinquelocularis, quinquerostrata, polysperma.

Sedae. Lam. ill. t. 390. Gärtn. t. 65.

939. LANCRETIA Delil. (1813.)

Cal. quinquepartitus.

Pet. quinque, calyce paullo longiora.

Stam. hypogyna, alterna breviora. *Antherae* cordatae.

Styli breves, filiformes.

Caps. quinquelocularis, marginibus valvarum introflexis et dissepimenta et placentas constituentibus.

Hypericeae. Delil. fl. aeg. t. 25.

940. CORIARIA Nissol. (1711.)

Flores diclini aut polygami.

Cal. campanulatus, decemfidus, laciniis alternis minoribus callosis.

Cor. nulla.

Stam. receptaculo inserta. *Antherae* biloculares.

Styl. nullus. *Stigmata* quinque subulata.

Caryopses quinque, laciniis internis calycis auctis tectae.

— — Mém. de l'ac. de Paris, 1711. t. 12.

941. ASTRANTHUS Lour. (1790.)

Cal. breviter tubulosus; limbo decem - quatuordecimfido, laciniis alternis minoribus.

Cor. nulla.

Stam. decem aut plura. *Antherae* subrotundae.

- Styli capillares.*
Caryopate in tubo calycis inclusa.
 — Bot. mag. 2659.

D E C A G Y N I A.

1942. *NEVNADA* L. (1748.)

- Cal.* quinquepartitus superus.
Pet. quinque calyci inserta.
Caps. calyce echinato cincta, decemlocularis, decaspe
Rosaefolia? Lam. ill. t. 393. Gärtn. t. 32.

1943. *GASTONIA* Commers. Juss. (1789.)

- Cal.* margine elevato, integro.
Pet. quinque aut sex.
Stam. ad basin petali cuiusvis bina.
Styli minimi, basi iuncti.
Caps. infera, decemlocularis.
Araliace.

1944. *PHYTOLACCA* Tournef. (1719.)

- Cal.* quinquepartitus, coloratus, persistens, inferus.
Stam. et *Styli* numero varii.
Bacca depressa, sulcata, decemlocularis. Sem. solida
Chenopodeae. Schk. t. 126. Gärtn. t. 77.

CLASSIS XI. ODECANDRIA.

CONSPECTUS

CLASSIS UNDECIMAE.

MONOGYNAE.

oribus perfectis.
 Corolla monopetala.
 Fructibus superis. 1945 — 1947.
 Corolla pleiopetala.
 Fructibus superis. 1948 — 1976.
 Fructibus inferis. 1977 — 1981.
 oribus imperfectis.
 Fructibus superis. 1982 — 1987.,
 Fructibus inferis. 1988.
 Perianthio nullo. 1989.

DIGYNAE.

1990. — 1993.

TRI - TETRAGYNAE.

1994 — 2000.

PENTAGYNAE.

2001 — 2008.

HEXAGYNAE.

2009, 2010.

DODECAGYNAE.

2011.

BOLYGNAE.

2012 — 2014.

MONOGYNIA.

I. *Flores perfecti.*

A. COROLLA MONOPETALA.

Fructus superus.

HOPEA Roxb. (1819.)

l. pentasepalus, sepalis duobus demum in alas fructus
 xrescentibus.

r. quinquepartita, contorta.

vm. Filamenta alterna bifida.

gma simplex.

Caps. monosperma.

— — *Roxb. corom. 3. t. 210.*

1946. *BASSIA* L. (1771.)

Cal. tetrasepalus, coriaceus, persistens.

Cor. campanulata, tubo ventricoso; limbo octopartito, erectiusculo.

Stam. tot fauci, quot tubo corollae inserta. Antherae lineares, sagittatae, introrsum villosae.

Styl. subulatus, corolla longior.

Bacca carnosae, quinquelocularis; loculis monospermis.

— — *Roxb. corom. 1. t. 13. Gärtn. t. 104.*

1947. *BAITARIA* Ruiz et Pav. (1794.)

Cal. disepalus, bibracteatus, persistens.

Cor. tubulosa, limbo quinquefido.

Stam. tubo corollae inserta, inclusa.

Styl. simplex. Stigma trifidum.

Caps. trigona, trilocularis, polysperma. Sem. dissepimentis affixa.

B. COROLLA PLEIOPETALA.

a. *Fructus superi.*

1948. *ACIDANDRA* Mart. (1826.)

Cal. integer, coriaceus, lateraliter fissus, demum deciduus.

Pet. quinque obovata, duo superiora paullo breviora.

Stam. hypogyna. Filamenta brevia subulata. Antherae lineares, sagittatae.

Styl. brevis, recurvus. Stigma acutum.

Legumen stipitatum, oligospermum.

Leguminosae. [Zollernia Nees in nov. act. nat. cur. 13 t. C. D. Krameria glabra N. E.]

1949. *XEROPETALUM* Delil. (1826.)

Cal. quinquefidus, patens.

Pet. quinque, obovata, emarginata, nervosa, persistentia.

Stam. viginti, quorum quinque abortiunt. Filamenta basi coalita. Antherae lineares.

Styl. brevis. Stigmata duo aut tria filiformia, spiratum reflexa.

Caps. trilocularis, trivalvis, loculis oligospermis.

Tiliaceae.

1950. *HELIOCARPUS* L. (1737.)

Cal. tetrasepalus.

Pet. quatuor, calyce breviora.

Stam. hypogyna. Antherae lineares, incumbentes, didymae.

Styl. bifidus. Stigmata acuta.

Caps. stipitata, compressa, longitudinaliter radiata, bilocularis bivalvis; dissepimento contrario. Sem. solitaria.
Tiliaceae. Linn. hort. Cliff. t. 16. Gärtn. t. 49.

951. *VATICA* L. (1771.)

Cal. quinquepartitus, laciniis lanceolatis.

Pet. obovata, subcontorta.

Stam. Antherae sessiles, quadriloculares; loculis exterioribus spinula interiecta terminatis.

Styl. cylindricus. Stigma obtusum.

Caps. trilocularis, trisperma?

Tiliaceae. Smith. ic. ined. t. 36.

952. *TRIUMFETTA* Plum. (1703.)

Cal. pentasepalus; sepalis sub apice mucronatis, deciduis.

Pet. quinque, aut rarius nulla.

Stam. Filamenta subulata erecta. Antherae simplices.

Styl. simplex. Stigma bifidum.

Glandulae quinque perigynae.

Caps. globosa, setis uncinatis echinata; quadrilocularis; loculis mono-dispermis.

Tiliaceae. Lindl. coll. t. 29. Humb. nov. gen. 5. t. 488. Gärtn. t. 111.

953. *CRATEVA* L. (1737.)

Cal. tetrasepalus, caducus.

Pet. unguiculata, saepe ad idem latus deflexa.

Stam. disco perigyno glanduloso inserta. Filamenta longissima. Antherae lineares, subcurvatae, basi emarginatae.

Germen longe stipitatum. Styl. nullus. Stigma sessile.

Bacca pleiosperma.

Capparideae. Andr. repos. t. 176. [Ubi petala parva aut nulla sunt, dicitur Niebuhria Cand. et Boscia Lam. Podoria Pers. Absentia glandulosae indolis disci differre dicitur. Othrys Noron. apud Thuars.]

954. *GARCINIA* L. (1737.)

Cal. tetrasepalus, persistens.

Pet. quatuor, subrotunda, concava.

Stam. hypogyna, basi subconnata. Antherae ovatae.

Styl. nullus. Stigma quadri-oculobum.

Bacca quadri-sex-oculocularis; loculis monospermis.

Sem. testa duplici obvoluta. Cotyledones conferruminatae, ut monocotyledonea mentiantur.

Guttiferae. Lam. ill. t. 405. [Mangostana Gärtn. t. 105. Oxycarpus Lour. Brindonia Thuars.]

955. *CRENEA* Aubl. (1775.)

Cal. turbinatus, semiquadrifidus, persistens.

Pet. subrotunda, calyci inserta.

Stam. ibidem inserta. Filamenta, capillaria. Antherae subrotundae.

Styl. filiformis, incurvus. Stigma oblongum crassiusculum.
Caps. uni- aut quinquelocularis, calyce vestita, polysperma.
Salicariae. *Aubl. guian.* t. 209.

1956. *ANTHERYLUM* Rohr. Vahl. (1792.)

Cal. quadripartitus, aequalis.

Pet. cum laciniis calycis alternantia.

Stam. imo calyci inserta.

Styl. filiformis.

Caps. tri-quadrialvis, polysperma. Placenta crassa tritetragona.

Salicariae. *Act. soc. hist. nat. havn.* 2. t. 8.

1957. *LYTRUM* L. (1737.) *Diosc.* 4, 3. *Λυτρώμιον*, οὐ λύτρον καλοῦσι.

Cal. tubulosus, angulato-striatus; limbo duodecimdentato; dentibus alternis longioribus ab externo calycis integumento, minoribus alternis ab interna eius membrana productis.

Pet. ut plurimum sex, calycis limbo imposita.

Stam. e medio aut imo calyce orta, alterna longiora, sena vel duodena.

Styl. filiformis. Stigma capitatum.

Caps. calyce vestita, bilocularis, polysperma.

Salicariae. (*Schk.* t. 128.)

1958. *CUPHEA* P. Brown. (1756.)

Cal. tubulosus, basi gibbus; limbo sub sex-duodecimdentato; dente supremo latiori.

Pet. sex, calyci inserta.

Stam. decem ad quatuordecim, fauci calycis inserta, inaequalia. Antherae biloculares.

Stylus filiformis. Stigma simplex aut subbilobum. Glaccula ad basin ovarii.

Caps. calyce oblecta, unilocularis, demum per placentam deflexam simul cum calyce fissa. Sem. lenticularia.

Salicariae. *Cav. ic.* 4. t. 380—382. *Humb. nov. gen.* 6 t. 550—552. *Hook. exot. fl.* t. 161. *Bot. reg.* 182. 852.

[*Melanium* et *Parsonsia* P. Brown. *javn.* t. 21. f. 2. *Balsamona* Vand. in *Röm. script.* t. 4.]

1959. *PEMPHIS* Forst. (1775.)

Cal. campanulatus, sulcatus; limbo ampliato duodecimdentato; dentibus alternis minoribus.

Pet. sex obovata, calyci inserta.

Stam. duodecim medio calyci inserta, alterna breviora.

Styl. brevis. Stigma capitatum.

Caps. calyce vestita, sexvalvis, polysperma. Placenta triangularis, basilaris.

Salicariae. *Forst. char.* t. 35.

1960. *NESAEA* Commers. Juss. (1789.)

Cal. campanulatus, limbo duodecimdentato; dentibus alternis mucronatis.

Pet. sex.

Stam. medio calyci inserta.

Styl. filiformis, persistens. Stigma capitatum.

Caps. calyce oblecta, quadrilocularis; polysperma. Placenta globosa centralis.

Salicariae. Reichenb. hort. 2. t. 166. [Heimia Link. ic. t. 28. Decodon Gmel. staurinup. duorum. et loculi unius defectu differt.]

961. GINORIA Jacqu. (1763.)

Cal. campanulatus, persistens, limbo sexfido patente, laciniis lanceolatis dorso dentato - cornutis.

Pet. subrotunda, longe unguiculata, oculo calycis inserta.

Stam. calyci inserta. Filamenta filiformia. Antherae reniformes, quadriloculares.

Caps. unilocularis, quadrivalvis, apice dehiscens, polysperma. Placenta centralis magna.

Salicariae. Jacqu. amer. t. 91.

962. DODECAS L. fil. (1781.)

Cal. quadrifidus.

Pet. quatuor, calyci inserta, suborbiculata.

Stam. tubo calycis adnata. Antherae oblongae.

Caps. globosa, unilocularis, semiquadrivalvis, polysperma. Placenta centralis globosa.

Salicariae. E. Meyer in nov. act. nat. cur. 12. p. 800.

963. DIPLODON * (Diplusodon Pohl. 1827.)

Cal. bibracteatus, hemisphaerico - campanulatus, nervosus, laciniis sex triangularibus; dentibus dorsalibus vel mucronatis, vel linearibus subulatis, vel lanceolato - linearibus elongatis.

Petala subrotundo - obovata, undulato - crenata, calyci inserta.

Stam. calyci inserta. Filamenta filiformia. Antherae semilunatae.

Styl. simplex. Stigma concavum.

Caps. bilocularis, bivalvis. Sem. marginata.

Salicariae. Pohl. fl. brasil. t. 66 - 81. [Friedlandia Schlechtend.]

964. SYMMETRIA Blum. (1826.)

Cal. campanulatus, sexfidus.

Pet. sex parva, calyci inserta.

Stam. alterna breviora.

Styl. crassus. Stigma quadrangulare.

Drupa baccata, calyce vestita, uni - trilocularis; loculis - monopyrenis.

Salicariae.

965. BLAKEA P. Brown. (1756.)

Cal. membranaceus, sexangularis, limbo repando, basi cinctus involucre tetra - aut hexaphyllo.

Pet. sex.

Stam. hypogyna. Antherae magnae in orbem coalitae, apice biporosae, basi appendiculatae.

Styl. simplex. Stigma peltatum.

Caps. baccata, calyce vestita, sexlocularis. Sem. ovato-angulata.

Melastomeae. *P. Brown*. *Jam.* t. 35. *Aubl. guian.* t. 210.

[*Topobea* *ib.* t. 189. *Valdesia* *R. et P. prodr.* t. 11.]

1966. *AXINAEA* *R. et P.* (1794.)

Cal. urceolatus, basi nudus, limbo persistente, quinque-sexdenticulato.

Pet. quinque aut sex.

Antherae liberae, apice biporosae, basi appendiculatae.

Caps. libera, quinque-sexlocularis.

Melastomeae.

1967. *BEFARIA* *Mutis. L.* (1774.)

Cal. septempartitus, persistens.

Pet. septem, obovata.

Stam. hypogyna. Antherae obovatae, basi poro gemino dehiscentes.

Styl. elongatus. Stigma capitatum.

Caps. globosa, calyce cineta, stylo coronata, septemlocularis; loculis polyspermis.

Briceae. *Michaux. fl. bor. amer.* t. 26. *Humb. plant. aequin.* 2. t. 118—121. 177.

1968. *ELAEOCARPUS* *Burm.* (1737.)

Cal. quinquepartitus.

Pet. quinque trifida, vel multifido-laciniata.

Discus glandulosus perigynus.

Stam. decem ad triginta, disco glanduloso inserta. Filamenta capillaria brevissima. Antherae lineares, apice bivalves; valvis apice setigeris vel ciliatis.

Styl. subulatus. Stigma simplex.

Drupa oblonga, putamine bi-trivalvi, biloculari interdum, raro quadriloculari.

Elaeocarpeae *Juss. Burm. zeyl.* t. 40. *Bot. mag.* 1737.

Bot. reg. 657. *Labill. nov. holl.* t. 155. [*Gauitrus* *Gärtn.*

t. 139. *Adenodus* et *Craspedum* *Lour.* *Dicera* *Forst.* t. 40.

Friesia *Cand.* *Eriostenum* *Coll. hort. ripul.* t. 30.]

1969. *TRICUSPIS* *Pers.* (1805.)

Cal. quinquedentatus.

Pet. quinque tricuspidata.

Discus glandulosus perigynus decagonus.

Stam. Antherae muticae, apice dehiscentes.

Caps. trilocularis, trivalvis, oligosperma.

Elaeocarpeae. [*Tricuspidaria* *Ruiz et Pav. prodr.* t. 36.]

1970. *EURYA* *Thunb.* (1784.)

Cal. pentasepalus, bracteis binis suffultus.

Pet. quinque, subrotunda.

- Annulus* glandulosus perigynus.
Stam. annulo inserta. Filamenta brevissima. Antherae lineares, tetragonae.
Styl. subulatus. Stigmata tria reflexa.
Caps. baccata tri-quinquelocularis. Sem. triangularia.
Ternströmiae Cand. Thunb. fl. jap. t. 25. [Geeria Blum.]
971. *ARISTOTELEA* Herit. (1784.)
Cal. quinquepartitus.
Pet. quinque obovata, calycem subaequantia, receptaculo inserta.
Stam. hypogyna. Filamenta brevissima. Antherae lineares.
Styl. filiformis, exsertus. Stigmata tria, reflexa.
Bacca globosa, trilocularis.
 — — Herit. stirp. t. 16.
972. *GRANGERIA* Commers. Juss. (1789.)
Cal. quinquepartitus, laciniis revolutis.
Pet. quinque unguiculata.
Stam. calycis limbo inserta. Filamenta basi dilatata, subconnata, apice curvata. Antherae subrotundae, biloculares.
Stylus lateralis.
Drupa monopyrena.
Rosaceae. Lam. ill. t. 427.
973. *NITRARIA* L. (1758.)
Cal. quinquefidus, brevissimus, persistens.
Pet. quinque canaliculata, apice fornicata, mucrone inflexo.
Stam. Filamenta hypogyna, basi dilatata. Antherae dorso incumbentes, biloculares, loculis subdiscretis.
Styl. simplex.
Drupa conica, putamine apice [saxariam] dehiscente, monosperma.
 — — Linn. in nov. comm. acad. petrop. 7. t. 10. Gärtn. t. 58.
974. *PEGANUM* Theophr. L. (1737.)
Cal. quinquepartitus, persistens; laciniis linearibus elongatis, basi subdentatis.
Pet. quinque ovali-oblonga, nervosa.
Stam. hypogyna. Filamenta basi arcuata, incrassata. Antherae oblongae, biloculares, basi emarginatae.
Styl. filiformis. Stigma incrassatum, trigonum, apice trifidum.
Caps. trilocularis, trivalvis.
Rutaceae. Schk. t. 127. Gärtn. t. 95.
975. *TALINUM* Adans. (1763.)
Cal. disepalus, deciduus.
Pet. quinque.
Stam. hypogyna. Antherae biloculares.
Styl. filiformis. Stigmata tria.

Caps. unilocularis, trivalvis, polysperma. Placenta centralis. Sem. saepe alata.

Portulacaceae. Bot. mag. 1543. *Gärtn.* t. 128. [Rülingia Ehrh. Anacampseros Sims. Phemeranthus Rafin.]

1976. *LEWISIA* Pursh. (1814.)

Cal. hepta- enneapetalus, coloratus, scariosus.

Pet. numero dupla, lanceolata, sepalis longiora.

Stam. tot quot petala, iisque opposita, hypogyna. Antherae oblongae, erectae.

Styl. filiformis. Stigmata tria, bifida.

Caps. trilocularis, loculis dispermis.

— —

b. *Fructus inferi.*

1977. *PORTULACA* Varr. Plin.

Cal. bipartitus, caducus.

Pet. quatuor, quinque aut sex.

Stam. numero varia, hypogyna. Filamenta basi cohaerentia.

Styl. tri - quinquepartitus, aut Stigmata sessilia tria, quinque, octo.

Caps. unilocularis, medio circumscissa, polysperma. Sem. basi funiculis singulis singula affixa.

Portulacaceae. Schk. t. 130. *Gärtn.* t. 128. Bot. mag. 2885.

1978. *SYMPHYSLIA* Presl. (1827.)

Cal. urceolatus, coriaceus, sexdentatus.

Calyptra apice sexradiata, integra secedens.

Pet. aut squamae duodecim lanceolatae, ciliatae.

Stam. totidem epigyna, petalis basi affixa.

Filamenta nulla. Antherae lineares, conniventes, quadriloculares, apice bicornes et poro gemino dehiscentes.

Styl. cylindricus. Stigma obtusum.

Bacca calyce coronata, sexlocularis, loculis polyspermis.

Ericaceae? Presl. in epist. ad Jacqu. cum ic.

1979. *AGATHOPHYLLUM* Juss. (1789.)

Cal. brevissimus, edentulus.

Pet. sex, calyci inserta, intus villosa.

Stam. alterna sex calyci, alterna petalorum basi inserta.

Filamenta brevia. Antherae subrotundae.

Styl. brevis, simplex.

Drupa exsucca, putamine semisexloculari, evalvi. Sem. unicum grande, sexlobum.

— — [Ravensara Sonner. it. ind. t. 127. *Evodia* *Gärtn.* t. 103.]

1980. *CARALLIA* Roxb. (1814.)

Cal. basi globosus, limbo sex - septemfido.

Pet. sex, septem, orbicularia.

Stam. calyci inserta. Antherae erectae.

Styl. simplex. Stigma trilobum.

Bacca globosa, monosperma, calyce coronata.

Rhizophoraceae R. Br. *Roxb.* coron. 3. t. 211. [Huc, auctore R. Brown, cong. p. 18, *Baraldea Thuarea*, *Barraultia* Steud. * syst.]

981. CHAMAEMELES Lindl. (1822.)

Cal. truncatus, quinquedentatus.

Pet. parva, calyci inserta.

Stam. calyci adnata, exserta. Antherae incumbentes, subreniformes, sulcatae.

Styl. simplex. Stigma obtusum.

Fructus baccatus? dispersmus.

Rosaceae. Lindl. in *Linn. transact.* 13. t. 11.

II. Flores imperfecti.

A. *Fructus superi.*

1982. HUDSONIA L. (1767.)

Cal. corollinus tri- vel quinquepartitus, laciniis duabus dorso appendiculatis, seu laciniis quarta et quinta calycinis cum duabus corollinis confluentibus.

Stam. irregulariter disposita. Antherae subrotundae, biloculares.

Styl. simplex. Stigma obtusum.

Caps. unilocularis, trivalvis, mono - trisperma.

Cisteae. Gärtn. t. 210. [Linnaei, Gärtneri et Nuttallii observationes cum meis consentiunt. Willdenovius hort. berol. t. 15. calycem tripartitum corollamque pentapetalam icone expressit.]

1983. BOCCONIA Plum. (1703.)

Cal. disepalus, caducus.

Stam. hypogyna. Filamenta brevia. Antherae lineares, erectae.

Styl. brevis, bifidus.

Caps. stipitata, siliculaeformis, stylo coronata, monosperma, basi valvis duabus dehiscens, persistente sutura annulari.

Papaveraceae. Bot. mag. 1905. *Humb.* plant. aequin. t. 35. Gärtn. t. 44.

1984. MABEA Aubl. (1775.)

Flores monoeci: pedunculi bracteati, masculi tri - feminei uniflori.

Cal. quinquefidus.

Stam. receptaculo inserta. Filamenta subnulla. Antherae cordatae, biloculares.

Styl. filiformis. Stigmata tria longissima, revoluta.

Caps. cortice crasso tecta, tricocca. Cocci bivalves, monospermi.

Tricoccae. Aubl. guian. t. 334.

1985. *APACTIS* Thunb. (1784.)

Cal. corollinus tetrasepalus. Sepala subrotunda, crenata, opposita latiora.

Styl. simplex.

Fructus ?

1986. *PTEROTUM* Lour. (1790.)

Cal. pentasepalus, coriaceus, persistens.

Stam. Filamenta plana, calyce longiora. Antherae subrotundae, biloculares.

Styl. nullus. Stigma simplex.

Fructus oblongus, acutus, coriaceus, monospermus, latere dehiscens. Sem. ala multifida, basi adnata instructum.

1987. *STIXIS* Lour. (1790.)

Cal. corollinus hexasepalus: sepalis carnosius, revolutis.

Stam. receptaculo inserta, exteriora breviora. Antherae oblongae, stantes.

Styl. brevis, crassus. Stigmata tria, rotunda.

Drupa ovata, monosperma, cortice punctato.

B. *Fructus inferi.*

1988. *ASARUM* Diosc.

Cal. corollinus, germini adnatus, urceolatus vel campanulatus, limbo trifido erecto vel revoluto.

Stam. epigyna. Filamenta subulata. Antherae sub apice filamentorum adnatae, biloculares, loculis discretis.

Styl. crassus, brevis. Stigma sexlobum.

Caps. coriacea, sexlocularis; loculis oligospermis. Sem. hinc substantia glandulosa (arillo?) praedita, qua demta, semilunata conspiciuntur.

Aristolochieae. Schk. t. 127. Gärtn. t. 14.

C. *Perianthia nulla.*

1989. *THOA* Aubl. (1775.)

Flores monoeci, spicati, superne masculi. Spica nodosa; nodi urceolati, proliferi, staminiferi.

Stam. Filamenta brevissima. Antherae exiles. Flores feminei ad basin spicae.

Styl. brevis. Stigmata tria, obtusa.

Caps. monosperma; cortice intus setis urentibus farcto.

— — Aubl. guian. t. 336.

D I G Y N I A.

990. AGRIMONIA Cels. Plin.

Cal. superus, quinquefidus, setis uncinatis obvallatus.

Pet. quinque calyci inserta.

Stam. calycis ori imposita. Filamenta filiformia. Antherae subrotundae, ambitu circinatim dehiscentes.

Styli stigmatibus capitatis.

Sem. duo, calyce indurato inclusa.

Rosaceae. Schk. t. 128. Gärtn. t. 73.

991. CALLICOMA Andr. (1809.)

Flores capitati. Receptaculum globosum villosum. Involucrum tetraphyllum.

Cal. tetrasepalus.

Stam. hypogyna. Filamenta capillaria, elongata. Antherae ovales, dorso affixae, biloculares.

Styli divaricati.

Fructus dispermus.

Cunoniaceae R. Br. Andr. repos. 566.

992. EUCLEA L. fil. (1781.)

Flores dioeci.

Cal. quadri - quinquedentatus.

Cor. quadri - quinquefida.

Stam. brevica, basi corollae inserta. Antherae erectae, tetragonae, sulcatae, apice latere utroque perforatae.

Styli crassiusculi. Stigmata emarginata.

Caps. baccata, mono - trisperma.

— — *Burm. afr. t. 84. f. 1. Jacqu. fragm. 3. t. 1. f. 5.*

993. PLEURANDRA Labill. (1806.)

Cal. pentasepalus.

Pet. quinque.

Stam. unilateralia, receptaculo inserta. Filamenta brevica, raro basi connata. Antherae lineares adnatae, longitudinaliter dehiscentes.

Styli filiformes.

Achenia duo. Sem. arillatum.

Dilleniaceae. Labill. nov. holl. t. 143. 144. Lessert. sel. ic. t. 78 — 82. Reichenb. hort. t. 79.

T R I - T E T R A G Y N I A.

994. RESEDA Plin.

Cal. inferus, quadri - quinque - sexpartitus, patens, persistens.

Pet. numero varia, ut plurimum sex, in plerisque trifida, multifida etiam aut inaequaliter laciniata; suprema nectario squamaeformi imposita.

Stam. Filamenta filiformia. Antherae basi emarginatae, bicrures, reflexae, biloculares, longitudinaliter dehiscetes.

Styli stigmatibus clavatis.

Caps. lignosa, quinquelocularis, loculis dispermis. Sem. margine alata.

Rosaceae. Humb. nov. gen. 6. t. 562. b.

2005. *GLINUS* Löff. (1758.)

Cal. quinquepartitus, intus corollinus.

Pet. quinque aut plura, linearia, fissa.

Stam. hypogyna. Antherae biloculares.

Stigmata quinque.

Caps. quinquelocularis, quinquevalvis, polysperma. Placenta nulla. Funiculi longissimi umbilicales ad angulos loculorum firmati. Sem. reniformia. Embryo spiralis.

Aizoideae. N. L. Burm. ind. t. 36. Gärtn. t. 130.

2006. *ORYGIA* Forsk. (1775.)

Cal. quinquepartitus, persistens.

Pet. plurima, lanceolata.

Stigmata filiformia.

Caps. quinquelocularis, quinquevalvis, polysperma.

Aizoideae.

2007. *MILTUS* Lour. (1790.)

Cal. quinquepartitus, patens, coloratus.

Pet. nulla.

Stam. fundo calycis inserta. Antherae erectae, didymae

Stigmata linearia, reflexa.

Caps. quinque, connatae, monospermae.

Aizoideae.

2008. *BRUNELLIA* Ruiz. et Pav. (1794.)

Cal. quinquepartitus.

Pet. nulla.

Stam. disco crasso perigyno inserta. Antherae cordatae, dorso affixae, biloculares.

Styli quinque.

Folliculi aut cocci bivalves, monospermi.

Terebintheae. Ruiz et Pav. prodr. t. 12. Humb. plant. aequin. t. 59 — 62.

HEXAGYNIA.

2009. *STRATIOTES* Dodon. (1557.)

Flores dioeci.

Cal. tripartitus.

Pet. tria orbiculata.

Stam. receptaculo inserta, exteriora abortientia. Antherae utrinque filamentis adnatae, apice cuspidatae. In corolla ♀ sola filamenta sterilia numerosa.

DODECANDRIA. DODECAGYNIA. POLYGYNIA. 401

Styli sex bipartiti, lateribus stigmatosis.
Caps. infera baccata, hexagona, sexlocularis. Sem. numerosa, ad dissepimenta fixa.

Hydrocharideae. Schk. t. 335. Gärtn. t. 14.

010. CEPHALOTUS Labill. (1806.)

Cal. sexpartitus, coloratus.

Cor. nulla.

Stam. basi calycis inserta. Antherae didymae, dorso glanduliferae.

Styli teretiusculi.

Caps. sex, monospermae.

Rosaceae? Labill. nov. holl. t. 145. R. Brown. ad Flinders it. t. 4.

D O D E C A G Y N I A.

011. SEMPERVIVUM Plin.

Cal. sex - duodecimpartitus.

Pet. sex ad duodecim, lanceolata, basi squamata.

Stam. hypogyna.

Stigmata acuta.

Caps. sex ad duodecim, folliculares, polyspermae.

Sedae. Schk. t. 130. Gärtn. t. 65. Sturm. fl. germ.

P O L Y G Y N I A.

012. CERATOCEPHALUS Mönch. (1794.)

Cal. pentasepalus.

Pet. unguiculata.

Caryopses cornutae, spicatae.

Ranunculeae. Lessert. ic. sel. 1. t. 23.

013. ISOPYRUM L. (1748.)

Cal. pentasepalus, coloratus, deciduus.

Pet. quinque tubulosa, bilabiata, labio exter. bifido.

Styli longitudinaliter stigmatosi.

Caps. folliculares polyspermae.

Ranunculeae. Schk. t. 153. Gärtn. t. 65.

014. LAURELIA Juss. (1809.)

Flöres monoeci.

Cal. campanulatus, lobis septem ad tredecim duplici serie dispositis.

Cor. nulla.

Stam. calycis fundo inserta. Filamenta basi biglandulifera, squamis ternis munita. Antherae laterales, apice valva dehiscentes.

Germina plura stylis coronata, basi squamis stipata.

Caryopses elongatae, stylo viloso coronatae.

Monimieae. [Pavonia Ruiz et Par. prodr. t. 28. Thiga Molin.]

CLASSIS XII. ICOSANDRIA

CONSPECTUS

CLASSIS DUODECIMAE.

MONOGYNIA.

I. Flores perfecti.

A. Corolla tetra-pentapetala.

a. Fructus superi.

Pomaceae. n. 2015 — 2017.

Spiraeaceae. 2018. 2019.

b. Fructus inferi.

Myrtaceae. 2020 — 2033.

Philadelphaeae. 2034.

Granataeae. 2035.

Legum. 2036 — 2039.

B. Corolla hexa-pleiopetala. 2040 — 2047.

II. Flores imperfecti.

A. Fructus superi. 2048 — 2051.

B. Fructus inferi. 2052 — 2054.

DI - PENTAGYNIA.

I. Fructus superi. 2055 — 2061.

II. Fructus inferi. 2062 — 2072.

DECA - POLYGYNIA.

* *Rosaceae.* 2073 — 2082.

** *Calycanthaceae* Lindl. 2083. 2084.

*** *Urticaceae* affines. 2085 — 2088.

MONOGYNIA.

I. Flores perfecti.

A. COROLLA TETRA - PENTAPETALA.

a. *Fructus superi.*

POMACEAE.

2015. *AMYGDALUS* Theophr.

Cal. campanulatus, sursum coloratus, quinquelobus, deciduus.

Pet. quinque, calyci inserta, plerumque oblonga.

Stam. calycis ori inserta: filamenta filiformia. Antherae erectae ovato-subrotundae, vel oblongae, biloculares.

Styl. simplex. Stigma capitatum.

Drupa ovato-subrotunda, pubescens aut tomentosa, sulco hinc longitudinali. Nux compressa, suturis prominulis, reticulata, foraminulosa. Embryo inversus. Cotyledones carnosae.

Tournef. instit. t. 402. Schk. t. 31. Gärtn. t. 93.

2016. *PRUNUS* Varr.

Cal. subcampanulatus, quinquefidus, deciduus.
Pet. quinque, subrotunda, calyci inserta.
Stam. calycis ori inserta: filamenta filiformia. *Antherae*
 erectae, ovato-subrotundae, vel oblongae, biloculares.
Styl. simplex. Stigma capitatum.
Drupa ovato-subrotunda, glabra. Nux compressa, su-
 turis prominulis, laeviuscula. Embryo inversus. Coty-
 ledones carnosae.

Tournef. inst. t. 398. Schk. t. 132. Gärtn. t. 93.

2017. *CHRYSOBALANUS* L. (1737.)

Cal. campanulatus, quinquefidus, marcescens.
Pet. quinque, oblonga, calyci inserta.
Stam. calycis ori inserta: filamenta filiformia. *Antherae*
 ovato-subrotundae, didymae.
Styl. lateralis. Stigma simplex, obtusum.
Drupa succulenta. Nux compressa, quinquesulca, quin-
 quevalvis.

*P. Brown. jam. t. 17. f. 5. Jacqu. amer. t. 94. [Icaco Plum.
 gen. t. 5. ic. t. 158.]*

S P I R A E A C E A E.

2018. *NEILLIA* Don. (1825.)

Cal. campanulatus quinquefidus, laciniis ovatis cuspidatis.
Pet. quinque subrotunda, fauci calycis inserta.
Stam. calyci duplici serie inserta: filamenta filiformia:
Antherae subrotundae, biloculares.
Styl. filiformis. Stigma simplex, obtusum.
Caps. follicularis unilocularis, intus dehiscens, polysperma,
 stylo persistente coronata. Semina sphaerica, testa
 duplici, exteriore crustacea. Albumen carnosum. Em-
 bryo erectus: cotyledones ovaes.

2019. *KUNZIA* * (1817.)

Cal. campanulatus, quinquepartitus.
Pet. quinque obovata, subunguiculata.
Stam. calyci inserta. *Antherae* oblongae, biloculares.
Styl. brevis, stigmatate capitato.
Caps. follicularis monosperma.

[*Tigarea Pursh. flor. Amer. sept. t. 15. Purshia Cand.*]

b. *Fructus inferi.*

M Y R T E A E.

2020. *MYRTUS* Cato.

Cal. tubo ventricosus, limbo quinque-raro quadripartito,
 persistens.
Pet. quinque, raro quatuor, calycis fauci inserta.

Stam. annulo calycis inserta: filamenta subulata, erecta.
Antherae subrotundae.

Styl. simplex, filiformis. Stigma obtusum.

Bacca ovalis vel subglobosa, bi-trilocularis. Sem. plura in quovis loculo, reniformi-incurva, subossca. Embryo teretiusculus, curvatus: cotyledonibus semicylindricis.

Tournef. inst. t. 409. *Gärtn.* t. 38. *Lam. ill.* t. 410.

2021. MYRCIA Cand. (1826.)

Cal. persistens, tubo ovato aut subgloboso; limbo quinquepartito.

Pet. quinque, calycis fauci inserta.

Stam. annulo calycis inserta: filamenta subulata, erecta.
Antherae subrotundae.

Styl. simplex, filiformis. Stigma obtusum.

Bacca unilocularis, submonosperma. Semen subglobosum, testa laevi. Cotyledones foliaceae contortuplicatae.

[*Eugenia Aubl. guian.* t. 196. 197. 199. 200. 201. *Myrtus Kunth. Humb. nov. gen.* 6. t. 544.]

2022. EUGENIA Michel. (1729.)

Cal. persistens, tubo oblongo, limbo quadripartito.

Pet. quatuor, calycis fauci inserta.

Stam. annulo calycis inserta: filamenta subulata, erecta.
Antherae subrotundae.

Styl. simplex, filiformis. Stigma obtusum. Germen biloculare, ovula plura in quovis loculo.

Bacca unilocularis, submonosperma. Sem. grande, laeviusculum. Cotyledones crassissimae, conferruminatae.

Michel. nov. gen. t. 108. *P. Brown. jam.* t. 25. f. 3. [*Greggia Gärttn.* t. 33. *Plinia Plum. L. Guapurium Just. Myrtus Kunth. Humb. nov. gen.* 6. t. 540 — 543.]

2023. CARYOPHYLLUS Val. Cord. (1561.)

Cal. coriaceus, coloratus, persistens, tubo cylindrico, limbo quadripartito.

Pet. quatuor, apice cohaerentia, decidua.

Stam. numerosissima, in quatuor phalanges divisa, vallo quadrangulari infra calycis limbum inserta. Filamenta filiformia. *Antherae* subrotundae, biloculares.

Styl. conicus, stigmate simplici. Germen biloculare, ovulis pluribus in quovis loculo.

Bacca elliptica, monosperma. Cotyledones inaequales, sinuosae, altera alteram amplectente.

Guimpel et Schlechtend. ic. pharmac. t. 72. *Gärtn.* t. 33. *Hook. bot. mag.* 2750. *Labill. nov. Caled.* t. 63.

2024. NELITRIS Gärttn. (1788.)

Cal. tubo obovato, limbo quatuor-quinquedentato.

Pet. quatuor vel quinque, calycis fauci inserta.

Stam. annulo angusto infra calycis limbum inserta. Filamenta filiformia. Antherae subrotundae, biloculares.

Styl. filiformis. Stigma capitatum.

Bacca subglobosa, unilocularis, polysperma. Sem. duplici ordine in orbem disposita, oblonga, subossea. Embryo inversus, levissime curvatus. Cotyledones minutae.

Radix maxima, oblonga.

Gärtn. t. 27. [*Decaspermum* Forst. char. t. 37.]

025. *CAMPOMANESIA* Ruiz. et Pav. (1794.)

Cal. tubo subgloboso, limbo quinquepartito.

Pet. quinque, ovato-subrotunda, calyci inserta.

Stam. calycis fauci inserta. Filamenta subulata. Antherae oblongae, didymae.

Styl. conico-subulatus. Stigma subpektatum.

Bacca globosa, succulenta, subdecemlocularis, loculis pleiospermis. Sem. subreniformia, testa granulosa, resinifera. Embryo arcuatus.

Ruiz. et Pav. prodr. t. 13. *Kunth.* nov. gen. 6. t. 541.

026. *PSIDIUM* L. (1737.)

Cal. campanulatus vel urceolatus, limbo demum quinquefido.

Pet. quinque, ovata, calyci inserta.

Stam. calyci inserta.

Styl. filiformis. Stigma capitatum.

Bacca grandis, succulenta, multilocularis, dissepimentis divis. Sem. numerosa, in pulpa nidulantia, reniformia, testa dura. Embryo arcuatus, teretiusculus.

Lam. in transact. Linn. soc. 11. t. 37. *Raddi* in opusc. scientif. di Bologn. 4. t. 7. Bot. reg. 622. *Aubl.* guian. t. 191. [*Guaiava* Tournef. inst. t. 443. *Gärtn.* t. 38.]

027. *GLAPHYRIA* Jack. (1823.)

Cal. persistens, quinquefidus.

Pet. quinque.

Stam. numerosa.

Styl. simplex.

Bacca quinquelocularis, polysperma. Sem. duplici serie affixa axi.

028. *CALYTHRIX* Labillard. (1806.)

Cal. basi bibracteatus, tubo cylindrico, elongato, limbo quinquefido, laciniis ovatis, singulis seta longa terminatis.

Pet. quinque, calyci inserta.

Stam. calycis fauci inserta: filamenta subulata: antherae subrotundae.

Styl. filiformis.

Caps. monosperma, non dehiscens. Semen subclavato-cylindricum.

Labillard. nov. holl. t. 146. Bot. reg. 409.

2029. ANGOPHORA Cav. (1795.)*Cal.* turbinatus, quinquecostatus, quinqueidentatus.*Pet.* quinque, calyci inserta.*Stam.* numerosa, calycis fauci inserta: filamenta subulata: antherae subrotundae.*Styl.* filiformis.*Caps.* trilocularis, trivalvis, trisperma. Sem. subrotunda plano-convexa, dorso appendiculata. Cotyledones orbiculatae, foliaceae.*Cav.* ic. 4. t. 336. 337. Bot. mag. 1960. [Metrosideros costata Gärtn. t. 34.]**2030. METROSIDEROS Soland. Gärtn. (1788.)***Cal.* ovatus, vel tubulosus, limbo quinquefido.*Pet.* quinque, calyci inserta.*Stam.* numerosa, calycis fauci inserta, longissima: antherae subrotundae.*Styl.* filiformis, stigmatē simplici.*Caps.* bi-trilocularis, loculis polyspermis.*Gärtn.* t. 34. *Lindl.* coll. t. 18.**2031. CALLISTEMON R. Br. (1819.)***Cal.* tubo hemisphaerico, limbo quinquelobo.*Pet.* quinque.*Stam.* numerosa, fauci calycis insidentia, elongata.*Styl.* filiformis, stigmatē capitato.*Caps.* trilocularis polysperma, cum calycis tubo incrassata connata.*Bot. reg.* 393.**2032. LEPTOSPERMUM Forst. (1775.)***Cal.* turbinatus, limbo quinquefido.*Pet.* quinque, calyci inserta.*Stam.* calycis fauci inserta, petalis breviora.*Styl.* filiformis, stigmatē capitato.*Caps.* quadri-quinquelocularis, loculis polyspermis. Sem. exigua, scobiformia.*Forst.* char. 36. *Gärtn.* t. 35. *Labill.* nov. holl. t. 147. 148.*Wendland* sert. hannov. t. 15.**2033. FABRICA Gärtn. (1788.)***Cal.* semisuperus, quinquefidus.*Pet.* quinque exungiculata.*Stam.* calycis fauci inserta: filamenta subulata: antherae subrotundae.*Styl.* simplex, stigmatē capitato.*Caps.* multilocularis. Sem. pauca, alata.*Gärtn.* t. 35. *Bot. mag.* 1304.**PHILADELPHEAE Don.****2034. PHILADELPHUS Rupp. (1718.)***Cal.* turbinatus, limbo quadri-quinquepartito.

Pet. quatuor vel quinque, calyci inserta.
Stam. fauci calycis inserta, petalis breviora.
Styl. quadri- aut quinquefidus, stigmatibus oblongis.
Caps. quadri- quinquelocularis, polysperma: placentae angulares. Sem. numerosa, scobiliformia, testa lata membranacea, apice dentato, cincta.
Schk. t. 131. *Gärtner* t. 35.

G R A N A T E A E.

035. PUNICA Caesalp. (1583.)

Cal. coloratus, campanulatus, carnosus - coriaceus.
Pet. quinque vel plura, calyci inserta.
Stam. numerosa, calyci inserta: filamenta filiformia: antherae subglobosae, rima duplici semicirculari dehiscentes.
Styl. conicus, stigmatibus papilloso.
Fructus grandis, subglobosus, diaphragmate transverso in duas cameras inaequales divisus. Camera superior multilocularis, dissepimentis parietalibus: inferior trilocularis, dissepimentis irregularibus. Sem. baccata. Cotyledones foliaceae spiratim convolutae.
Schk. t. 131. b. *Gärtner* t. 38. *Trew.* ic. pict. t. 71.

L O A S E A E Juss.

036. KLAPROTHIA Kunth. (1823.)

Cal. quadripartitus persistens, laciniis ovatis, aequalibus.
Pet. quatuor, calyci inserta, subrotunda, concava, breviter unguiculata.
Stam. calyci inserta, quinque exteriora sterilia. Filamenta capillaria. Antherae subrotundae biloculares.
Styl. apice quadrifidus.
Fructus baccatus, tri- tetraspermus.
Humb. nov. gen. 6. t. 537.

037. LOASA Adans. (1763.)

Cal. quinquepartitus, persistens.
Pet. quinque concava. Squamae totidem petaloidae cymbaeformes, singulae duo filamenta sterilia intus foventes.
Stam. calyci inserta, in fasciculis subquinis disposita, exteriora sterilia. Filamenta filiformia. Antherae ovales, biloculares.
Styl. apice trifidus.
Caps. turbinata, calyce coronata, unilocularis, apice trivalvis, valvis margine placentiferis. Sem. ovalia reticulata.
Juss. in ann. du mus. 5. t. 1 — 5. *Hook.* exot. flor. t. 83.

2038. *BLUMENBACHIA* Schrad. (1823.)*Cal.* quinquepartitus, marcescens, tubo spiratim striato.*Pet.* quinque compresso - cucullata. Squamae totidem petaloideae cymbaeformes, singulae duo filamenta sterilia foventes.*Stam.* calyci inserta, in fasciculis quinis disposita, exteriora sterilia. Filamenta filiformia. Antherae ovales, biloculares.*Styl.* stigmatate simplici obtuso.*Caps.* in partes decem basi spirales dehiscens: quinque alternae tenuiores dissepimenta sistunt; quinque crassiores, spongiosae. Sem. sulcata.*Reichenb. hort.* 2. t. 124. *Bot. mag.* 2865.2039. *MENTZELIA* Plum. (1703.)*Cal.* quinquepartitus, persistens, tubo cylindrico, sulcato.*Pet.* quinque, calyci inserta.*Stam.* calyci inserta, in fasciculis disposita. Antherae ovales, biloculares.*Styl.* apice trifidus.*Caps.* unilocularis, apice trivalvis. Placentae parietales.*Plum. gen.* t. 6. ic. t. 174. *Bot. mag.* 1760.

B. COROLLA HEXA - PLEIOPETALA.

2040. *PYGEUM* Gärtn. (1788.)*Cal.* cyathiformis sexdentatus, inferus: basi annuliformi persistente.*Pet.* sex, obovata, villosa.*Stam.* calyci inserta, filamenta capillaria: antherae subrotundae.*Styl.* filiformis. Stigma truncatum.*Drupa* sicca, transverse oblonga. Nux subreniformis, medio contracta. Embryo inversus, cotyledonibus crassis.*— Gärtn.* t. 46. *Colebrook. in Linn. transact.* 12. t. 18.2041. *DECUMARIA* Phil. Conr. Fabric. (1759.)*Cal.* campanulatus, limbo 7 — 10 - dentato.*Pet.* septem ad decem, oblonga.*Stam.* numerosa: filamenta filiformia: antherae didymae.*Styl.* cylindricus, stigmatate peltato multiradiato.*Caps.* evalvis, stylo stigmatateque coronata, multilocularis.

Sem. oblonga, membrana obvoluta.

Philadelphaeae. Lam. ill. t. 403. *Bosc. in act. soc. nat. paris.* 1. t. 13.

SALICARIAE.

2042. *LAFOËNSIA* Vandell. (1788.)*Cal.* campanulatus, basi bibracteatus, apice stellato - plicatus, 10 — 12 dentatus.

Pet. decem aut duodecim, summo limbo calycis inserta, oblonga, unguiculata, sublobata.

Stam. calyci affixa, longissime exserta. Filamenta apice inflexa. Antherae dorso affixae, oblongae, interne longitudinaliter dehiscentes.

Styl. filiformis, stigmatē obtuso.

Bacca globosa, polysperma. Sem. fundo affixa, complanata, subalata.

Vandell. in *Römer.* script. t. 7. f. 13. *Cand.* in mém. de la soc. de Genève 3. t. 1. [*Calypsectus Ruiz.* et *Par.* *Kunth.* *Humb.* nov. gen. 6. t. 548.]

2043. *PHYSOCALYMMA* Pohl. (1827.)

Cal. campanulatus, inflatus, octodentatus, basi bibracteatus.

Pet. octo, unguiculata, inter dentes calycis inserta.

Stam. basi calycis affixa, filiformia, exserta. Antherae incumbentes, semilunares.

Styl. filiformis, stigmatē subgloboso.

Caps. ? globosa.

Pohl. plant. brasil. t. 82. 83.

2044. *BARTONIA* Lewis (1804.)

Cal. quinquepartitus, persistens.

Pet. decem, calyci inserta, ovata, unguiculata.

Stam. numerosissima, corolla breviora, calyci inserta. Antherae oblongae, biloculares.

Styl. filiformis, spiratim striatus.

Caps. oblongo - cylindrica, unilocularis, 5 — 7 - valvis; placentae parietales. Sem. numerosa, biserialia, compressa. Embryo erectus.

Loaseae. Bot. mag. 1487.

2045. *SONNERATIA* L. fil. (1781.)

Cal. semiinferus, campanulatus, sexfidus.

Pet. sex patentia.

Stamina basi calycis inserta, numerosa. Filamenta subulata. Antherae globosae.

Styl. filiformis, stigmatē obtuso.

Bacca subglobosa, multilocularis, loculis polyspermis.

Sem. curvata, in pulpa carnosā nidulantia.

Myrteae. [Pagapate *Sonnerat* nov. Guin. t. 10. 11. *Aubletia Gärtn.* t. 78.]

2046. *CACTUS* L. (1737.)

Cal. campanulatus, imbricatus vel aequabilis.

Pet. sex aut plura, calyci adnata, corollam campanulatam vel subrotatam formantia.

Stam. numerosissima, petalis calycique adnata. Filamenta subulata. Antherae ovaes, biloculares.

Styl. stigmatibus vel radiatis, vel rectis, vel conglomeratis.

Bacca unilocularis, polysperma. Placentatio parietalis.
Sem. exalbuminosae.

- a. *Mammillaria Haw.* Plantae aphyllae, globosae, oblongae, ovatae, tuberculis obsitae. Sem. cotyledonibus germinare dicuntur. [Nuttall gen. plant. Amer. bor. 1. p. 296.] *Cand.* in mém. du mus. 17. t. 2 — 5.
- β. *Melocactus Tourn.* Plantae aphyllae, hemisphaericae, oblongae, angulosae, spinosae. Flores in spadice terminali. [Cand. in mém. du mus. 17. t. 6.]
- γ. *Echinocactus Ott.* Plantae aphyllae, hemisphaericae, oblongae, angulosae, spinosae. Flores in summis angulis. [Cand. in mém. du mus. 17. t. 7 — 10.]
- δ. *Cereus C. Bauh.* Plantae aphyllae, elongatae, continuae, plerumque angulatae. [Cand. in mém. du mus. 17. t. 11 — 14.]
- ε. *Opuntia Tourn.* Plantae aphyllae, ramosae, compressae, articulatae: corollae subradiatae. Stigmata recta. [Cand. in mém. du mus. 17. t. 15. 16.]
- ζ. *Peirescía Plum.* Plantae succulentae, foliosae. Corolla subradiata. Stigmata subradiata. [Cand. in mém. du mus. 17. t. 17 — 20.]

2047. RHIPSALIS Gärtn. (1788.)

Cal. tri-sexpartitus brevissimus.

Pet. sex biserialia, marcescentia.

Stam. basi petalorum inserta.

Styli filiformes. Stigmata tria, sex, patentia.

Bacca globularis, calyce coronata. Placenta centralis.

Sem. exalbuminosa.

Gärtn. t. 28. Hook. exot. fl. t. 2. *Cand.* in mém. du mus. 17. t. 21.

II. Flores apetalis.

A. Fructus supéri.

2048. PHOEBROS Lour. (1790.)

Cal. decemfidus, persistens: laciniis ovatis, concavis, alternis maioribus.

Stam. numerosa, basi calycis insidentia, exserta. Filamenta capillaria. Antherae subovatae, minimae.

Styl. crassus.

Bacca ovata, unilocularis, tetrasperma.

— — [Oxyacantha javana Rumph. amboin. auctor. t. 19. f. 3.]

2049. ABATIA Ruiz et Pav. (1794.)

Cal. quadripartitus, interne coloratus.

Stam. numerosa, basi calycis affixa. Filamenta exteriora sterilia, interiora flexuosa elongata. Antherae cordato-ovatae, biloculares.

Styl. simplex, filiformis.

Caps. unilocularis, bivalvis, polysperma. Placentae duae parietales. Sem. oblonga.

Tiliaceae, Ruiz et Pav. prodr. t. 14. Humb. nov. gen. 5. t. 486.

1050. *POLYLEPIS* Ruiz. et Pav. (1794.)

Cal. turbinatus, tri-quadrangularis, superne dentibus spiniformibus instructus, fauce constricta, limbo tri-quadrupartito.

Stam. fauci calycis inserta. Antherae villosae.

Styl. brevis. Stigma multifidum.

Achenium calyce indurato tectum, limbo eius persistente coronatum. Semen appensum.

Rosaceae, Ruiz. et Pav. prodr. t. 15.

1051. *CERCOCARPUS* Kunth. (1823.)

Cal. coloratus, tubo cylindrico persistente, limbo quinquelobo, deciduo.

Stam. limbo calycis affixa. Filamenta subulata. Antherae dorso adnatae, ovaes, biloculares.

Styl. longissimus, sericeus. Stigma simplex, subclavatum.

Fructus membranaceus, monospermus, stylo persistente caudatus, tubo calycis involutus.

Rosaceae, Humb. nov. gen. 6. t. 559.

B. *Fructus inferi.*

1052. *CALYPTRANTHES* Sw. (1788.)

Cal. tubo obovato, limbo integro, ante anthesin circumscisso, calyptram deciduam formans.

Stam. tubo calycis adnata: filamenta capillaria: antherae subrotundae, biloculares.

Styl. simplex.

Bacca unilocularis, oligosperma.

Myrteae. Sw. flor. ind. occid. t. 15. Hilar. plant. us. brasil. 3. t. 14. [Suzygium P. Brown. jam. t. 7. f. 2.

Chytraculia Ej. t. 37. f. 2.]

1053. *EUCALYPTUS* Herit. (1788.)

Cal. medio circumscissus, parte superiore operculum, inferiore cupulam sistente. In operculo saepe vestigia corollae adsunt.

Stam. numerosa, annulo cupulae inserta. Filamenta capillaria. Antherae subrotundae.

Styl. simplex.

Caps. quadrilocularis, polysperma. Placentatio parietalis.

Myrteae. Link et Otto ic. t. 45. Labill. iter, 1. t. 13. 20. nov. holl. t. 150 — 154.

1054. *FOETIDIA* Commers. Juss. (1789.)

Cal. tubo tetragono-hemisphaerico; limbo quadripartito, persistente, demum reflexo.

Stam. multiserialia: filamenta capillaria.

412 ICOSANDRIA. DI-PENTAGYNIA.

Styl. filiformis, stigmate quadrifido.

Drupa exsucca, superne plana, inferne tetragona, quadrilocularis, loculis submonospermis.

Myrtaceae. Lam. ill. t. 419.

DI - PENTAGYNIA

I. *Fructus superi.*

2055. *BRAYERA* Kunth. (1824.)

Cal. persistens, tubo turbinato, extus sericeo; limbo duplici, utroque quinquepartito, lobis exterioribus magnis oblongis, interioribus spathulatis brevioribus.

Pet. quinque, squamaeformia, linearia.

Stam. petalis breviora.

Styli exserti, stigmatibus subpeltatis crenato-lobatis.

Caryopses duae in fundo calycis. Sem. pendula.

Rosaceae. Dictionn. class. hist. nat. 5. p. 501. ic.

2056. *WALDSTEINIA* Willd. (1799.)

Cal. campanulatus, decemfidus, laciniis alternis minoribus.

Pet. subrotunda, calyci inter lacinias maiores inserta.

Stam. numerosa, calyci insidentia. Filamenta capillaria.

Antherae versatiles.

Styli subulati.

Caryopses binae aut ternae in fundo calycis.

Rosaceae. Cl. Rich. in Nestl. Potent. t. 1. *Kitaib.* pl. hung. 1. t. 77.

2057. *SPIRAEA* Theophr.

Cal. quinquefidus, persistens.

Pet. quinque, calyci adnata.

Stam. calyci inserta.

Styli tres, quatuor, quinque, simplices.

Capsulae tres aut plures, membranaceae, folliculares, bivalves, pleiospermae. Sem. suturae internae affixa.

Rosaceae. Schk. t. 134. Sturm. fl. germ. Engl. bot. 284. 960. 1468.

2058. *GILLENIA* Mönch. (1794.)

Cal. tubuloso-campanulatus, ore contractus, quinquefidus.

Pet. quinque, lineari-lanceolata, elongata, inaequalia, ori calycis inserta.

Stam. inclusa.

Caps. quinquelocularis, loculis dispermis.

Rosaceae. [Spiraea Bot. mag. 489. Cambessedes in ann. des sc. nat. 1. t. 28]

2059. *KERIA* Cand. (1817.)

Cal. quinquefidus: laciniis ovatis, tribus obtusis, duabus apice calloso-submucronatis.

Pet. orbiculata, calyci inserta.

Stam. calyci inserta. Filamenta filiformia. Antherae ovatae.

Styli subquini filiformes.

Achenia subquina, globosa, glabra, stylis coronata.

Rosaceae. [Corchorus Andr. bot. rep. 587. Bot. mag. 1296.]

060. *SESUVIUM* L. (1758.)

Cal. campanulatus, quinquepartitus, persistens: intus corollinus, superficie corollina interdum producta ultra calycis limites.

Cor. 0.

Stam. tubo calycis adnata.

Styli tres, quinque.

Caps. circumscissa, tri-rarius quinquelocularis. Placenta centralis. Sem. numerosa, curvata.

Aizoideae. Lam. ill. t. 434. Bot. mag. 1701.

061. *AIZOON* L. (1737.)

Cal. quinquepartitus, persistens, intus corollinus.

Cor. 0.

Stam. per fasciculos calyci affixa.

Stigmata crassa, sessilia.

Caps. quinquelocularis, pentagona, quinquevalvis, polysperma.

Aizoideae. Gärtn. t. 76. Lam. ill. t. 437.

II. *Fructus inferi.*

062. *TETRAGONIA* L. (1737.)

Cal. germi adhaerens, intus subcorollinus, plerumque quinquefidus, post anthesin auctus.

Cor. 0.

Stam. numero varia, calyce fere breviora.

Styli brevissimi tres aut plures.

Nux calyce vestita, non dehiscens, tetragona, inermis, alata, vel cornuta, plerumque quadrilocularis, loculis submonospermis.

Aizoideae. Herit. stirp. nov. t. 39. Gärtn. t. 127. 179. Bot. mag. 2362.

063. *MESSEMBRIANTHEMUM* Dillen. (1732.)

Cal. plerumque quinquefidus, persistens, lacinii interdum inaequalibus.

Pet. numerosa, angusta, saepe multiserialia, basi secum et cum calyce concreta.

Stam. numerosa, multiserialia, calyci insidentia, cum petalis basi concreta.

Stigmata quatuor, quinque, aut plura.

Caps. multilocularis, apice stellatim dehiscens, calyci adnata. Sem. reniformi - triangularia.

Aizoideae. Gärtn. t. 126. Dillen. hort. Eltham. t. 179-215. Bot. reg. 260. 358. 493. 494. 582. 863.

POMACEAE.

2064. PYRUS (Pirus Virg. Plin.)

Cal. limbo quinquefido, persistente.

Pet. quinque, subrotunda.

Stam. annulo ad faucem calycis inserta. Filamenta subulata. Antherae biloculares.

Styli subquini (interdum pauciores).

Pomum clausum quinqueloculare, loculis dispermis, dissepimentis cartilagineis. Seminum testa subcartilaginea.

Schk. t. 134. Gärtn. t. 87. [*Sorbus* L. *Schk.* t. 133.]

2065. CYDONIA Tournef. (1700.)

Cal. quinquepartitus persistens, laciniis subfoliaceis.

Pet. subrotunda.

Stam. fauci calycis inserta. Filamenta subulata. Antherae biloculares.

Styli quinque.

Pomum clausum, quinqueloculare, loculis polyspermis, dissepimentis cartilagineis. Seminum testa mucilaginosa.

Tournef. inst. t. 405. *Guimpel et Schlechtend. ic. pharmacop.* t. 17. Bot. reg. 905.

2066. COTONEASTER F. C. Medicus (1793.)

Flores polygami.

Cal. turbinatus, obtuse quinqueidentatus.

Pet. brevia erecta.

Stam. calyci adnata, calycem subaequantia. Antherae biloculares.

Styli 2, 3, staminibus breviores.

Sem. corticata, parietalia, calyce inclusa.

Lindl. in Linn. transact. 13. t. 9. Bot. reg. 1187. [*Mespilus* *Cotoneaster* L. fl. dan. 112.]

2067. MESPILUS Plin.

Cal. quinquepartitus, laciniis foliaceis.

Pet. subrotunda.

Discus magnus, mellifluus.

Styli duo, tres, quinque glabri.

Pomum turbinatum, apertum, quinqueloculare. Seminum testa ossea.

Smith. exot. bot. 1. t. 18. Gärtn. t. 87.

2068. CRATAEGUS Tourn. (1700.)

Cal. tubo urceolato, limbo quinquefido.

Pet. subrotunda.

Stam. annulo calycis glanduloso imposita.

Styli duo, tres, quinque glabri.

Pomum carnosum, subglobosum, clausum, quinqueloculare. Sem. solitaria vel bina in quovis loculo. Putamen osseum.

Fl. dan. 334. [*Mespilus* Schk. t. 133.]

069. *RAPHIOLEPIS* Lindl. (1820.)

Cal. infundibuliformis, coloratus, deciduus, limbo quinquefido.

Pet. ovalia.

Stam. calyci inserta: filamenta filiformia.

Styli duo.

Pomum disco incrassato clausum, putamine chartaceo.

Sem. gibbosa, testa coriacea crassissima.

Bot. reg. 468. 652.

070. *PHOTINIA* Lindl. (1821.)

Cal. quinqueidentatus.

Pet. reflexa.

Stam. annulo calycis imposita.

Styli duo glabri.

Pomum biloculare, calyce carnosio inclusum. Sem. panca, testa cartilaginea.

Lindl. in *Linn. transact.* 13. t. 10. Bot. reg. 491. *Mespilus* glabra Coll. hort. ripul. t. 36.

071. *ERIOBOTRYA* Lindl. (1821.)

Cal. lanatus, obtuse quinqueidentatus.

Pet. barbata.

Stam. calyci adnata eumque aequantia.

Styli quinque filiformes, pilosi.

Pomum clausum, tri-quinqueloculare, loculis monospermis.

[*Mespilus japonica* Thunb. Bot. reg. 365. *M. lanuginosa* Ruiz et Pav. fl. peruv. t. 425.]

072. *OSTEOMELES* Lindl. (1821.)

Cal. persistens, tubo ovali, limbo quinquefido.

Pet. spathulato-oblonga, patentia.

Stam. calycis fauci inserta, erecta.

Styli quinque, inferne barbati, persistentes.

Pomum clausum, lanatum, quinqueloculare, loculis monospermis. Sem. ossea.

Linn. transact. 13. t. 8. ? *Kunth. in Humb. nov. gen.* 6. t. 553. 554.

DECA - POLYGYNIA.

* *Rosaceae.*

073. *COWANIA* Don. (1825.)

Cal. tubo turbinato, limbo quinquefido.

Pet. quinque obovata, fauci calycis inserta.

Stam. duplici serie fauci calycis inserta: filamenta capillaria: antherae biloculares.

Styli quinque aut decem, longissimi, plumosi.

Caryopses quinque aut decem (undecim) stylis persistentibus coronatae.

Don. in *Linn. transact.* 14. t. 22.

2074. DALIBARDA L. (1753.)

Cal. tubo brevi concavo, limbo subquinquefido, laciniiis dentatis.

Pet. quinque.

Stam. fauci calycis inserta: filamenta capillaria: antherae subrotundae vel ovae.

Styli quini vel deni, filiformes, decidui.

Caryopses siccae, subdrupaceae.

Michaux flor. boreali-amer. t. 27. 28. *Richard* in *Nestl.* *Potent.* t. 1. *Bot. mag.* 1567. [*Rubus* *Smith.* ic. Ined. t. 19. 20. 63. *Comaropsis* *Rich.* in *Nestl.* *Potent.* t. 1.]

2075. RUBUS Virg. Plin.

Cal. quinquefidus.

Pet. quinque.

Stam. calyci inserta.

Styli numerosi, sublaterales, capillares.

Baccae plurimae confluentes: receptaculum conicum.

Schk. t. 135. *Gärtn.* t. 73. *Weihe* *Rubi* germ. t. 1 — 4. *Engl. bot.* 715. 716. 826. 827. 1585. 2233. 2442. 2572. *Bot. reg.* 461. 469. 854.

2076. FRAGARIA Brunfels. (1531.)

Cal. decemfidus, laciniiis alternis minoribus.

Pet. quinque.

Stam. basi calycis adnata.

Styli plurimi decidui, sublaterales.

Receptaculum succulentum, post anthesin turgidum, *caryopses* impositas numerosas ferens.

Schk. t. 135. *Gärtn.* t. 73. [*Duchesnea* *Smith.* fructus fert insipidos, flores flavos. *Bot. reg.* 61.]

2077. POTENTILLA Brunfels. (1531.)

Cal. decemfidus, laciniiis alternis minoribus.

Pet. quinque.

Stam. basi calycis inserta.

Styli numerosi laterales.

Receptaculum exsuccum, villosum, pilosum vel glabrum.

Caryopses plurimae.

Schk. t. 136. *Lehmann* hist. *Potentill.* t. 1 — 20. [*Comarum* et *Tormentilla* L.]

2078. GEUM C. Gesn. (1565.)

Cal. decemfidus, laciniiis alternis minoribus.

Pet. quinque.

Stam. basi calycis inserta.

Styli persistentes, post anthesin geniculati, uncinati aut barbati, sublaterales.

Receptaculum exsuccum, caryopses plurimas stylis caudatas gerens.

Schk. t. 137. *Sturm.* fl. germ. *Reichenb.* ic. 1. t. 3.

1079. *SIEVERSIA* Willd. (1811.)

Cal. decemfidus, laciniis alternis minoribus.

Pet. quinque.

Stam. fauci calycis inserta.

Styli persistentes, post anthesin continui.

Receptaculum exsuccum, pilosum, caryopses plurimas stylis caudatas gerens.

Linn. transact. 14. t. 22. *Bot. mag.* 2858. 2863. [*Geum montanum* et réptans *L.* *Sturm.* fl. germ.]

1080. *DRYAS* L. (1735.)

Cal. octopartitus, laciniis aequalibus.

Pet. octo.

Stam. calyci inserta.

Styli capillares, lanati, persistentes.

Rec. exsuccum. Caryopses plurimae, stylis plumosis caudatae.

Gärtn. t. 74. *Fl. dan.* 31. t. 1216. *Engl. bot.* 451.

1081. *ROSA* Varr.

Cal. tubus unceolatus, ampullaceus, turbinatus, persistens; limbus quinquepartitus, laciniis alternis saepe pinnatifidis.

Pet. quinque.

Stam. plurima, calycis collo inserta: filamenta capillaria: antherae biloculares.

Styli plurimi, sublaterales, vel liberi, vel in columnam coacti: stigmata capitata. Calycis tubus baccatus semina plurima subcrustacea, parietibus affixa, partim vero pedicellis partialibus suffulta, stylisque coronata, gerens.

Schk. t. 134. *Gärtn.* t. 73. *Lindley* Ros. t. 1—18.

1082. *ANTHODISCUS* G. F. W. Meyer. (1818.)

Cal. planus, subintegerrimus, persistens.

Pet. quinque oblonga.

Stam. numerosa, annulo calycis inserta. Filamenta capillaria, tortuosa. Antherae minutae, subrotundae, didymae.

Styli plures subulati, apice incurvi.

Bacca corticata, umbilicata, radiatim sulcata.

— — *Meyer* essequieb. p. 193.

** *Calycantheae* Lindl.

1083. *CALYCANTHUS* L. (1753.)

Perianthium multiplici serie imbricatum, coloratum, laciniis interioribus petala mentientibus.

Cor. nulla.

418 ICOSANDRIA. DECA-POLYGYNIA.

Stam. numerosa, collo calycis inserta. Antherae obli-
biloculares, filamentis adnatae, apiculatae.

Styli plures compressi, stigmatibus glandulosis.

Cal. post anthesin baccatus, semina plurima c-
fovens.

Schk. t. 138. Bot. reg. 404.

2084. *MERATIA* Loisel. (1817.)

Perianthium imbricatum: lobi exteriores bractearu
teriores petalorum forma.

Cor. nulla.

Stam. fauci carnosae perianthii inserta: perfecta ex
paucis, antheris bilocularibus adnatis: interiora fili
pilosa.

Styli plures filiformes.

Perianthium post anthesin incrassatum semina pau-
longa cornea gerit. Cotyledones foliaceae, s-
convolutae.

[*Chimonanthus* Lindl. Bot. reg. 451. Calycanthus
Ait. hort. kew. 2. t. 10.]

*** *Urticeis affines.*

2085. *ATHEROSPERMA* Labill. (1806.)

Flores monoeci.

Cal. ♂, involucro diphylo, caduco: campanulatus,
lobus, lobis exterioribus maioribus.

Cor. nulla.

Stam. plurima, basi calycis inserta. Filamenta
capillaria, basi subcoalita. Antherae adnatae, obli-
biloculares.

Cal. ♀, involucro diphylo, caduco: hemisphaericus
tufidus, laciniis brevibus acutis.

Styli capillares, plumosi, persistentes.

Caryopses plurimae, stylis caudatae.

Labillard. nov. holl. t. 224.

2086. *CITROSMA* Ruiz. et Pav. (1794.)

Flores plerumque dioeci.

Cal. hemisphaericus vel urceolatus, quadri- vel octo-
utrique sexui communis.

Stam. plurima, calyci inserta. Filamenta laxa, cor

Styli totidem, exserti, subulati.

Caryopses plurimae baccatae, calyci carnosio immer-

2087. *HEDYCARYA* Forst. (1775.)

Flores dioeci.

Cal. planus, octo - decemfidus.

Stam. filamentis nullis. Antherae plurimae, sessiles,
longae, quadrisulcae, apice barbatae.

Germina pedicellata, subglobosa, stigmatibus sessi-
papillosis terminata.

Nuculae globosae, monospermae.

Forst. char. 64.

2088. EUPOMATIA R. Br. (1814.)

Perianthium superum, limbo integerrimo operculi forma circumscisso.

Stam. numerosa, perianthio inserta; interiora sterilia, petaloidea; exteriora lineari-subulata, ultra antheras adnatas, lineares, biloculares producta.

Stigma sessile, planiusculum, areolis tot, quot loculi germinis.

Bacca polysperma. Sem. albumine carnosio lobato.

— — *R. Brown* app. ad *Flinders* it. t. 2.

CLASSIS XIII. POLYANDRIA.

CONSPECTUS CLASSIS DECIMAE TERTIAE.

MONOGYNIA.

- I. Flores perfecti.
 A. Monopetali. 2089 — 2095.
 B. Tripetali. 2096 — 2098.
 C. Tetrapetali. 2099 — 2114.
 D. Pentapetali. 2115 — 2153.
 E. Hexa - polypetali. 2154 — 2174.
 II. Flores imperfecti. 2175 — 2196.

DIGYNIA.

2197 — 2206.

TRIGYNIA.

2207 — 2225.

TETRAGYNIA.

2226 — 2232.

PENTAGYNIA.

2233 — 2241.

POLYGYNIA.

1. Flores perfecti. 2242 — 2267.
 2. Flores imperfecti. 2268 — 2278.

MONOGYNIA.

I. FLORES PERFECTI.

A. Monopetali.

2089. OMPHALOCARPUS Pal. Beauv. (1805.)

Cal. imbricatus squamis plurimis concavis, obtusis, e villosis.

Cor. monopetala, regularis: tubo intus squamis sex briatis inter stamina munito: limbo sex - aut sept fido, laciniis ovalibus undulatis.

Stam. numerosa, antheris elongatis subulatis rectis.

Styl. filiformis, persistens. Stigma capitatum.

Fructus lignosus, umbilicatus, non dehiscens, multil laris, loculis monospermis. Sem. ossea, lucida, pulpa nidulantia,

Sapotaceae, Pal. Beauv. fl. owar. t. 5.

2090. FOUQUIERA Kunth. (1823.)

Cal. pentasepalus, coloratus, persistens, foliis imbricatis, subcoriaceo - scariosis, exterioribus minoribus.

Cor. monopetala, hypogyna; tubo cylindrico arcuato; limbo quinquepartito subregulari, reflexo.

Stam. hypogyna, exserta. Filamenta inferne villosa, subcohaerentia. Antherae cordatae, biloculares.

Styl. apice trifidus.

Germen superum, sessile, uniloculare: ovula octodecim, placentis tribus parietalibus affixa.

Portulacaeis affinis. Humb. nov. gen. 6. t. 527.

2091. SWARTZIA Schreb. (1789.)

Cal. irregulariter dehiscens in valvas binas, ternas aut quinas.

Petalum hypogynum, laterale, planum, quandoque nullum.

Stam. numero varia, hypogyna; interdum duo aut quatuor maiora sterilia, reliqua basi saepe subconnexa.

Antherae biloculares.

Styl. persistens, rectus aut incurvus.

Legumen pedicellatum, oligospermum.

Leguminosae. Cand. mém. sur les légum. t. 58 — 60.

[Rittera Schreb. Vahl. eclog. t. 9. symb. 2. t. 34. Possira

Aubl. guian. t. 355. Tounatea ib. t. 218.

2092. MARCGRAVIA Plum. (1703.)

Cal. imbricatus, sexpartitus, coriaceus, persistens.

Cor. hypogyna, conica, calyptraeformis, integra, post anthesin basi circumscissa.

Stam. numerosa, membranulae perigynae imposita. Antherae oblongae, longitudinaliter dehiscentes.

Styl. subnullus. Stigma crassum, persistens.

Caps. coriacea, baccata, multivalvis, valvis medio septiferis, dissepimentis basi et apice coalitis, medio distantibus. Sem. minutissima, in pulpa nidulantia.

Marcgraviaeae Juss. Plum. ic. t. 173. P. Brown. Jam.

t. 26. Ann. du mus. 14. t. 25. Hook. exot. fl. t. 160.

2093. ANTHOLOMA Labill. (1800.)

Cal. tetra - raro disepalus, ovatus, caducus.

Cor. ovato - cylindrica, margine crenata.

Stam. numerosissima, disco fungoso alveolato inserta.

Antherae oblongae, apice dehiscentes.

Styl. elongatus. Stigma acutum.

Caps. immatura quadrilocularis.

Marcgraviaeae. Labill. iter 2. t. 41.

2094. ASTERANTHUS Desfont. (1820.)

Cal. monophyllus, basi cupularis, limbo plano multidentato, dentibus obtusis vel acutis.

Cor. regularis, rotata, basi calycis affixa, nervosa, ambitu multifida.

Stam. numerosissima, basi corollae inserta. *Fils* subulata. *Antherae* oblongae, obtusae, biloc erectae.

Stylus columnaris, basi sexcruis. *Stigma* sexlobu
Fructus?

— — Mém. du mus. 6. t. 3.

2095. *ANNESLEA* Wallich. (1829.)

Cal. bibracteatus, quinquepartitus, persistens.

Cor. ventricosa, fauce contracta, limbo quinquefid

Stam. numerosissima, duplici serie disco perigyno i

Antherae lineares, mediis filamentis adnatae.

Styl. teres, corolla longior. *Stigmata* tria su recurva.

Bacca infera, calyce coronata, trilocularis, loculis trispermis. Sem. pendula, arillata.

Ternströmiaceae. Wallich. plant. asiat. rar. 1. t.

B. *Tripetali.*

2096. *EMPEDOCLEA* Hilar. (1825.)

Cal. arcte imbricatus, sepalis subduodemis obtusis

Pet. hypogyna, subunguiculata, spathulata, calyce

Stam. hypogyna. Filamenta filiformia. *Anthera* culares.

Styl. subulatus. *Stigma* capitatum.

Fructus baccatus?

Dilleniaceae. Hilar. fl. brasil. t. 3.

2097. *DOLIOCARPUS* Roland. (1756.)

Cal. pentasepalus persistens, sepalis concavis inaeq

Pet. tria aut quinque, subrotunda.

Stam. receptaculo inserta. Filamenta apice dilata theae adnatae, introrsum dehiscentes.

Styl. subincurvus.

Bacca globosa, disperma.

Dilleniaceae. Roland. in act. holm. 1756. t. 9. [et *Calinea* Aubl. t. 219. 221.]

2098. *TRILIX* L. (1771.)

Cal. trisepalus, persistens, sepalis ovatis acutis pl

Pet. lanceolata, minuta, calyce minora.

Stam. numerosa, hypogyna. Filamenta capillaria theae minutae, subrotundae, didymae.

Styl. cylindricus. *Stigma* simplex.

Bacca subpentagona, quinquelocularis, calyce tecta numerosa, subrotunda, minuta.

— — [? *Ascrea* H. Schott.]

C. *Tetrapetali.*

2099. *PAPAVER* Virgil. Plin.

Cal. disepalus, caducus.

Pet. quatuor, in aestivatione irregulariter plicata.

Stam. hypogyna. Filamenta filiformia. Antherae erectae, biloculares.

Styl. nullus. Stigma peltatum, stellatum, persistens.

Caps. plerumque obovata, medio unilocularis, divisa in ambitu ope septorum abruptorum placentigerorum in plures versus centrum apertos loculos: foraminibus sub stigmate patulis. Sem. numerosissima, reniformia, albugine carnoso, oleoso.

Papavereae. Schk. t. 140. Gärtn. t. 60. Lam. ill. t. 451.

100. ARGEMONE Tourn. (1700.)

Cal. di-trisepalus, sepalis sub apice mucronatis, deciduis.

Pet. quatuor vel sex.

Stam. hypogyna. Antherae oblongae, biloculares.

Styl. nullus. Stigma capitatum, lobatum, persistens.

Caps. unilocularis, valvis apice dehiscens. Placentae filiformes, apice fornicatim ope stigmati persistentis cohaerentes. Sem. sphaerica, scrobiculata.

Papavereae. Tourn. inst. t. 121. Gärtn. t. 60. Schk. t. 141. [Echthrus Lour.]

101. STYLOPHORUM Nuttall. (1818.)

Cal. disepalus, deciduus.

Pet. quatuor.

Stam. hypogyna.

Stylus brevis, demum tortus. Stigma capitatum, quadrilobum.

Caps. unilocularis, subquadrivalvis, valvis revolutis. Placentae filiformes, marginales, apice fornicatim ope stigmati persistentis cohaerentes. Sem. numerosa.

Papavereae. [Papaver cambricum L. Engl. bot. t. 66. Meconopsis Viguier. Cand.]

102. GLAUCIUM Tourn. (1700.)

Cal. disepalus, deciduus.

Pet. quatuor.

Stam. hypogyna.

Styl. nullus. Stigma bilamellatum, persistens.

Caps. siliquaeformis, bi- (raro tri- aut quadri) valvis, valvis ab apice ad basin dehiscentibus. Dissepimentum subspongiosum, foveolatum. Sem. scrobiculata.

Papavereae. Gärtn. t. 115. Engl. bot. 1433. Reichenb. ic. 3. t. 227. [Chelidonium L. Engl. bot. 201. Römeria Medic. Cand.]

103. CHELIDONIUM Diosc.

Cal. disepalus, deciduus.

Pet. quatuor.

Stam. hypogyna.

Styl. brevissimus, subnullus. Stigma bilobum.

Caps. siliquaeformis, torulosa, bivalvis, valvis a basi ad apicem dehiscentibus. Placentae stigmatibus bilobo apice cohaerentes. Sem. strophiliolo glanduloso instructa.

Papaveraceae. Schk. t. 140. Gärtn. t. 115.

2104. CAPPARIS Diosc.

Cal. quadripartitus.

Pet. quatuor.

Stam. subperigyna, numerosa, vel pauciora. Filamenta filiformia. Antherae oblongae, versatiles.

Germen pedicellatum. Stylus nullus. Stigma obtusum, sessile.

Bacca corticosa, stipitata, polysperma: subrotunda vel oblonga; saepe siliquaeformis carnosa, vel etiam dehiscentis. Sem. reniformia, exalbuminosa.

Capparideae. Tourn. inst. t. 139. Schk. t. 139. Jacqu. amer. t. 98—106. Humb. nov. gen. 5. t. 437.

2105. CORYNANDRA Schrad. (1826.)

Cal. tetrasepalus, sepalis ovatis, acutis.

Pet. quatuor, obovata.

Stam. hypogyna. Filamenta clavata, colorata. Antherae oblongae, reflexae.

Styl. stigmatibus simplicibus.

Caps. siliquaeformis, unilocularis, bivalvis, valvis a basi dehiscentibus. Sem. verrucosa.

Capparideae. Reichenb. hort. 2. t. 147.

2106. ACTAEA L. (1737.)

Cal. tetrasepalus.

Pet. quatuor.

Stam. hypogyna. Antherae adnatae, biloculares, introrsum dehiscentes.

Stigma sessile.

Bacca mollis, hinc sulcata, unilocularis. Sem. plana, lateri alteri baccae longitudinaliter affixa.

Ranunculeae. Fl. dan. 498. Gärtn. t. 114. Sturm. fl. germ. [Christophoriaua Tourn. inst. t. 154.]

2107. GRIAS L. (1764.)

Cal. germini adhaerens, cyathiformis, limbo parvo quadrifido persistente.

Pet. quatuor, coriacea.

Stam. numerosa, disco quadrato inserta. Filamenta setacea; basi subconnata. Antherae reniformes, minimae.

Styl. nullus. Stigma crassiusculum, tetragonum, cruciatum, excavatum.

Drupa magna, utrinque attenuata, calyce coronata, octo-sulcata: nucleo oblongo.

Myrteis affinis. Sloane hist. 2. t. 217. [Calophyllum P. Brown.]

2108. CALOPHYLLUM L. (1737.)*Cal.* di-tetrasepalus, coloratus.*Pet.* quatuor, subrotunda, patentia.*Stam.* numerosa, hypogyna. Filamenta filiformia basi subcoalita. Antherae oblongae, erectae.*Styl.* simplex. Stigma concavo-depressum.*Drupa* carnosae, globosae, monosperma: putamine intus fungoso. Sem. exalbuminosum.*Cotyledones* maximae, crassae, carnosae, oleosae.*Guttiferae.* Gärtn. t. 43. Jacqu. amer. t. 165.**2109. STALAGMITES Murr. (1789.)***Cal.* tetrasepalus, caducus, inaequalis.*Pet.* quatuor, obovata, subcoriacea; duo maiora.*Stam.* numerosa, receptaculo carnosae insertae. Filamenta basi subcoalita, clavata, truncata. Antherae didymae.*Styl.* crassus. Stigma quadrilobum.*Bacca* subtrisperma.*Guttiferae.* Murr. in commentat. gotting. 9. p. 175.**2110. MAMMEA L. (1753.)***Cal.* disepalus, coloratus, coriaceus, deciduus.*Pet.* quatuor, coriacea.*Stam.* numerosa, receptaculo insertae. Filam. brevissima, setacea. Antherae oblongae, erectae.*Styl.* cylindricus. Stigma capitatum.*Bacca* maxima, styli parte terminata, unilocularis, tetrasperma.*Guttiferae.* Plum. ic. t. 170.**2111. SPARRMANNIA L. fil. (1781.)***Cal.* tetrasepalus.*Pet.* quatuor.*Stam.* numerosissima, hypogyna: exteriora sterilia, torulosa, colorata. Antherae cordato-ovatae.*Styl.* elongatus, filiformis. Stigma truncatum papillosum.*Caps.* quinquangularis, quinquelocularis, echinata: loculi dispermi. Sem. albumine carnosae.*Tiliaceae.* Bot. mag. 516.**2112. ENTELEA R. Br. (1824.)***Cal.* tetra-pentasepalus.*Pet.* quatuor.*Stam.* omnia perfecta, hypogyna.*Styl.* simplex.*Caps.* echinata, quinquelocularis, polysperma.*Tiliaceae.* Bot. mag. 2480.**2113. HASSELTIA Kunth. (1825.)***Cal.* quadripartitus.*Pet.* quatuor spatulata.*Stam.* perigyna. Filamenta capillaria, persistentia. Antherae biloculares.

Styl. subulatus.

Fructus subglobosus, calyce persistente cinctus, non dehiscent, subdispermus.

Tiliaceae. Humb. nov. gen. 7. t. 651.

2114. ECHINOCARPUS Blum. (1825.)

Cal. quadripartitus, deciduus.

Pet. quatuor, incisa.

Stam. Antherae cuspidatae.

Styl. stigmati simplici.

Caps. lignosa, echinata, quadrivalvis, intus pulpa farinacea farcta.

Bixaeae.

D. Pentapetali.

BIXEAE Kunth.

2115. BIXA Oviéd. (1535.)

Cal. pentasepalus, sepalis aestivatione imbricatis, basi tuberculatis, deciduis.

Pet. obovata.

Stam. hypogyna. Filamenta filiformia. Antherae ovatae.

Styl. elongatus, apice lingulato-compressus.

Caps. ovato-cordata, aculeato-hispida, bivalvis: valvis sulco medio longitudinali praedita, ad quem semina extus pulpa farinacea cincta affixa sunt. Albumen carnosum. Cotyledones foliaceae, flexuosae.

Commel. hort. amstelod. 1. t. 33. Gärtn. t. 61. Bot. mag. 1456.

TILIACEAE.

2116. TILIA Virg. Plin.

Cal. quinquepartitus. Bractea integerrima, colorata, pedunculo adnata.

Pet. quinque, quandoque duplicata.

Stam. hypogyna. Filamenta basi subconnata. Antherae didymae.

Styl. simplex. Stigma capitatum, quinquelobum.

Nux coriacea subcostata mono- aut disperma. Cotyledones foliaceae sinuatae, dentatae.

Schk. t. 141. Ventenat monograph. t. 1 — 5.

2117. BROWNLOWIA Roxb. (1819.)

Cal. campanulatus, quadri-quinquedentatus.

Pet. quinque obliqua.

Stam. hypogyna. Filamenta quinque sterilia inter stamina ac germen.

Styl. simplex.

Caps. tres, bivalves, monospermae.

Roxb. corom. 3. t. 265.

2118. PORPA Blum. (1825.)*Cal.* quinquepartitus.*Pet.* quinque, intus basi tomentosa.*Stam.* annulo membranaceo cincta.*Stigma* tridentatum.*Caps.* octolocularis, loculis monospermis.**2119. DIPTEROCARPUS C. J. Gärtn. (1805.)***Cal.* quinquefidus, laciniis duabus latioribus elongatis, post anthesin in alas longissimas reticulato-venosas ex-crescentibus.*Stam.* Filamenta brevissima. Antherae elongatae subulatae.*Caps.* monosperma, calyce alato vestita. Sem. grande, pyramidatum. Cotyledones maximae, carnosae, foliaceae, intricatissime contortuplicatae.*Gärtn.* suppl. t. 187. *Roxb.* corom. 3. t. 213. *Blum.* flor. Jav. fasc. 7. 8. tab. 1—6.**2120. SHOREA Roxb. Gärtn. (1805.)***Cal.* quinquefidus, post anthesin alis auctus.*Pet.* quinque, lanceolata, patentia.*Stam.* filamenta basi dilatata. Antherae bilobae, mucronatae.*Styl.* persistens.*Caps.* monosperma, stylo coronata, calyce quinquealato vestita. Sem. ovato-acuminatum. Cotyledones crassae, bilobae, inaequales.*Gärtn.* suppl. t. 186. *Roxb.* corom. 3. t. 212.**2121. BERRIA Roxb. (1814.)***Cal.* inaequaliter quinquefidus.*Pet.* subobliqua.*Stam.* basi leviter cohaerentia. Antherae incumbentes bilobae.*Styl.* apice trifidus.*Caps.* trilocularis, trivalvis: valvis singulis dorso alas binas horizontales gerentibus. Sem. bina in quovis loculo, globosa hispida.*Roxb.* corom. 3. t. 264.**2122. COLUMBIA Pers. (1807.)***Cal.* pentasepalus, intus coloratus, deciduus.*Pet.* quinque, intus ad basin squama aucta.*Recept.* pentagonum, stipitifforme.*Stam.* numerosa.*Styl.* staminibus duplo longior.*Fructus* globosus, quadrilocularis, quadrialatus, alis duplicatis, loculis monospermis.*[Colona Cav. ic. 4. t. 370.]***2123. GREWIA L. (1737.)***Cal.* pentasepalus, coriaceus, intus coloratus.*Pet.* intus ad basin squama aut glandula praedita.

Recept. stipitiforme.

Stam. numerosa. Antherae subrotundae.

Styl. simplex. Stigma quadrilobum.

Drupa quadriloba, tetra- aut dipyrenā. Nuces biloculares, dispermae. Sem. albuminosum. Cotyled. foliaceae, planae.

Gärtn. t. 106. *Juss.* in ann. du mus. 4. t. 47 — 51. [*Microcos* et *Mallocoeca* *Forst.* *Chadara* *Forst.*]

2124. DIPLOPHRACTUM Desf. (1819.)

Cal. pentasepalus, tomentosus.

Pet. quinque, basi squama aucta.

Stam. numerosa.

Styl. simplex. Stigmata quinque aggregata.

Caps. globosa, non dehiscens, quinquealata, decemlocularis, loculis per septa transversa divisis, dispermis. Sem. parietibus affixa.

Mém. du mus. 5. t. 1.

2125. MUNTINGIA Plum. (1703.)

Cal. quinquepartitus, deciduus.

Pet. quinque, subrotunda, calyci inserta.

Stam. hypogyna. Filamenta brevissima, capillaria. Antherae subrotundae.

Styl. nullus. Stigma capitato-radiatum, persistens.

Bacca globosa, quinque-octolocularis. Sem. numerosissima, minuta, globosa, albuminosa, in pulpa nidulantia.

Jacqu. amer. t. 107. *Gärtn.* t. 59.

2126. AUBLETIA Schreb. (1789.)

Cal. quinquepartitus, intus coloratus, deciduus.

Pet. quinque, subunguiculata, calyce minora.

Stam. brevissima. Antherae ovato-oblongae, extus gibbae, apice in lobulos binos foliaceos abeuntes: exteriores steriles.

Styl. elongatus, apice dilatatus in stigma infundibuliforme dentatum.

Caps. globosa, depressa, coriacea, tuberculis setisque obsita, multilocularis. Sem. minuta, subglobosa, in pulpa nidulantia.

[*Apeiba Aubl. guian.* t. 213 — 216. *Gärtn.* t. 121.]

2127. CORCHORUS Lobel. (1576.)

Cal. pentasepalus, deciduus.

Pet. quinque.

Stam. hypogyna. Filamenta capillaria; corolla breviora. Antherae parvae.

Styl. subnullus. Stigmata duo, tria, quinque.

Caps. vel subrotundo-ovata, vel oblonga, vel elongata, siliquaeformis, quinquelocularis, quinquevalvis, valvis

medio septiferis, apice interdum cornutis. Sem. biserialia, angulata.

Gärtn. t. 64. 129. Plum. ic. t. 103. 104. Jacqu. hort. vindob. 1. t. 85. 2. t. 173. 3. t. 57. 59. Humb. nov. gen. 5. t. 487.

C I S T E A E.

2128. CISTUS Theophr.

Cal. pentasepalus, sepalis subinaequalibus.

Pet. quinque hypogyna, caduca.

Stam. hypogyna, erecta. Filamenta capillaria. Antherae subrotundae.

Styl. filiformis. Stigma capitatum.

Caps. calyce oblecta, quinquelocularis, quinquevalvis, valvis medio septiferis. Sem. ovato-angulata. Embryo filiformis, spiralis.

Tourn. instit. t. 136. Gärtn. t. 76.

2129. HELIANTHEMUM Val. Cord. (1565.)

Cal. trisepalus, bibracteatus, seu pentasepalus, sepalis duobus exterioribus accessorius minoribus:

Pet. quinque.

Stam. hypogyna. Filamenta capillaria. Antherae biloculares.

Stylus simplex, rectus vel obliquus. Stigma capitatum.

Caps. unilocularis, trivalvis: valvis medio placentiferis.

Sem. angulata. Embryo uncinato-inflexus.

Tourn. inst. t. 128. Schk. t. 145.

T E R N S T R Ö M I E A E Mirb.

2130. TERNSTRÖMIA L. fil. (1781.)

Cal. pentasepalus, bibracteatus, persistens.

Pet. quinque aut sex, sepalis opposita.

Stam. duplici serie basi corollae inserta. Filamenta filiformia. Antherae lineares, erectae, elongatae.

Styl. cylindricus. Stigma capitatum.

Bacca exsucca, bilocularis, loculis subtetraspermis. Placenta centralis. Sem. angulata, hinc gibba inde plana.

Humb. nov. gen. 5. t. 463. [Taouabo Aubl. guian. t. 227. 228.]

2131. CLEYERA Thunb. (1784.)

Cal. quinquepartitus, bibracteatus, coriaceus, persistens.

Pet. quinque, sepalis alterna.

Stam. hypogyna, corolla breviora. Filamenta capillaria, inaequalia. Antherae didymae, subulatae, hirtae.

Styl. filiformis. Stigmata duo.

Bacca exsucca, calyce cincta, bilocularis. Sem. reniformia, albuminosa. Embryo curvatus.

[Mukoki Kämpfer amoen. p. 874. ic. Banks. ic. Kämpf. t. 33.]

2132. FREZIERA Sw. (1800.)*Cal.* pentasepalus, bibracteatus, persistens.*Pet.* quinque, ovato-subrotunda.*Stam.* hypogyna, subfasciculata. Filamenta filiformia.

Antherae ovales, adnatae.

Styl. filiformis, apice trifidus.*Bacca* exsucca, trilocularis, polysperma. Sem. oblonga, compressiuscula.*Sw.* ind. acid. t. 19. *Humb.* plant. aequinoct. t. 5 — 9. nov. gen. 5. t. 463. f. 2. [*Eroteum Soland.* *Sw.* prodr.]**2133. CARAIPA Aubl. (1775.)***Cal.* quinquepartitus, persistens.*Pet.* quinque, inaequilatera.*Stam.* hypogyna: filamenta filiformia, persistentia, basi interdum subcohaerentia. Antherae vel erectae lineares, vel ovatae, incumbentes, apice glandula notatae.*Styl.* simplex. Stigma capitato-trilobum.*Caps.* ovato-trigona, calyce staminibusque persistentibus cincta, trilocularis, trivalvis, valvis lignosis. Placentae centralis trigono-alata. Sem. compressa.*Aubl.* guian. t. 223. *Mart.* nov. gen. t. 64.**2134. GODOYA Ruiz. et Pav. (1794.)***Cal.* polysepalus, imbricatus, coloratus, deciduus.*Pet.* quinque.*Stam.* hypogyna, exteriora sterilia: interiora perfecta brevissima. Antherae adnatae, elongatae, apice poro gemino dehiscentes.*Styl.* brevis: stigma obsoletum.*Caps.* lignosa, tri-pentagona, tri-quinquelocularis: valvis introflexis septiferis. Placentae centrales lineares.

Sem. biserialia, alata.

Mart. nov. gen. t. 74.**2135. COCHLOSPERMUM Kunth. (1822.)***Cal.* pentasepalus persistens, sepalis duobus exterioribus minoribus.*Pet.* quinque.*Stam.* hypogyna. Filamenta filiformia. Antherae basi affixae, elongatae, quadriloculares, apice poro gemino dehiscentes.*Styl.* simplex.*Caps.* quinquelocularis, prismatica, coriacea, polysperma.

Sem. reniformia, lanata.

[*Wittelsbachia Mart.* nov. gen. t. 55.]**ELAEocarpeae Juss.****2136. VALLEA Mutis. L. fil. (1781.)***Cal.* quinquepartitus, coloratus, deciduus.

Pet. quinque, triloba, aequalia.

Stam. hypogyna, biserialia. Filamenta arcuata, pilosa, germen arcte cingentia. Antherae lineares, tetragonae, biloculares, apice poro gemino dehiscentes.

Styl. filiformis, apice quadrifidus.

Caps. ovata, muricata, sublignosa, quadrilocularis, quadrivalvis, loculis dispermis.

Humb. nov. gen. 5. t. 489.

CHLAENACEAE Thuars.

2137. SARCOCHLAENA Thuars. (1806.)

Involucrum uniflorum, carnosum, persistens.

Cal. trisepalus, membranaceus.

Pet. quinque obovata.

Urceolus carnosus stamina ambiens.

Stam. Filamenta filiformia, erecta. Antherae quadrangulares, latere dehiscentes, dorso affixae.

Styl. cylindricus. Stigma trilobum.

Caps. involucreo baccato cincta, trilocularis, non dehiscent. Placenta centralis. Sem. in quovis loculo solitaria vel bina, ovata, albuminosa. Embryo inversus. Cotyledones foliaceae cordatae.

Aubert du Pet. Thouars isles austr. d'Afr. t. 9. 10.

2138. SCHIZOCHLAENA Thuars. (1806.)

Involucrum membranaceum, biflorum, crenatum.

Cal. trisepalus, membranaceus.

Pet. quinque.

Urceolus basi intus staminifer.

Stam. Filamenta apice dilatata. Antherae adnatae, biloculares, loculis discretis.

Styl. simplex. Stigma trilobum, suburceolatum.

Caps. trilocularis, trivalvis, involucreo ampliato, inciso, viscoso cincta. Placenta centralis. Sem. subsolitaria, ovata, cartilaginea, albuminosa. Embryo inversus.

Cotyledones foliaceae, cordatae.

Aubert du Pet. Thouars isles austr. d'Afr. t. 12.

GUTTIFERAE

2139. BONNETIA Schreb. (1789.)

Cal. pentasepalus, coriaceus, subimbricatus, sepalis exterioribus minoribus.

Pet. quinque.

Stam. hypogyna. Filamenta filiformia. Antherae oblongae, biloculares.

Styl. cylindricus. Stigma capitatum subtrilobum.

Caps. conico-prismatica, trilocularis, trivalvis: valvis introflexis septifera. Sem. bifariam imbricata, compressa.

Mart. nov. gen. t. 100. Nov. act. nat. cur. vol. 12. t. 6.
[*Mahurea Aubl.* guian. t. 222. *Marila Sw.* prodr. *Kielmeyera Mart.* nov. gen. t. 68. 69.]

2140. CLUSIA Plum. (1703.)

Flores polygami.

Cal. penta - (tetra - hexa -) sepalus, imbricatus, persistens.

Pet. quinque, (quatuor, sex,) concava.

Stam. hypogyna, corolla breviora, quandoque pauciora.

Antherae apicis lateri adnatae.

Styl. nullus. *Stigma* peltatum, radiatum, persistens.

Caps. unilocularis, subquinquevalvis, valvis introflexis ad placentas triangulares fixis. Sem. numerosa, ovata, pulpa obtecta.

Plum. ic. t. 87. [*Quapoya Aubl.* guian. t. 343. 344. *Xantho Willd.* *Havetia Kunth.* *Humb.* nov. gen. 5. t. 462.]

2141. SINGANA Aubl. (1775.)

Cal. pentasepalus.

Pet. quinque, (tria,) unguiculata, denticulata.

Stam. hypogyna. *Antherae* subrotundae, biloculares.

Styl. staminibus longior, apice incurvus. *Stigma* capitatum, concavum.

Caps. cylindrica, unilocularis, polysperma. *Placentae* tres laterales. Sem. grandia, sibi incumbentia, pulpa involuta.

Aubl. guian. t. 230. [*Sterbeckia Schreb.*]

2142. VATERIA L. (1753.)

Cal. quinquepartitus, persistens.

Pet. quinque, ovata.

Stam. Filamenta brevissima. *Antherae* elongatae.

Styl. simplex. *Stigma* capitatum.

Caps. ovata, coriacea, unilocularis, trivalvis, monosperma.

Roxb. corom. 3. t. 228.

D I L L E N I E A E.

2143. DELIMA L. (1747.)

Cal. pentasepalus, persistens.

Pet. quinque subrotunda.

Stam. numerosa. Filamenta capillaria. *Antherae* subrotundae.

Styl. cylindricus. *Stigma* simplex.

Caps. unilocularis, coriacea, subdisperma. Sem. arillo spongioso laciniato praedita.

N. L. Burm. fl. ind. t. 37. f. 1. *Gärtn.* t. 106.

2144. OTHLIS H. Schott. (1827.)

Cal. pentasepalus, persistens, bibracteatus, sepalis bracteisque imbricatis, concavis.

Pet. quinque.

Stam. hypogyna. Filamenta filiformia. Antherae lineares, acutae, cum filamentis continuæ, biloculares.

Styl. simplex.

Caps. monosperma. Sem. arillatum,

WINTEREAE R. Br.

2145. TASMANNIA R. Br. Cand. (1818.)

Flores dioeci vel polygami.

Cal. disepalus.

Pet. quinque (aut duo).

Stam. hypogyna.

Stigma lateri interiori germinis adnatum.

Bacca polysperma. Sem. lateraliter affixa, biserialia, incurva.

Lessert. ic. sel. 1. t. 84.

AURANTIAE Corr.

2146. AEGLE Corr. (1800.)

Cal. quinquedentatus, deciduus.

Pet. quinque, patentia.

Stam. Filamenta libera. Antherae lineares, elongatae, mucronatae.

Stigma subsessile.

Fructus baccatus multilocularis, polyspermus. Loculi intus carnosos - spongiosi.

Roxb. coron. 2. t. 143. [Crataeva Marmelos L.]

OCHNEAE Cand.

2147. OCHNA L. (1753.)

Cal. pentasepalus, persistens.

Pet. quinque, interdum duplicata.

Stam. disco hypogyno inserta. Filamenta filiformia, persistentia. Antherae lineares, rima duplici longitudinali dehiscentes.

Styl. filiformis, persistens.

Gynobasis carnosa, baccas monospermas quinque articulatis fulciens.

Cand. in ann. du mus. 17. t. 11 — 16.

AFFINITATES DUBIAE.

2148. SARACENIA Tournef. (1700.)

Cal. duplex, persistens: exter. trisepalus, inter. pentasepalus, coloratus, apicibus incurvis.

Pet. quinque, decidua.

Stam. receptaculo inserta. Filamenta brevia, filiformia.
Antherae dorso affixae, oblongae, biloculares, apice
 dehiscentes.

Styl. brevis filiformis. *Stigma* grande, umbraculiforme,
 quinquangulare, stamina obtegens.

Caps. quinquelocularis, quinquevalvis, polysperma: valvis
 medio septiferis. Placenta centralis pentagona.

Tournef. inst. t. 476. *Bot. mag.* 780. 849. 1710. *Smith.*
exot. bot. 1. t. 53. *Hook. exot. fl.* t. 13.

2149. CASSIPOUREA Aubl. (1775.)

Cal. campanulatus quinquefidus.

Pet. quinque, basi attenuata, margine lacero-fimbriata.

Stam. receptaculo inserta. *Antherae* oblongae erectae.

Styl. persistens. *Stigma* obtusum.

Caps. trilocularis, trivalvis, oligosperma.

Aubl. guian. t. 211. [*Leguotis Sw. fl. ind. occid.* t. 17.
Richaëia Thuars.]

2150. VANTANEA Aubl. (1775.)

Cal. quinquefidus, brevis.

Pet. linearia, recurva, disco perigyno urceolato carnos
 inserta.

Stam. numerosa, ibidem inserta. Filamenta capilaria,
 elongata. *Antherae* subrotundae.

Styl. filiformis, elongatus. *Stigma* obtusum.

Caps. quinquelocularis, pentasperma.

Aubl. guian. t. 229. [*Lemniscia Schreb.*]

2151. TOUROULIA Aubl. (1775.)

Cal. turbinatus, quinquedentatus, basi bisquamatus.

Pet. quinque perigyna, subrotunda.

Stam. numerosa, perigyna. *Antherae* biloculares, locu-
 lis discretis, divaricatis.

Styl. nullus. *Stigma* oblongum, striatum.

Bacca supera, subglobosa, multistriata, septemlocularis,
 heptasperma. Sem. oblonga, villosa.

Aubl. guian. t. 194. [*Robinsonia Scop. Schreb. Willd.*]

2152. ARSIS Lour. (1790.)

Cal. pentasepalus, coloratus, deciduus.

Pet. quinque, oblonga, calyce breviora.

Stam. numerosa, receptaculo elongato inserta. Filamenta
 filiformia. *Antherae* subrotundae, quadriloculares.

Styl. subulatus, stigmatibus acuto.

Bacca stipitata, globosa, monosperma.

2153. DEINBOLLIA Schumach. (1827.)

Flores polygami.

Cal. pentasepalus, deciduus; sepala exteriora minora.

Pet. quinque, concava, sericea, receptaculo inserta.

Discus annularis perigynus.

Stam. receptaculo inserta. Filamenta pubescentia. Antherae oblongae, erectae.

Styl. cylindricus, persistens.

Bacca binae, coriaceae, pubescentes, monospermae.

Sem. grande, pulpa mucosa inclusum.

Schumacher plant. guineens. p. 242.

E. Flores hexa - polypetali.

1154. SANGUINARIA Dill. (1732.)

Cal. disepalus, caducus.

Pet. octo, decem, duodecim, patentia, alterna angustiora.

Stam. corolla breviora. Antherae oblongae.

Styl. nullus. Stigmata duo.

Caps. ventricosa, bivalvis, polysperma.

Papavereae. Bot. mag. 162. Gärtn. t. 184.

1155. PODOPHYLLUM L. (1737.)

Cal. trisepalus, coloratus, deciduus.

Pet. sex, octo, novem, patentia.

Stam. filamentis brevissimis, antheris oblongis, erectis.

Styl. nullus. Stigma obtusum, subpeltatum.

Bacca unilocularis, polysperma.

Podophylleae Cand. Lam. ill. t. 449. Bot. mag. 1819.

1156. BANARA Aubl. (1775.)

Cal. sexpartitus, persistens.

Pet. sex concava, receptaculo inserta.

Stam. disco glanduloso inserta. Filamenta capillaria.

Antherae subrotundae.

Styl. filiformis. Stigma capitatum.

Bacca subexsucca, stylo terminata, polysperma. *Sem.* minima, angulata, striata.

Bixaeae. Aubl. guian. t. 217.

1157. LECYTHIS Löff. (1756.)

Cal. turbinatus, sexlobus, persistens.

Pet. sex, calyci inserta, inaequalia.

Discus annularis, intus staminibus obsitus, hinc dilatatus in ligulam membranaceam papilloso-lamellatam.

Stam. centralia perfecta, lateralialia in ligulam illam eiusque papillas mutata.

Styl. brevis.

Caps. coriaceo-lignosa, demum circumscissa, operculata, quadri- aut sexlocularis. Placenta centralis. *Sem.* pauca, ovato-oblonga, membrana carnea tecta. Cotyledones crassae, arcte conferruminatae.

Myrteae. Jacqu. amer. f. 109. Aubl. guian. t. 283-289.

Poit. in mém. du mus. 13. t. 2. 3.

1158. BERTHOLLETIA Bonpl. (1808.)

Cal. bipartitus, deciduus.

Pet. sex.

Discus annularis cum staminibus, qualis in *Lecythide*.

Styl. incurvus. Stigma capitatum.

Caps. lignosa, evalvis, intus carnosa, operculo parvo papilloso. Placenta centralis. Sem. triangularia, rugosa.

Myrteae. Humb. plant. aequin. t. 36. Poit. in mém. du mus. 13. t. 4.

2159. *CURATARIA* (Couratari Aubl. 1775.)

Cal. tubo turbinato, limbo sexpartito.

Pet. sex, basi coalita.

Stam. Filamenta concreta in ligulam urceolato - concavam truncatam, intus antheriferam, unilateralem.

Styl. simplex, subulatus.

Fructus oblongus coriaceus, subtrigonus, unilocularis, evalvis. Placenta centralis, columnaris, apice dilatata in operculum convexum. Sem. oblongo - lanceolata, fundo fructus affixa.

Myrteae. Aubl. guian. t. 290. Richard. in ann. des scienc. nat. 1. t. 21.

2160. *ESPERA* Willd. (1801.)

Cal. inferus, quadripartitus, persistens.

Pet. sex, calyce longiora, persistentia.

Stam. Filamenta capillaria, corolla breviora. Antherae subrotundae.

Styl. filiformis, stigmati simplici.

Caps. oblonga, quadri - sexlocularis, totidem alis lanceolatis reticulatis aucta. Sem. solitaria, ovata, compressa, hirsuta.

Tiliaceae. Willd. in neuen Schr. der berl. Ges. naturf. Freunde, 3. p. 449.

2161. *LAGERSTRÖMIA* L. (1753.)

Cal. sexfidus, basi bibracteolatus, lobis sinu vacuo distinctis.

Pet. sex unguiculata, fauci calycis inserta.

Stam. tubo calycis inserta. Filamenta filiformia. Antherae ovales, incumbentes.

Caps. calyce cincta, tri - sexlocularis, tri - sexvalvis. Sem. alata.

Salicariae. Roxb. corom. 1. t. 65. 66. Bot. mag. 405. [Münchhausia L. Adambea Lam.]

2162. *RHODOCHLAENA* Thuars. (1806.)

Cal. trisepalus, membranaceus, viscosus, basi bibracteolatus.

Pet. sex, oblique convoluta, patentia.

Annulus discoideus stamina basi fulciens. Antherae subquadratae, dorso affixae, biloculares.

Styl. stigmati trilobo.

Caps. ? trilocularis.

Chlaenaceae. Aubert du Pet. Thouars, isl. d'Afr. austr. t. 13.

2163. THEA Kämpfer. (1712.)*Cal.* penta - hexasepalus, sepalis subrotundis.*Pet.* sex aut plura, concava, bi - aut triserialia.*Stam.* receptaculo inserta. Filamenta filiformia. Antherae subcordatae, dorso affixae.*Styl.* subtripartitus.*Caps.* tricoeca. Sem. cartilaginea, globoso - angulata.

Cotyledones crassae.

Camellieae. Kämpf. amoen. p. 606. Botan. mag. 998.

Lettson's nat. hist. of the Tea-tree. 1772.

2164. MONODORA Dunal. (1817.)*Cal.* tripartitus.*Pet.* sex, duplici serie, exteriora lanceolata acuta, interiora ovata, breviora, crassiora.*Stam.* Antherae subsessiles, confertissimae.*Styl.* nullus. Stigma sessile.*Bacca* grandis, subglobosa, unilocularis, polysperma.

Sem. oblonga, compressa. Albumen crassum, durum, rimis transversis parallelis incisum. Cotyledones foliaceae.

Anoneae. [Anona Myristica Gärtn. t. 125]**2165. ALANGIUM Lam. (1783.)***Cal.* turbinatus, limbo campanulato, quinque - decedentato.*Pet.* sex, decem, linearia, reflexa.*Stam.* filamenta filiformia. Antherae adnatae, lineares, biloculares.*Discus* carnosus ad basin limbi calycis.*Drupa* ovalis, costata, monosperma. Semen albuminosum.

— — Angolan Rheed. malabar. 4. t. 17. 26.

2166. MAYNA Aubl. (1775.)

Flores dioeci.

Cal. trilobus, concavus.*Pet.* octo, unguiculata, receptaculo inserta.*Stam.* receptaculo conico inserta. Filamenta brevia. Antherae oblongae, tetragonae, apice dehiscentes.*Bacca* grandis, polysperma, externe foliosa.*Magnolieae* Cand. *Bixae* Hilar.**2167. MICRANTHERA Chois. (1823.)**

Flores dioeci.

Cal. tetrasepalus, coloratus: sepala exteriora coriacea, interiora membranacea.*Pet.* octo, decem, inaequalia.*Stam.* numerosissima Filamenta brevia compressa. Antherae terminales, minutae, biloculares, loculis poro-
mentientibus.*Styl.* nullus. Stigma sessile quinquelobum.

Cor. 0.

Stam. numerosa, disco inserta. Antherae subrotundae.

Styl. filiformis, aut obsoletus. Stigma obtusum vel orbiculatum, planiusculum.

Bacca subexsucca, oligosperma; placentatione parietali.

Vahl. symb. 3. t. 64. *Lam. ill.* t. 465. *Bory it.* 2. t. 24.
Bot. reg. 972.

2178. *LAETIA* Löffl. L. (1756.)

Cal. quinquepartitus, coloratus, marcescens.

Cor. 0. (raro pentapetala).

Stam. hypogyna. Antherae ellipticae, biloculares, introrsum dehiscentes.

Styl. brevis. Stigma subcapitatum.

Caps. carnosa, stylo terminata, unilocularis, tri-quinquevalvis, polysperma. Sem. membrana pulposa involuta.

Jacqu. amer. t. 108. 183. f. 60. [*Guidonia P. Brown. jam.* t. 29. *Thamnia ib.* t. 25.]

FLACOURTIEAE Rich.

2179. *FLACOURTIA* Commers. Herit. (1784.)

Flores subdioeci.

Cal. quinque - septempartitus, deciduus.

Cor. 0.

Stam. receptaculo dense superposita. Antherae subrotundae.

Styl. subnullus. Stigma latum, persistens, quinque-novemradiatum, radiis supra sulcatis.

Bacca globosa, multifocularis, loculis dispermis. Sem. ossea.

Herit. stirp. t. 30. *Roxb. corom.* 1. t. 68. 69. 3. t. 222.

2180. *KUHLLIA* Kunth. (1825.)

Cal. subgloboso - campanulatus, sex - octofidus, persistens, laciniis interioribus maioribus.

Cor. 0.

Stam. creberrima, hypogyna. Antherae ellipticae, biloculares, erectae.

Styl. cylindricus. Stigma obtusum.

Bacca exsucca, subglobosa, unilocularis, polysperma.

Placentae tres parietales. Sem. obovata, albuminosa.

Humb. nov. gen. 7. t. 652. 653.

2181. *RYANIA* Vahl. (1796.)

Cal. coloratus, pentasepalus, persistens.

Cor. 0.

Urceolus perigynus corollinus, villosissimus.

Stam. receptaculo inserta. Filamenta subulata. Antherae erectae lanceolatae.

Styl. stamina aequans.

Stigmata quatuor.

Bacca polysperma, suberosa, polysperma. Placentae quinque parietales. Sem. arillo trialato, albuminosa.

Vahl. eclog. t. 9.

2182. *PATRISIA* Rich. (1792.)

Cal. coloratus, pentasepalus, persistens.

Cor. 0.

Urceolus perigynus nullus.

Stam. receptaculo inserta. Antherae lineares, erectae, biloculares.

Styl. apice quadri-quinquefidus, stigmatibus capitatis.

Caps. suberosa, tri-quinquevalvis. Placentae parietales.

D I L L E N I E A E.

2183. *DAVILLA* Vandell. (1788.)

Cal. penta-octosepalus, valde inaequalis; sepalis tri-quadriseptis; duobus maximis, oppositis, concavis; duobus aut tribus intimis coloratis, corollinis, caducis.

Cor. 0., nisi sepala intima sic appellaveris.

Stam. hypogyna, persistentia. Filamenta apice gradatim incrassata. Antherae continuae ellipticae, biloculares.

Styl. apice incrassatus, (interdum duo aut tres). Stigma peltatum, umbilicatum.

Fructus monospermus, sepalis duobus incrassatis, crustaceis, concretis inclusus. Sem. subglobosum, albuminosum.

Vandell. in Römer. script. t. 7. f. 14. Lessert. ic. sel. 1. t. 71. Hilari. flor. brasil. t. 2.

G U T T I F E R A E.

2184. *RHEEDIA* L. (van Rheedea Plum. 1703.)

Cal. corollinus, tetrasepalus, sepalis subrotundis patentibus.

Cor. 0.

Stam. perigyna, quatuor seriebus ex adverso sepalorum disposita. Filamenta breviuscula, filiformia. Antherae subrotundae.

Styl. subnullus. Stigma concaviusculum.

Bacca grandis, unilocularis, subtrisperma. Sem. magna, ovato-oblonga, carnosa, in pulpa nidulantia.

Plum. ic. t. 257.

2185. *MELANORRHOEA* Wall. (1829.)

Cal. penta-hexasepalus, coloratus, persistens. Sepala patentissima, lanceolata, obtusa, venosa.

Stam. disco convexo inserta.

Styl. lateralis. Stigma convexum.

Fructus stipitatus, coriaceus, monospermus. Sem. exalbuminosum. Cotyledones crassae.

Wallich. plant. asiat. rar. 1. t. 11. 12.

Cal. bivalvis, uniflorus: valvae ♂ minimae lanceolatae, aequales, ♀ corollam subaequantes, cartilagineae.

Cor. bivalvis: valvae ♂ cartilagineae, ovatae, obtusae: ♀ membranaceae, acutae.

Stam. circiter quadraginta, fundo corollae inserta. Filamenta capillaria. Antherae lineares.

Styli duo filiformes. Stigmata pilosa.

Caryopsis lanceolata, subfalcata.

Aubl. guian. t. 337. *Pal. Beauv. agrost.* t. 22. f. 11.

D I G Y N I A.

D I L L E N I E A E.

2197. CURATELLA Löffl. L. (1756.)

Cal. tetra-pentasepalus, persistens: sepalis subrotundis, inaequalibus.

Pet. subquaterna, subrotunda, concava.

Stam. receptaculo inserta. Filamenta filiformia. Antherae subrotundae, introrsum dehiscentes.

Styli filiformes. Stigmata capitata.

Caps. duae, bivalves, hirsutae, mono-dispermae. Sem. oblonga, nitida.

Aubl. guian. t. 232. *Vent. sel.* t. 49. *Hilar. plant. us. bras.* t. 24.

2198. HEMISTEMMA Juss. Thuars. (1811.)

Cal. pentasepalus, sepalis concavis extus villosiusculis.

Pet. quinque, duo remotiora.

Stam. unilateralia, exteriora abortientia. Filamenta brevissima. Antherae lineares, biloculares.

Styli duo simplices, divergentes.

Caps. duae, subrotundae, villosae, submonospermae. Sem. albuminosa, membrana cincta.

Lessert. ic. sel. 1. t. 74. 75.

2199. TEMUS Molin. (1782.)

Cal. trifidus.

Pet. duodeviginti linearia longissima.

Stam. petalis breviora. Antherae globosae.

Styli coaliti.

Baccae duae siccae. Sem. arillata.

Magnoliseae.

2200. BAUERA Salisb. (1793.)

Cal. sex-octopartitus, persistens.

Pet. sex, octo aut plura.

Stam. hypogyna. Filamenta filiformia. Antherae ovaes, biloculares.

Styli duo, divergentes.

Caps. didyma, bilocularis, polysperma. Sem. oblonga, tuberculata.

Saxifragae? Bot. mag. 715. *Salisb.* in ann. of bot. 1. t. 10.
Reichenb. hort. 1. t. 77.

2201. *TRICHOSPERMUM* Blum. (1825.)

Cal. pentasepalus, deciduus.

Pet. quinque ovalia.

Stam. disco hypogyno inserta. Antherae didymae.

Styl. 0. Stigmata duo emarginata.

Caps. hirsuta, reniformis, bivalvis, polysperma. Sem. lenticularia, arillata, ciliata, albuminosa. Cotyledones foliaceae, orbiculatae.

Bixae. Blum. bijdrag. tot de flor. van nederlandse Indië, p. 57.

2202. *POTERIUM* Lobel. (1570.)

Flores diclini.

Cal. germini accretus, basi bi- aut trisquamatus, limbo quadripartito.

Cor. 0.

Stam. calycis fauci inserta. Filamenta filiformia. Antherae subrotundae, biloculares.

Styli filiformes. Stigmata penicilliformia.

Drupa exsucca, subangulata, uni-bilocularis, mono-disperma. Sem. inversa.

Rosaceae. Schk. t. 300. *Kit.* hungar. 2. t. 198. *Desfont.* atl. 2. t. 251. [*Pimpinella Gärtn.* t. 32.]

2203. *CLIFFORTIA* Eichrodt in hort. Carolruh. (1733.)

Flores dioeci.

Cal. urceolatus, limbo trifido.

Cor. 0.

Stam. Filamenta capillaria. Antherae oblongae, didymae.

Styli duo. Stigmata elongata, plumosa.

Calycis tubus induratus sistit capsulam bilocularem, dispersam. Sem. erecta. Cotyledones foliaceae.

Rosaceae. Linn. hort. Cliffort. t. 30. 31. 32.

2204. *ABLANIA* Aubl. (1775.)

Cal. quadri-quinquepartitus.

Cor. 0.

Stam. hypogyna. Filamenta capillaria. Antherae parvae subrotundae.

Styli duo, apice bifidi.

Caps. setosa, quadrivalvis, unilocularis, polysperma. Placenta centralis libera. Sem. membrana mucosa obvoluta.

— — Aubl. guian. t. 234. [*Trichocarpus Schreb.*]

2205. *FOTHERGILLIA* L. (1774.)

Cal. campanulatus, truncatus, integer, persistens.

Cor. 0.

Stam. Filamenta superne incrassata. Antherae erectae, quadrangulares, biloculares.

P A L M A E.

2213. *LODOICEA* Commers. Labill. (1807.)

Flores dioeci.

Spadix masculus longissimus, densissime imbricatus squamis carnosis.*Cal.* trisepalus, corolla maior.*Cor.* tripetala.*Stam.* basi coalita. Antherae lineares.*Spadix* femineus, squamis grandibus erosis vestitus. Flores remotiusculi.*Sepala* persistentia ingenter incrassata, crenata.*Stigmata* tria sessilia.*Drupa* ingens trisperma. Putamen bilobum.*Labill.* in ann. du mus. 9. t. 13. *Hook.* in bot. mag. 2734-2738. [*Borassus Sonner.* nov. guin. t. 3—7.]2214. *LATANIA* Commers. Jus. (1789.)

Flores dioeci.

Spadix ramosus, amentis apice digitato-partitis. *Spathae* dense imbricatae.*Cal.* trisepalus.*Cor.* tripetala, calycem excedens..*Stam.* basi in columnam coalita. Antherae oblongae biloculares.*Bacca* tripyrena. Sem. albumine corneo. Embryo verticalis.*Jacqu.* fragm. t. 8. [*Cleophora Gärtn.* t. 120.]2215. *PTYCHOSPERMA* Labill. (1808.)

Flores hermaphroditi, sessiles.

Spadix ramosissimus. *Spatha* multivalvis, caduca.*Cal.* trisepalus.*Cor.* tripetala, calycem superans.*Stam.* receptaculo iuxta ac sepalorum basi inserta. Antherae biloculares, versatiles.*Stigmata* tria.*Bacca* oblonga, mucronata, calyce suffulta, putamine fibroso, monosperma. Sem. quinque-striatum. Embryo basilaris.*Labill.* in mém. de l'instit. 1808. p. 251.2216. *SEAFORTHIA* R. Br. (1810.)

Flores polygamo-androgyni.

Spadix subramosus, spathis pluribus.*Cal.* trisepalus.*Cor.* tripetala.*Stigmata* tria sessilia, obtusa.*Bacca* ovalis. Semen striatum. Albumen rimis crebris radiatis exsculptum. Embryo basilaris.*R. Brown.* prodr. nov. holl. p. 267.

217. IRIARTEA Ruiz. et Pav. (1794.)

Flores polygami monoeci.

Spadix ramosissimus. *Spathae* plures imbricatae.*Cal.* trisepalus.*Cor.* tripetala, calycem superans.*Styli* tres.*Drupa* monosperma. Albumen aequabile. Embryo sub-basilaris.*Ruiz. et Pav. prodr. t. 32. Mart. palm. brasil. t. 33 — 37.*
[*Ceroxylon Humb. plant. aequin. t. 1. 2.*]**218. CARYOTA L. (1737.)**

Flores monoeci.

Spadix ramosissimus. *Spathae* plures.*Cal.* trisepalus.*Cor.* tripetala.*Stigmata* tria, sessilia, conniventia.*Bacca* disperma. Albumen rimis crebris radiatis ex-sculptum. Embryo dorsalis.*Gärtn. t. 7. Ann. du mus. 13. t. 3. f. 29.***219. JUBAEA Kunth. (1815.)**

Flores hermaphroditi, pedicellati.

Spadix ramosus. *Spatha* monophylla.*Cal.* trisepalus, corolla minor.*Cor.* tripetala.*Stam.* basi corollae inserta. *Antherae* lineares.*Styli* tres. *Stigmata* simplicia.*Drupa* obovata, monosperma. Nux apice triforata.*Humb. nov. gen. 1. t. 96.***220. DIPLOTHEMIUM Mart. (1824.)**

Flores androgyni, inter bracteas alveolares sessiles, in quincunces dispositi.

Spadix simplex. *Spatha* duplex.*Cal.* tripartitus.*Cor.* tripetala.*Stigmata* tria sessilia.*Drupa* fibrosa. Putamen monospermum, poris tribus basilaribus. Albumen aequabile.*Mart. palm. brasil. t. 75 — 78.***221. ATTALIA Kunth. (1815.)**

Flores androgyni, sessiles, secundi.

Spadix ramosus, radicalis. *Spatha* monophylla.*Cal.* trisepalus minimus.*Cor.* tripetala, carnosa.*Stam.* pistilli rudimento inserta.*Styli* tres.

Drupa fibrosa, trilocularis, loculis monospermis. Putamen lignosum, sulcatum, basi foraminibus tribus perivium. Albumen solidum.

Humb. nov. gen. 1. t. 95. 96.

2222. GOMUTUS Rumph. (1741.)

Flores monoeci sessiles.

Spadix ramosus. *Spatha* simplex.

Cal. trisepalus.

Cor. tripetala.

Stam. Antherae mucronatae.

Stigmata tria sessilia.

Bacca trisperma. Albumen aequabile. Embryo lateralis.

Rumph. herb. amboin. 1. t. 13.

2223. MANICARIA Gärtn. (1791.)

Flores androgyni, alveolis immersi.

Spadix ramosus. *Spatha* maxima, reticulato-fibrosa, aegerrime dehiscens.

Cal. monophyllus, lacerus.

Cor. tripetala, coriacea.

Stigmata tria sessilia.

Drupa echinata, tricocca. Putamen crustaceum, basi foratum. Embryo basilaris in albuminis aequabilis cavo.

Gärtn. t. 176. [*Pilophora Jacqu. fragm.* t. 35. 36.]

2224. BEGONIA Plum. (1703.)

Flores monoeci.

Cal. 0.

Pet. ♂ quaterna, duo maiora, minora saepe deficiunt. ♀ quinque aut plura, persistentia, supera.

Stam. Filamenta basi saepe iuncta. Antherae oblongo-ovales, biloculares, loculis extrorsum dehiscentibus.

Styli tres, brevissimi. *Stigmata* bipartita, oblonga, obtusa.

Caps. triquetra, alata, trilocularis, polysperma.

Begoniae, Polygoneis affines. *Dryand. in Linn. transact.*

1. t. 14. 15. *Smith. ic. ined.* t. 43. 44. 45. *Herit. stirp.*

t. 46. 47. 48. *Hook. exot. bot.* t. 17. 18. *Bot. mag.* 2900.

2920. *Link et Otto nov. ic.* 13. 19. 25.]

2225. MALLOTUS Lour. (1790.)

Flores dioeci.

Cal. trisepalus, villosus.

Cor. 0.

Stam. receptaculo inserta. Filamenta brevissima. Antherae bilobae.

Styl. 0. *Stigmata* tria, oblonga, reflexa, hirsuta.

Caps. supera, subrotunda, triloba, trilocularis, sexvalvis.

Sem. solitaria, angulata, pedicellata.

T E T R A G Y N I A

2226. *ESCHSCHOLZIA* Chamiss. (1820.)

Receptaculum ampliatus, hypocrateriforme.

Cal. mitraeformis, subinteger, caducus.

Cor. tetrapetala, receptaculo inserta, breviter unguiculata.

Stam. et receptaculo et basi petalorum inserta. Filamenta brevissima. Antherae lineares, continuae, biloculares, loculis extrorsum dehiscentibus.

Styli quatuor, simplices, basi confluentes.

Caps. siliquaeformis, angulata, bivalvis. Placentae parietales. Sem. numerosa.

Papaveraceae. Chamiss. in hor. phys. berol. t. 15. Bot. reg. 1165. Bot. mag. 2887.

2227. *CARYOCAR* Allamand. (1770.)

Cal. quinquepartitus, carnosus.

Cor. ampla, pentapetala, petalis oblongis, concavis, subcarnosis.

Stam. innumera, hypogyna. Filamenta basi coalita, filiformia, corolla longiora. Antherae oblongae, dorso affixae, biloculares.

Styli quatuor, filiformes, elongati.

Drupa quadrilocularis, tetrasperma. Noces subreniformes, durissimae, hinc acutae, sulcatae. Sem. reniformia, exalbuminosa, carnosio-oleosa.

Rhizophoraceae Cand. Hook. in bot. mag. 2727. 2728. [Pekoa Aubl. guian. t. 238. 239. *Rhizobolus* Gärtn. t. 98.]

2228. *PERRA* Mutis. (1784.)

Flores dioeci.

Cal. disepalus, minimus, inaequalis, deciduus.

Cor. monopetala, semigloboso-concava, dependens.

Stam. plurima, receptaculo inserta. Filamenta crassiuscula, biserialia, membranulis multifidis plicatis mixta. Antherae oblongae, tetragonae, erectae.

Stigmata quatuor, laciniis peltatis.

Caps. obovata, subtrigona, trilocularis, trivalvis, valvis bifidis.

— — *Mutis* in act. holm. 1784. t. 8. [Perula Schreb. Willd. *Peridium* H. Schott. in cur. post. p. 410.]

2229. *MARIALVEA* Vand. (1788.)

Cal. disepalus, sepalis concavis, demum reflexis.

Cor. tetrapetala, patens.

Stam. numerosa, disco inserta. Filamenta filiformia. Antherae biloculares.

Stigmata quatuor, obtusa.

Bacca corticata, quadrilocularis, tetrasperma.

Guttiferae. Vand. in Röm. script. lusit. t. 7. f. 16. [Tomvita Aubl. guian. t. 364. Beauharnoisia Ruiz. et Pav. in ann. du mus. 11. t. 9.]

2230. TETRACERA L. (1764.)

Cal. (tetra-) pentasepalus, sepalis coriaceis, exterioribus minoribus.

Cor. (tetra-) pentapetala.

Stam. hypogyna. Filamenta apice dilatata. Antherarum loculi discreti utrinque ad apicem filamentorum.

Stylus vel unicus, stigmate orbiculari, concavo: vel styli plures brevissimi, stigmatibus simplicibus.

Capsulae plures monospermae, stylis terminatae, bivalves.

Sem. albuminosum, arillo laciniato.

Dilleniaeae. Gärtn. t. 69. Lessert. ic. sel. t. 67—70.
[*Euryandra* Forst. char. 41. Tigarea Aubl. guian. t. 350.
351. *Wahlbomia* Thunb. act. holm. 1790. t. 9.]

2231. DRIMYS Forst. (1775.)

Cal. integer, vel bi-tripartitus.

Cor. hexapetala.

Stam. hypogyna. Filamenta brevissima, apice crassiora. Antherae didymae, apice adnatae, loculis subdiscretis lateralibus.

Styli 0. Stigmata subquaterna, plano-depressa.

Baccae quatuor pedicellatae, obovatae, uniloculares, tetra-vel pleiospermae. Sem. angulata.

Magnoliaceae. Forst. char. t. 42. Lam. ill. t. 494. Humb. pl. aequin. 1. t. 58. Lessert. ic. sel. 1. t. 83. [Wintera Murr. Winterana Soland. in Medic. obs. and inquir. 5. t. 1.]

2232. CIMICIFUGA L. (1751.)

Cal. tetrasepalus, deciduus.

Cor. tetrapetala. Petala saepe urceolata, unguiculata.

Stam. hypogyna. Filamenta filiformia. Antherae ovales, biloculares.

Styli simplices recurvi, persistentes.

Capsulae quaternae vel quinae, folliculares, stylis recurvis coronatae, pleiospermae. Sem. squamuloso-echinata.

Ranunculeae. Linn. amoen. 7. t. 6. Gmel. fl. sib. 4. t. 70. Gärtn. t. 140. Bot. mag. 1630. 2069. [Actaea Schk. t. 139. Lessert. ic. sel. 1. t. 66.]

P E N T A G Y N I A.

2233. AQUILEGIA Trag. (1560.)

Cal. pentasepalus, coloratus, deciduus.

Cor. pentapetala. Pet. apice hiantia, basi cornuta.

Stam. hypogyna, exteriora dilatata subpersistencia, abortientia, interiora filiformia. Antherae terminales, ovales, biloculares.

Styli simplices, subulati.

Caps. quinque folliculares, membranaceae, pleiospermae, stylis rostratae. Sem. glabra.

Ranunculeae. Schk. t. 146. Gärtn. t. 118. Lessert ic. sel. 1. t. 47 — 49.

2234. NIGELLA Trag. (1560.)

Cal. pentasepalus, corollinus, deciduus: sepalis unguiculatis.

Petala quinque minora, bilabiata, ungue foveolato nectarifera.

Stam. hypogyna. Filamenta filiformia. Antherae adnatae, biloculares, loculis discretis.

Styli subulati, persistentes.

Capsulae quinque folliculares, concreatæ, sutura dehiscentes, polyspermae. Sem. angulata, reticulato-rugosa.

Ranunculeae. Schk. t. 146. Gärtn. t. 118. Lessert ic. sel. 1. t. 45. 46.

2235. LAPLACEA Kunth. (1821.)

Cal. pentasepalus, coriaceus, persistens, sepalis suborbiculatis, concavis, exterioribus minoribus.

Pet. novem, ovalia, aequalia, patentia.

Stam. numerosissima, basi petalorum inserta. Filamenta subulata. Antherae terminales, didymae, loculis extrorsum dehiscentibus.

Styli subcoaliti.

Caps. pentagona, quinquelocularis, quinquevalvis, loculis oligospermis. Placenta centralis. Sem. oblonga, dorso alata, albuminosa.

Ternströmiace. Humb. nov. gen. 5. t. 461.

2236. GEERIA Blum. (1825.)

Flores dioeci.

Cal. pentasepalus, bibracteatus.

Cor. pentapetala.

Stam. numerosa, basi corollae inserta.

Styli basi coaliti.

Bacca quinquelocularis, polysperma. Placenta centralis.

Sem. biserialia, albuminosa.

Ternströmiace. Blum. bijdr. tot de flor. van nederl. Indië, 3. p. 124.

2237. GYNOCARDIA Roxb. (1819.)

Flores dioeci.

Cal. quinquelobus,

Pet. quinque, basi squamis oblongis ciliatis auctae.

Stam. receptaculo inserta. Eorum loco in ♀ corpuscula pinnatifida.

Styli quinque. Stigmata cordata.

Bacca sicca, polysperma.

Capparideae? Roxb. corom. 3. t. 299.

2238. WORMIA Rottb. (1783.)

Cal. pentasepalus, persistens, sepalis obtusissimis.

Pet. quinque, patentia, obovato - oblonga.

Stam. innumera, hypogyna. Filamenta brevissima. Antherae lineares, elongatae, utroque latere dehiscentes.

Styli filiformes, divergentes. Stigmata capitata.

Capsulae quinque, intus dehiscentes, pleiospermae. Sem. pulpa pellucida obvoluta.

Dillenieceae. Rottb. in nov. act. hafn. 2. t. 3. Lessert ic. sel. 1. t. 82. [*Dillenia Thunb.* in Linn. transact. 1. t. 20. *Roxb.* corom. 1. t. 20. *Colbertia Salisb. Cand. Blum.*

2239. PORCELIA Ruiz et Pav. (1794.)

Cal. tripartitus.

Pet. sex, interiora maiora.

Stam. Filamenta subnulla. Antherae lineares.

Styli subnulli. Stigmata obtusa.

Baccae tres aut sex, magnae, cylindricae, coriaceae, polyspermae, sutura dorsali longitudinali notatae. Sem. oblonga, reniformia, compressa, duplici serie.

Anoneae. Ruiz. et Pav. prodr. fl. per. t. 16.

2240. ROUMEA Poit. (1815.)

Flores dioeci.

Cal. campanulatus, quadri - quinquepartitus.

Pet. 0.

Stam. receptaculo inserta, basi glandula crenata cincta.

Styli persistentes. Stigmata umbilicata.

Bacca stylis coronata, oligosperma. Sem. cartilaginea.

Flacourtiaceae. Mém. du mus. 1. t. 4.

2241. REAUMURIA Hasselqu. L. (1756.)

Cal. quinquepartitus, bracteis minoribus cinctus.

Pet. quinque, ovato - oblonga, basi latiora. Squamae binariae petaloideae oblongae, fimbriato - ciliatae ad basin cuiusvis petali.

Stam. numerosa, hypogyna. Filamenta filiformia. Antherae globosae, didymae.

Styli filiformes.

Caps. quinquelocularis, quinquevalvis, loculis dispermis. Sem. villosissima.

Aizoideae. Salisb. parad. long. t. 18. Jacq. f. ecl. 1. t. 18. Bot. reg. t. 845. Bot. mag. 2057. [*Hypericum Labill.* syr. 2. t. 10.]

P O L Y G Y N I A.

1. Flores perfecti.

2242. SAGITTARIA Lobel. (1591.)

Flores monoeci, dioeci.

Cal. trisepalus.

Cor. tripetala.

Stam. receptaculo inserta. Filamenta filiformia. Antherae ovaes; didymae, loculis discretis, extrorsum dehiscentibus.

Styli numerosi, brevissimi. Stigmata acuta, persistentia.

Caryopses numerosae, durae. Semen exalbuminosum.

Embryo teres, conduplicatus.

Alismaceae Rich. Schk. t. 298. Gärtn. t. 84. Bot. mag. 1661. 1632. 1792.

243. *BRASENIA* Schreb. (1789.)

Cal. trisepalus.

Cor. tripetala, calyce maior.

Stam. receptaculo inserta. Filamenta subulata. Antherae oblongae, erectae.

Styli quinque, decem, apice inflexi, pubescentes. Stigmata obtusa.

Caps. plures, oblongae, compressae, evalves, dispermae.

Sem. albuminosa. Embryo operculatus, inversus.

Hydrocharideae. [Hydropeltis Mx. flor. bor. amer. 1. t. 29. Rich. in ann. du mus. 17. t. 5. f. 22. Bot. mag. 1147.]

244. *LIMNOCHARIS* Bonpl. (1808.)

Cal. trisepalus.

Cor. tripetala, minor.

Stam. receptaculo inserta. Filamenta in ambitu abortientia.

Caps. plures, uniloculares, polyspermae. Sem. scabra, exalbuminosa. Embryo curvatus.

Butomaceae Rich. Humb. pl. aequin. 1. t. 34. Rich. in mém. du mus. 1. t. 19. 20. Bot. mag. 2525. [Damasonium Plum. ic. t. 115.]

245. *NELUMBium* Juss. (1789.)

Cal. tetra - pentasepalus.

Pet. subvicena, multiserialia, oblonga.

Stam. basi germinis inserta. Filamenta filiformia super antheris in appendiculas planas producta. Antherae lineares, elongatae.

Receptaculum obconicum, truncatum, alveolatum. Noces monospermae, stylo stigmatique obtuso coronatae, in alveolis libere sedentes. Embryo crassus intra nucem germinans, viridis.

Nymphaeaceae. Bot. mag. 903. Rich. in ann. du mus. 17. t. 9. f. 49. 50. 57. Turpin in ann. du mus. 7. t. 11. f. 27. [Nymphaea Nelumbo Delil. fl. aeg. t. 61. Cyamus Sm. exot. bot. 1. t. 31. 32.]

246. *BARCLAYA* Wallich. (1827.)

Cal. pentasepalus, hypogynus.

Cor. basi tubulosa, germiini insidens, persistens, limbo octo - decemlobo, lobis imbricatis.

Stam. tubo corollae inserta, eoque inclusa. *Antherae* exappendiculatae.

Styli plures coaliti, fundo tubi radiatim inserti.

Bacca carnosae, globosae, multilocularis, polysperma, corolla persistente coronata. Sem. globosae, setis carnosae obiectae, albuminosa.

Nymphaeaceae. Linn. transact. 15. t. 18.

A N O N E A E.

2247. ROLLINIA Hilar. (1825.)

Cal. trisepalus, brevis, deciduus.

Cor. infra receptaculum inserta, monopetala, globosa, apice sexloba, infra lobos in alas tres obtusas dorsales producta.

Stam. basi receptaculi inserta. Filamenta brevissima, plana, apice incrassata. *Antherae* biloculares, dorso dehiscentes.

Styli coaliti.

Bacca squamosa, polysperma.

Hilar. flor. brasil. t. 5.

2248. GUATTERIA Ruiz. et Pav. (1794.)

Cal. tripartitus, laciniis subcordato-ovatis acutis, patentibus.

Pet. sex, ovata.

Stam. disco germinifero inserta. *Antherae* subsessiles.

Styli simplices.

Baccae plures stipitatae, oblongae, coriaceae, saecae, monospermae.

Ruiz. et Pav. prodr. t. 17. Dunal monogr. Anon. t. 29-33.

Lessert ic. sel. 1. t. 90. [Cananga Aubl. guian. t. 244.

Ej. Aberemoa t. 245. Uvaria Roxb. corom. 1. t. 33-34.]

2249. XYLOPIA L. (1762.)

Cal. tripartitus, vel tri-quinquefidus.

Pet. sex, exteriora maiora.

Stam. disco germinifero inserta.

Styli simplices.

Capsulae baccatae, breviter stipitatae, compressae, coriaceae, hivalves, mono-dispermae.

Dunal monogr. Anon. t. 15-20. Aublet. guian. t. 292.

Gärtn. t. 69. [Xylopicium P. Brown.]

2250. ARTABOTRYS R. Brown. (1820.)

Cal. tripartitus, corolla maior, laciniis lanceolatis.

Pet. sex, alterna subrotunda, alterna lanceolata.

Stam. hypogyna.

Styli brevissimi; simplices.

Baccae utrinque attenuatae, dispermae.

Bot. reg. 423. [Unona Dunal Anon. t. 12. 27.]

2251. UNONA L. fil. (1781.)

Cal. tripartitus vel trilobus.

Pet. sex, tria interiora minora.

Stam. disco germinifero inserta.

Styli simplices.

Baccae oblongae stipitatae, uni- vel pluriloculares, polyspermae. Sem. simplici serie disposita.

Dunal monogr. Anon. t. 21. 23. 24. 25. 26. 28. *Lessert* ic. sel. 1. t. 88. 89. *Roxb.* corom. 3. t. 290. *Vahl.* symb. 2. t. 36. *Labill.* nov. Caled. t. 56. [Melodorum et Desmos Lour.]

2252. UVARIA L. (1747.)

Cal. tripartitus.

Pet. sex, calycem excedentia.

Stam. disco germinifero inserta. Filamenta subnulla. Antherae oblongae, truncatae.

Baccae numerosae, substipitatae, obovato-oblongae, multiloculares, polyspermae.

Dunal monogr. Anon. t. 13. 14. *Roxb.* corom. 1. t. 35. *Gärtn.* t. 114.]

2253. ASIMINA Adans. (1763.)

Cal. tripartitus.

Pet. sex, interiora minora.

Stam. numerosa, disco germinifero inserta. Antherae subsessiles, subcoalitae.

Styli subterni.

Baccae ovatae, carnosae, sessiles, pleiospermae.

Dunal monogr. Anon. t. 9. [Annona L. *Schk.* t. 149.]

2254. ANONA Jos. Acosta (1591.) P. Hermann. (1687.)

Cal. tripartitus.

Pet. sex, interiora minora aut nulla.

Stam. disco germinifero inserta. Antherae plurimae subsessiles, apice capitato.

Styli exiles. Stigmata linearia.

Bacca solitaria sessilis, globosa; cortice muricato, squamoso aut reticulato; intus pulposa, ad ambitum multilocularis, polysperma.

Dunal monogr. Anon. t. 2 — 8. *Lessert* ic. sel. 1. t. 86. 87.

Hook. in bot. mag. 2911. 2912. *Tussac* fl. antill. t. 28.

Aubl. guian. t. 246 — 249.

2255. CADSURA Kämpf. (1712.)

Cal. tripartitus.

Pet. sex.

Stam. disco globoso germinifero inserta. Filamenta subnulla. Antherae oblongae.

Styli subnulli.

Baccae plurimae, coadunatae, dispermae.

Kämpf. amoen. p. 477. ic.

MAGNOLIEAE.

2256. MAGNOLIA Plum. (1703.)

Cal. trisepalus, deciduus.

Pet. novem aut plura, patentia.

Stam. disco hypogyno inserta. Filamenta brevia. Antherae adnatae, oblongae, biloculares.

Styli brevissimi, recurvi. Stigmata villosa.

Strobilus ovatus, tectus *Capsulis* confertis, subrotundis, bivalvibus, extrorsum dehiscentibus. Sem. baccata, albuminosa, integumentis crassissimis, funiculis umbilicibus longissimis pendula.

Schk. t. 148. *Gärtn.* t. 70. *Salisb.* parad. lond. t. 5. 38. 43. 87. [*Liriodendron liliifera* L. et duae species *Lour.*, *Talauma Juss.* *Hilar.* flor. bras. t. 4. *Blum.* fl. Jav. fasc. 19. 20. tab. 9—12.]

2257. LIRIODENDRON L. (1737.)

Cal. trisepalus, bracteatus.

Pet. sex patentia.

Stam. hypogyna. Filamenta subulata. Antherae lineares, erectae.

Styl. 0. Stigmata globosa.

Fructus. Samarae coriaceae, mono-dispermae, in conum collectae. Sem. funiculo umbilicali brevi incurvo pendula.

Schk. t. 147. *Gärtn.* t. 178. *Trew.* ic. Ehret. t. 10.

2258. MICHELIA L. (1747.)

Cal. trisepalus, corollinus, deciduus, bractea spathacea lateraliter dehiscente cinctus.

Pet. sex, aut plura, exteriora maiora.

Stam. disco clavato inserta. Filamenta brevissima. Antherae lineares, adnatae.

Styl. 0. Stigmata reflexa, obtusa.

Caps. plures, baccatae, bivalves, di-pleiospermae, in strobilum oblongum collectae.

Lam. ill. t. 493. *Gärtn.* t. 137. *Lessert* ic. sel. 1. t. 85. *Blum.* fl. Jav. fasc. 17. 18. tab. 1—3. [*Sarcocarpon* et *Sphaerostemma* et *Manglietia* *Blum.* fl. Jav. fasc. 19. 20. tab. 6.]

2259. ILLICIAM L. (1749.)

Cal. tri-hexapetalus, deciduus: sepalis inaequalibus.

Pet. novem aut plura, inaequalia.

Stam. receptaculo inserta. Filamenta brevissima. Antherae erectae, oblongae, emarginatae.

Styli brevissimi. Stigmata lateralia, oblonga.

Caps. plures, in orbem dispositae, lignoso-coriaceae, superne dehiscentes, monospermae. Sem. dura, albuminosa.

Ellis in phil. transact. vol. 60. t. 12. *Vent.* Cels, t. 22. *Gärtn.* t. 69. Bot. mag. 439. [*Skimmia Kämpf.* amoen. p. 881. ic.]

† 2260. AROMADENDRON Blum. (1825.)

Cal. monophyllus spathaceus.

Pet. numerosa, angusta, multiserialia,

Stam. Filamenta subulata. Antherae extrorsum dehiscentes.
Caps. dispermae in strobilum lignosum coalitae.

Blum. fl. Jav. fasc. 19. 20. tab. 7. 8.

DILLENEAE.

2261. DILLENIA L. (1737.)

Cal. pentasepalus, coriaceus, persistens.

Pet. quinque, subcoriacea.

Stam. numerosissima, receptaculo inserta. Antherae oblongae, erectae.

Styl. 0. Stigmata plana, persistentia, in orbem disposita.

Bacca multilocularis, pleiosperma.

Smith. exot. bot. t. 92. 93.

2262. CAPELLIA Blum. (1825.)

Cal. pentasepalus, persistens.

Pet. quinque.

Stam. exteriora breviora.

Styli subulati, divergentes.

Fructus membranacei, intus dehiscentes, polyspermi. Sem. duplici serie margini affixa.

Blum. bijdr. tot de flor. van nederl. Indië, 1. p. 5.

RANUNCULEAE.

2263. RANUNCULUS Plin.

Cal. pentasepalus deciduus: (interdum trisepalus).

Pet. plerumque quinque (nonnunquam decem, raro tria) basi foveola squamaque nectarifera instructa.

Stam. hypogyna. Filamenta filiformia. Antherae oblongae, erectae, biloculares.

Styli breves, persistentes. Stigmata incrassata.

Caryopses plurimae, subovatae, compressiusculae, glabrae vel muricatae, echinatae.

Schk. t. 152. *Gärtn.* t. 74. [*Ficaria Dillen.* Hall. *Mönek.*
Cand. *Hecatonis Lour.* *Casalea Hilar.*]

2264. ADONIS Matthiol. (1559.)

Cal. pentasepalus deciduus.

Pet. quinque, sex aut plura, unguibus nudis.

Stam. hypogyna. Filamenta filiformia. Antherae oblongae, biloculares.

Styli brevissimi, laterales. Stigmata lateralia, papillosa.

Caryopses laeves, spicatae, stylis brevibus coronatae.

Schk. t. 152. *Gärtn.* t. 74.

2265. KNOWLTONIA Salisb. (1796.)

Cal. pentasepalus.

Pet. quinque aut plura, unguibus nudis.

Stam. hypogyna.

Styli decidui.

Caryopsees plurimae baccatae, muticae, receptaculo globoso insertae.

Lessert. ic. sel. 1. t. 19. [Anamenia Vent. malmais. 1. t. 22.]

2266. ATRAGENE L. (1747.)

Cal. tetrasepalus, deciduus.

Pet. dena aut duodena, linearia.

Stam. receptaculo inserta. Filamenta brevissima. Antherae oblongae, acuminatae.

Styli villosi cum stigmatibus persistentes.

Caryopsees plures, stylis villosis caudatae.

Schk. t. 150. Gärtn. t. 74. Roxb. corom. 2. t. 188. [Narvelia Herm. Cand.]

2267. HAMADRYAS Commers. Juss. (1789.)

Flores dioeci.

Cal. penta - hexasepalus.

Pet. dena aut duodena, linearia, acuminata, elongata.

Stam. numerosa, brevia, receptaculo inserta. Filamenta filiformia. Antherae subrotundae, biloculares.

Stigmata sessilia.

Caryopsees plures, ovatae.

Lessert ic. sel. 1. t. 22.

2. Flores imperfecti.

2268. TROLLIUS C. Gesn. (1565.)

Cal. corollinus, polysepalus; sepalis subrotundo - oblongis conniventibus.

Nectaria petaloidea, angusta, basi tubulosa.

Stam. receptaculo inserta. Filamenta filiformia. Antherae oblongae, biloculares,

Styli cylindrici, laterales, persistentes.

Capsulae plures, membranaceae, folliculares, pleiospermae, stylis coronatae. Sem. suturae adnata.

Schk. t. 153. Gärtn. t. 118.

2269. HELLEBORUS Theophr.

Cal. penta - (hexa -) sepalus, subcorollinus, persistens aut marcescens.

Nectaria ad basin sepalorum tubulosa, brevia.

Stam. hypogyna. Filamenta subulata. Antherae erectae, ovaes, biloculares.

Styli cylindrici, sublaterales. Stigmata capitata.

Caps. plures, coriaceae, folliculares, pleiospermae, stylis coronatae. Placenta ligamentosa utrinque ad suturam.

Sem. duplici serie, arillo. angulato elongato.

Schk. t. 154. Gärtn. t. 65. Schmidel ic. t. 6. Desfont. sel. t. 45. Kit. hungar. 2. t. 101. 3. t. 271. Rochel bannat. t. 10.

2270. ERANTHIS Salisb. (1807.)

Involucrum multifidum.

Cal. penta - octosepalus, deciduus.

- Nectaria* ad basin sepalorum tubulosa, brevissima, ore bilabiato.
Stam. numerosa. Filamenta filiformia. Antherae ovaes, biloculares.
Styli brevissimi.
Caps. plures, pedicellatae, folliculares. Sem. globosa, simplici serie.
 [Helleborus hyemalis L. Jacqu. fl. austr. t. 202.]
271. *CORRIS* Salisb. (1807.)
Cal. penta - hexasepalus, deciduus.
Nectaria parva, cucullata.
Stam. hypogyna.
Styli persistentes.
Caps. subsenae aut denae, longe pedicellatae, folliculares, stylis acuminatae, tetra - aut hexaspermae.
 [Helleborus trifolius L. amoen. ac. 2. t. 4. f. 18. Anemone grönlandica O. F. Müll. fl. dan. t. 566.]
272. *CALTHA* C. Gesn. (1565.)
Cal. corollinus pentasepalus.
Nect. 0.
Stam. hypogyna. Filamenta subulata. Antherae oblongae biloculares.
Styl. 0. Stigmata oblonga, lateralia, papillosa.
Caps. plures folliculares, membranaceae, pleiospermae.
 Sem. duplici serie suturae affixa: arillo elongato.
 Schk. t. 154. Sturm. fl. germ. Gärtn. t. 118. Pursh fl. amer. sept. t. 17.
273. *HYDRASTIS* Ellis. L. (1762.)
Cal. trisepalus.
Nect. 0.
Stam. hypogyna. Filamenta linearia.
Styli brevissimi, persistentes. Stigmata latiuscula.
Caryopses baccatae, monospermae, in capitalum aggregatae.
274. *THALICTRUM* Diosc.
Cal. corollinus, tetra - aut pentasepalus, caducus.
Nect. 0.
Stam. hypogyna. Filamenta filiformia, saepe colorata vel clavata, vel apice dilatata. Antherae lineares, apice mucronatae, vel apiculatae, biloculares.
Styli subnulli. Stigmata crassiuscula, papillosa.
Caryopses ovato - oblongae, striatae, vel angulatae.
 Schk. t. 151. Gärtn. t. 74. Lessert ic. sel. 1. t. 6 - 12.
275. *ANEMONE* Diosc.
 Flores involucrati: involucri plerumque remota.
Cal. corollinus pleiosepalus.
Nect. 0.

Stam. hypogyna. Filamenta filiformia, saepe apice dilatata. Antherae oblongae biloculares.

Styli filiformes, breves. Stigmata capitata vel dilatata.

Caryopses nudae.

Sturm. fl. germ. *Lessert* tc. sel. 1. t. 13 — 18. *Engl. bot.* 355. 1062. 1484. [*Hepatica Dill. Cand.*]

2276. PULSATILLA Matthiol. (1559.)

Flores involucriati, involucrio remoto.

Cal. corollinus, penta-hexasepalus, deciduus. Glandulae stipitatae, nectariferae ad basin sepalorum.

Stam. hypogyna. Filamenta filiformia. Antherae ovales, biloculares.

Styli persistentes, villosi.

Caryopses caudatae.

[*Anemone Pulsatilla*, pratensis, patens, vernalis, alpina L. *Schk.* t. 150. *Sturm.* fl. germ.]

2277. CLEMATIS Diosc.

Cal. corollinus, tetra-hexasepalus.

Nect. 0.

Stam. hypogyna. Filamenta apice dilatata. Antherae biloculares.

Styli persistentes, villosi.

Caryopses caudatae.

Schk. t. 151. *Gärtn.* t. 74. *Bot. reg.* 97. 238. 599.

2278. GYROSTEMON Desf. (1820.)

Flores dioeci.

Cal. inferus, patens, subsexlobus.

Pet. 0.

Stam. receptaculo gyris concentricis imposita. Filamenta nulla. Antherae cuneiformes, subquadrilobae, biloculares, longitudinaliter dehiscentes.

Styli plures, crassiusculi, acuti, gyro unico dispositi.

Caps. plures in orbem dispositae, membranaceae, monospermae. Sem. albuminosa.

— — *Desfont.* in mém. du mus. 6. t. 6. 8. 10.





